


FLUCTUATIONS OF RANDOM TILINGS AND DISCRETE BETA-ENSEMBLES

ALICE GUIONNET


CNRS (ÉNS Lyon)


Advances in Mathematics and Theoretical Physics, Roma, 2017

Joint work with A. Borodin, G. Borot, V. Gorin, J.Huang



Consider an hexagon with a hole and take a tiling at random. How does it look ?






Petrov's picture.

When the mesh of the tiling goes to zero, one can see a “frozen” region and a “liquid” region. Limits, fluctuations ?


Tiling of the hexagon



Kenyon's picture.

Cohn, Larsen, Propp 98': When tiling an hexagon, the shape of the tiling converges almost surely as the mesh goes to zero.


General domains



Kenyon-Okounkov's picture.

Cohn-Kenyon-Propp 00' and Kenyon- Okounkov 07': The shape of the tiling (e.g the height function) converges almost surely for a large class of domains.


Fluctuations of the surface



Conjecture. (Kenyon-Okounkov) The recentered height function converges to the **Gaussian Free Field** in the liquid region in general domains.

- ▶ (Kenyon-06') A class of domains with no frozen regions
- ▶ (Borodin–Ferrari-08') Some *infinite* domains with frozen regions
- ▶ (Boutillier-de Tilière-09', Dubedat-11') On the torus
- ▶ (Petrov-12', Bufetov-G.-16') A class of simply-connected polygons
- ▶ (Berestycki-Laslier-Ray-16+) *Flat* domains, some manifolds
- ▶ Borodin-Gorin-G.-16', and Bufetov-Gorin.-17' Polygons with holes — *trapezoid gluings*.


Local fluctuations of the boundary of the liquid region




Ferrari-Spohn 02', Baik-Kriecherbauer-McLaughlin-Miller 03':
appropriately rescaled, a generic point in the boundary of the liquid
region converges to the [Tracy-Widom distribution](#) in the random
tiling of the hexagon, the distribution of the fluctuations of the
largest eigenvalue of the GUE.

G-Huang. -17': This extends to polygonal domains obtained by
trapezoid gluings (on the gluing axis).

Trapezoids gluings




Trapezoids gluings



- ▶ We can glue arbitrary many trapezoids, where we may cut triangles or lines,
- ▶ Always along a **single** vertical axis

What is good about trapezoids?




Fact: The total number of tilings of trapezoid with **fixed** along the border horizontal lozenges $l_N > \dots > l_1$ is proportional to

$$\prod_{i < j} \frac{l_j - l_i}{j - i}$$

Indeed Tilings = Gelfand–Tsetlin patterns, enumerated through combinatorics of Schur polynomials or characters of unitary groups

Distribution of horizontal tiles



$H = 4$ cuts

The distribution of horizontal lozenges $\{\ell_i^h\}$ along the axis of gluing has the form: $\ell_{i+1}^h \geq \ell_i + 1$

$$P_N^{\Theta, w}(\ell) = \frac{1}{Z_N^{\Theta, w}} \prod_{i < j} (\ell_i - \ell_j)^{2\Theta[h(i), h(j)]} \prod_{i=1}^N w(\ell_i)$$

$h(i)$ — number of the **cut**. Θ — symmetric $H \times H$ matrix of 1's, 1/2's, and 0's with 1's on the diagonal.

Discrete β -ensembles ($\beta = 2\theta$)

For configurations ℓ such that $\ell_{i+1}^h - \ell_i^h - \theta_{h,h} \in \mathbb{N}$, it is given by:

$$P_N^{\theta,w}(\ell) = \frac{1}{Z_N} \prod_{1 \leq h \leq h' \leq H} \prod_{\substack{1 \leq i \leq N_h \\ 1 \leq j \leq N'_h, i < j}} I_{\theta_{h,h'}}(\ell_j^{h'}, \ell_i^h) \prod w(\ell_i^h),$$

$$\text{where } I_{\theta}(\ell', \ell) = \frac{\Gamma(\ell' - \ell + 1)\Gamma(\ell' - \ell + \theta)}{\Gamma(\ell' - \ell)\Gamma(\ell' - \ell + 1 - \theta)}$$

Note that $I_{\theta}(\ell', \ell) \simeq |\ell' - \ell|^{2\theta}$ with \simeq if $\theta = 1, 1/2$.

Discrete β -ensembles ($\beta = 2\theta$)

For configurations ℓ such that $\ell_{i+1}^h - \ell_i^h - \theta_{h,h} \in \mathbb{N}$, it is given by:

$$P_N^{\theta,w}(\ell) = \frac{1}{Z_N} \prod_{1 \leq h \leq h' \leq H} \prod_{\substack{1 \leq i \leq N_h \\ 1 \leq j \leq N_{h'}, i < j}} I_{\theta_{h,h'}}(\ell_j^{h'}, \ell_i^h) \prod w(\ell_i^h),$$

$$\text{where } I_{\theta}(\ell', \ell) = \frac{\Gamma(\ell' - \ell + 1)\Gamma(\ell' - \ell + \theta)}{\Gamma(\ell' - \ell)\Gamma(\ell' - \ell + 1 - \theta)}$$

Note that $I_{\theta}(\ell', \ell) \simeq |\ell' - \ell|^{2\theta}$ with \simeq if $\theta = 1, 1/2$.

- ▶ We can study the convergence, global fluctuations of the empirical measures

$$\hat{\mu}_N^h = \frac{1}{N_h} \sum_{i=1}^{N_h} \delta_{\ell_i^h/N}, 1 \leq h \leq H$$

and fluctuations of the extreme particles.

Discrete β -ensembles ($\beta = 2\theta$)

For configurations ℓ such that $\ell_{i+1}^h - \ell_i^h - \theta_{h,h} \in \mathbb{N}$, it is given by:

$$P_N^{\theta,w}(\ell) = \frac{1}{Z_N} \prod_{1 \leq h \leq h' \leq H} \prod_{\substack{1 \leq i \leq N_h \\ 1 \leq j \leq N_{h'}, i < j}} I_{\theta_{h,h'}}(\ell_j^{h'}, \ell_i^h) \prod w(\ell_i^h),$$

$$\text{where } I_{\theta}(\ell', \ell) = \frac{\Gamma(\ell' - \ell + 1)\Gamma(\ell' - \ell + \theta)}{\Gamma(\ell' - \ell)\Gamma(\ell' - \ell + 1 - \theta)}$$

Note that $I_{\theta}(\ell', \ell) \simeq |\ell' - \ell|^{2\theta}$ with \simeq if $\theta = 1, 1/2$.

- ▶ We can study the convergence, global fluctuations of the empirical measures

$$\hat{\mu}_N^h = \frac{1}{N_h} \sum_{i=1}^{N_h} \delta_{\ell_i^h/N}, 1 \leq h \leq H$$

and fluctuations of the extreme particles.

- ▶ Bufetov-Gorin 17': the fluctuations of the surface of the whole tiling follows from the fluctuations on the gluing axis.

Discrete β -ensembles ($\beta = 2\theta$): law of large numbers

Assume $w(x) \simeq e^{-NV(x/M)}$, $(\theta_{h,h'})_{h,h'} \geq 0$, $\theta_{h,h} > 0$.

► Fixed heights : $N_h/N \mapsto \varepsilon_h$, Then

$$\lim_{N \rightarrow \infty} \frac{1}{N_h} \sum_{i=1}^{N_h} \delta_{\ell_i^h/N} \rightarrow \mu_{\varepsilon}^h \quad \text{a.s.},$$

► Random heights : $\sum_h N_h = N$, Then $N_h/N \rightarrow \varepsilon_h^*$ and

$$\lim_{N \rightarrow \infty} \frac{1}{N_h} \sum_{i=1}^{N_h} \delta_{\ell_i^h/N} \rightarrow \mu_{\varepsilon^*}^h \quad \text{a.s.},$$

Indeed,

$$P_N^{\theta,w}(\ell) \simeq e^{-N^2 \mathcal{E}(\frac{1}{N_h} \sum_{i=1}^{N_h} \delta_{\ell_i^h/N}, 1 \leq h \leq H)}$$

where \mathcal{E} has a unique minimizer.

Assumption on the equilibrium measures towards fluctuations

Note that for all h :

$$0 \leq \frac{d\mu_\varepsilon^h}{dx} \leq \theta_{hh}^{-1}$$

We shall assume

- ▶ The liquid region $\{0 < \frac{d\mu_\varepsilon^h}{dx} < \theta_{hh}^{-1}\}$ are connected,
- ▶ The equilibrium measures are **off critical**: at the boundary of the liquid region they behave like a **square root**.
- ▶

$$\frac{w(x)}{w(x-1)} = \frac{\phi_N^+(x)}{\phi_N^-(x)}, \quad \phi_N^\pm \text{ analytic}, \phi_N^\pm = \phi^\pm + \frac{1}{N}\phi_1^\pm + o\left(\frac{1}{N}\right)$$

Rmk: Off-criticality should be generically true.

Global fluctuations: fixed heights

Assume $N_h/N \mapsto \varepsilon_h$,

Theorem (Borodin-Gorin-G 15' Borot-Gorin-G 17')

Then for any analytic functions f_h :

$$\left(\sum_{i=1}^{N_h} (f_h(\ell_i^h/N) - \mathbb{E}[f_h(\ell_i^h/N)]) \right)_h \Rightarrow N(0, \Sigma(f)).$$

Global fluctuations: Random Heights

Assume $\sum N_i = N$. Then [WIP Borot-Gorin-G]



$$\frac{N_i}{N} \rightarrow \varepsilon_i^*$$

▶ The heights are equivalent to discrete Gaussian ':

$$P_N^{\theta,w}(N_h - \mathbb{E}[N_h] = x) \simeq \frac{1}{Z} e^{-\frac{1}{2\sigma}(x)^2}$$

Global fluctuations: Random Heights

Assume $\sum N_i = N$. Then [WIP Borot-Gorin-G]



$$\frac{N_i}{N} \rightarrow \varepsilon_i^*$$

▶ The heights are equivalent to discrete Gaussian ':


$$P_N^{\theta,w}(N_h - \mathbb{E}[N_h] = x) \simeq \frac{1}{Z} e^{-\frac{1}{2\sigma^2}(x)^2}$$



$$\sum_{i=1}^{N_h} (f_h(\ell_i^h/N) - \mathbb{E}[f_h(\ell_i^h/N)]) - \sum_k (N_k - \mathbb{E}[N_k]) \partial_{\varepsilon_k} \mu_{\varepsilon}^h(f_h)|_{\varepsilon=\varepsilon^*}$$

converges towards a centered Gaussian variable.

Discrete β -ensembles, edge fluctuations [Huang-G 17']



Under the previous assumptions, the boundary fluctuates like a Tracy-Widom distribution.

If $\frac{1}{N_1} \sum \delta_{\ell_i^1/N}$ converges towards μ^1 with liquid region $[a, b]$, $\mu^1((-\infty, a)) = 0$, then for all t real

$$\lim_{N \rightarrow \infty} P_N^{\theta, w} \left(N^{2/3} (\ell_1^1/N - a) \geq t \right) = f_{2\theta_{11}}(t)$$

with $f_{2\theta}$ the 2θ -Tracy-Widom distribution appearing in the **continuous** β -ensembles.

Corollary: Fluctuations of the first rows of Young diagrams under Jack deformation of Plancherel measure.

Continuous β -ensembles

The distribution of continuous β -ensembles is given by

$$dP_N^{\beta, V}(\lambda) = \frac{1}{\tilde{Z}_N^{\beta, V}} \prod_{i < j} |\lambda_i - \lambda_j|^\beta e^{-\beta N \sum_{i=1}^N V(\lambda_i)} \prod d\lambda_i.$$

- ▶ When $V(x) = \frac{1}{4}x^2$, and $\beta = 1$ (resp. $\beta = 2, 4$), $P_N^{\beta, V}$ is the distribution of the eigenvalues of a symmetric (resp. Hermitian, resp. symplectic) $N \times N$ matrix with centered Gaussian entries with covariance $1/N$, the **GOE** (resp. **GUE**, resp. **GSE**) .

Continuous β -ensembles

The distribution of continuous β -ensembles is given by

$$dP_N^{\beta,V}(\lambda) = \frac{1}{\tilde{Z}_N^{\beta,V}} \prod_{i < j} |\lambda_i - \lambda_j|^\beta e^{-\beta N \sum_{i=1}^N V(\lambda_i)} \prod d\lambda_i.$$

- ▶ When $V(x) = \frac{1}{4}x^2$, and $\beta = 1$ (resp. $\beta = 2, 4$), $P_N^{\beta,V}$ is the distribution of the eigenvalues of a symmetric (resp. Hermitian, resp. symplectic) $N \times N$ matrix with centered Gaussian entries with covariance $1/N$, the **GOE** (resp. **GUE**, resp. **GSE**) .
- ▶ when $w(x) \simeq e^{-\beta NV(x/N)}$,

$$Z_N^{\theta,w} P_N^{\theta,w}(\ell) \simeq \tilde{Z}_N^{2\theta,V} \frac{dP_N^{2\theta,V}}{d\lambda}(\ell/N)$$

Some techniques to deal with β -ensembles

- ▶ **Integrable systems.** In some cases, these distributions have particular symmetries allowing for special analysis, e.g when $\beta = 2\theta = 2$ the density is the square of a determinant, allowing for orthogonal polynomials analysis [Mehta, Deift, Baik, Johansson etc]
- ▶ **General case.**
 - Dyson-Schwinger (or loop) Equations.* Use equations for the correlators, e.g the moments of the empirical measures, and try to solve them asymptotically [Johansson, Shcherbina, Borot-G, etc]
 - Universality.* Compare your law to one you can analyze [Erdős-Yau et al, Tao-Vu]. An important step is to prove *rigidity* (particles are very close to their deterministic limit), see [Bourgade-Erdős-Yau]

Continuous β -ensembles: Dyson-Schwinger equations(DSE)

Let

$$\hat{\mu}^N = \frac{1}{N} \sum_{i=1}^N \delta_{\lambda_i} : \quad \hat{\mu}_N(f) = \frac{1}{N} \sum_{i=1}^N f(\lambda_i)$$

If f is a smooth test function, then **integration by parts** yields the DSE:

$$N\mathbb{E} \left[\hat{\mu}_N(V'f) - \frac{1}{2} \int \frac{f(x) - f(y)}{x - y} d\hat{\mu}_N(x) d\hat{\mu}_N(y) \right] = \left(\frac{1}{\beta} - \frac{1}{2} \right) \mathbb{E}[\hat{\mu}_N(f_0)']$$

Continuous β -ensembles: Dyson-Schwinger equations

Linearizing this equation around its limit, we get

$$N\mathbb{E}[(\hat{\mu}_N - \mu)(\Xi f)] = \left(\frac{1}{\beta} - \frac{1}{2}\right)\mathbb{E}[\hat{\mu}_N(f')] \\ + \frac{1}{2N}\mathbb{E}\left[\int \int \frac{f(x) - f(y)}{x - y} dN(\hat{\mu}_N - \mu)(x) dN(\hat{\mu}_N - \mu)(y)\right]$$

where

$$\Xi f(x) = V'(x)f(x) - \int \frac{f(x) - f(y)}{x - y} d\mu(y).$$

If we can show that the last term is negligible and Ξ is invertible (off-criticality), we can solve asymptotically this equation to get

$$N\mathbb{E}[\hat{\mu}_N - \mu](f) \simeq \left(\frac{1}{\beta} - \frac{1}{2}\right)\mathbb{E}[\hat{\mu}_N((\Xi^{-1}f)')]$$

We can get an infinite system of DSE for all moments of $\hat{\mu}_N$ and get large N expansion up to any order. This gives the CLT.

Continuous β -ensembles: Fluctuations at the edge

Dumitriu-Edelman 02': Take $V(x) = \beta x^2/2$. Then $P_N^{\beta, \beta x^2/2}$ is the law of the eigenvalues of

$$H_N^\beta = \begin{pmatrix} Y_1^\beta & \xi_1 & 0 & \cdots & 0 \\ \xi_1 & Y_2^\beta & \xi_2 & 0 & \vdots \\ 0 & \ddots & \ddots & \vdots & \vdots \\ 0 & \cdots & \ddots & \ddots & \vdots \\ 0 & 0 & \cdots & \xi_{N-1} & Y_N^\beta \end{pmatrix}$$

where ξ_i are iid $N(0, 1)$ and $Y_i^\beta \simeq \chi_{i\beta}$ independent.

Continuous β -ensembles: Fluctuations at the edge

Dumitriu-Edelman 02': Take $V(x) = \beta x^2/2$. Then $P_N^{\beta, \beta x^2/2}$ is the law of the eigenvalues of

$$H_N^\beta = \begin{pmatrix} Y_1^\beta & \xi_1 & 0 & \cdots & 0 \\ \xi_1 & Y_2^\beta & \xi_2 & 0 & \vdots \\ 0 & \ddots & \ddots & \vdots & \vdots \\ 0 & \cdots & \ddots & \ddots & \vdots \\ 0 & 0 & \cdots & \xi_{N-1} & Y_N^\beta \end{pmatrix}$$

where ξ_i are iid $N(0, 1)$ and $Y_i^\beta \simeq \chi_{i\beta}$ independent.

Ramirez-Rider-Viràg 06': The largest eigenvalue fluctuates like [Tracy-Widom \$\beta\$ distribution](#).

Bourgade-Erdős-Yau 11', Shcherbina 13', Bekerman-Figalli-G 13': [Universality](#): This remains true for general potentials provided off-criticality holds.

Discrete β -ensembles: Nekrasov's equation

Recall $l_{i+1} - l_i - \theta \in \mathbb{N}$ and set

$$P_N^{\theta, w}(\ell) = \frac{1}{Z_N} \prod_{1 \leq i < j \leq N} \frac{\Gamma(l_j - l_i + 1) \Gamma(l_j - l_i + \theta)}{\Gamma(l_j - l_i) \Gamma(l_j - l_i + 1 - \theta)} \prod w(l_i, N)$$

and assume there exists ϕ_N^\pm analytic so that

$$\frac{w(x, N)}{w(x-1, N)} = \frac{\phi_N^+(x)}{\phi_N^-(x)}.$$

Then

$$\phi_N^-(\xi) \mathbb{E}_{P_N^{\theta, w}} \left[\prod_{i=1}^N \left(1 - \frac{\theta}{\xi - l_i} \right) \right] + \phi_N^+(\xi) \mathbb{E}_{P_N^{\theta, w}} \left[\prod_{i=1}^N \left(1 + \frac{\theta}{\xi - l_i - 1} \right) \right]$$

is analytic.

Discrete β -ensembles: Nekrasov's equation

Recall $l_{i+1} - l_i - \theta \in \mathbb{N}$ and set

$$P_N^{\theta, w}(\ell) = \frac{1}{Z_N} \prod_{1 \leq i < j \leq N} \frac{\Gamma(\ell_j - \ell_i + 1) \Gamma(\ell_j - \ell_i + \theta)}{\Gamma(\ell_j - \ell_i) \Gamma(\ell_j - \ell_i + 1 - \theta)} \prod w(\ell_i, N)$$

and assume there exists ϕ_N^\pm analytic so that

$$\frac{w(x, N)}{w(x-1, N)} = \frac{\phi_N^+(x)}{\phi_N^-(x)}.$$

Then

$$\phi_N^-(\xi) \mathbb{E}_{P_N^{\theta, w}} \left[\prod_{i=1}^N \left(1 - \frac{\theta}{\xi - l_i} \right) \right] + \phi_N^+(\xi) \mathbb{E}_{P_N^{\theta, w}} \left[\prod_{i=1}^N \left(1 + \frac{\theta}{\xi - l_i - 1} \right) \right]$$

is analytic.

$$\Rightarrow \text{Gives asymptotic equations for } G_N(z) := \frac{1}{N} \sum_{i=1}^N \frac{1}{z - \frac{l_i}{N}}$$

which can be analyzed as DSE.

Consequences of Nekrasov's equation

- ▶ One can estimate all moments of $N(G_N(z) - G(z))$ for $z \in \mathbb{C} \setminus \mathbb{R}$, proving CLT,

Consequences of Nekrasov's equation

- ▶ One can estimate all moments of $N(G_N(z) - G(z))$ for $z \in \mathbb{C} \setminus \mathbb{R}$, proving CLT,
- ▶ One can estimate all moments of $N(G_N(z) - G(z))$ for $\Im z = \frac{1}{N^{1-\delta}}$ proving rigidity : for any $a > 0$

$$P_N^{\theta, w} \left(\sup_i |\ell_i - N\gamma_i| \geq \frac{N^a}{\min\{i/N, 1 - i/N\}^{1/3}} \right) \leq e^{-(\log N)^2}$$

where $\mu((-\infty, \gamma_i)) = i/N$.

Consequences of Nekrasov's equation

- ▶ One can estimate all moments of $N(G_N(z) - G(z))$ for $z \in \mathbb{C} \setminus \mathbb{R}$, proving CLT,
- ▶ One can estimate all moments of $N(G_N(z) - G(z))$ for $\Im z = \frac{1}{N^{1-\delta}}$ proving rigidity : for any $a > 0$


$$P_N^{\theta, w}(\sup_i |\ell_i - N\gamma_i| \geq \frac{N^a}{\min\{i/N, 1 - i/N\}^{1/3}}) \leq e^{-(\log N)^2}$$

where $\mu((-\infty, \gamma_i)) = i/N$.

- ▶ One can compare the law of the extreme particles, at distance of order $N^{1/3} \gg 1$ (the mesh of the tiling) with the law of the extreme particles for the continuous model and deduce the 2θ -Tracy-Widom fluctuations.

Some open questions

- ▶ **Critical case.** In continuous setting, Bekerman, Leblé, Serfaty 17' derived CLT for functions in $\text{Im}(\Xi)$. What about the discrete case ? Can we get universality of local fluctuations ?
- ▶ **Universality for CLT.** What if we have true Coulomb gas interaction in the discrete case ?
- ▶ **General domains ?**
- ▶ **Local fluctuations in the bulk.** When $\theta = 1$ correlations functions in the bulk converge to discrete Sine process. What about other θ ?



Many thanks to Alexei Borodin, Vadim Gorin and Leonid Petrov for the pictures.

And thank you for your attention!