# A REMARK ON THE INTERSECTION OF PLANE CURVES

C. CILIBERTO, F. FLAMINI, AND M. ZAIDENBERG

ABSTRACT. Let D be a very general curve of degree  $d = 2\ell - \varepsilon$  in  $\mathbb{P}^2$ , with  $\varepsilon \in \{0, 1\}$ . Let  $\Gamma \subset \mathbb{P}^2$  be an integral curve of geometric genus g and degree  $m, \Gamma \neq D$ , and let  $\nu : C \to \Gamma$  be the normalization. Let  $\delta$  be the degree of the *reduction modulo* 2 of the divisor  $\nu^*(D)$  of C (see § 2.1). In this paper we prove the inequality  $4g + \delta \ge m(d - 8 + 2\varepsilon) + 5$ . We compare this with similar inequalities due to Geng Xu ([88, 89]) and Xi Chen ([17, 18]). Besides, we provide a brief account on genera of subvarieties in projective hypersurfaces.

## Contents

Introduction	2
1. Focal loci	3
2. Double planes	5
2.1. The $\delta$ -invariant	5
2.2. Basics on a certain weighted projective 3-space	6
2.3. The even degree case	7
2.4. The odd degree case	8
3. The main result	9
3.1. Constructing appropriate families	9
3.2. Proof of Theorem 3.1	10
4. Genera of subvarieties: a survey	13
References	16

<sup>2010</sup> Mathematics Subject Classification. Primary: 14J70, 14D05; Secondary: 14C17, 14C20, 14H30.

Key words and phrases. Projective hypersurfaces, intersection number, foci, geometric genus, algebraic hyperbolicity.

Acknowledgements: This research started during a visit of the third author at the Dept. of Mathematics of Univ. Roma "Tor Vergata" (supported by this Department, the INdAM "F. Severi" in Rome and the cooperation program GDRE-GRIFGA) and of the second author at the Institut Fourier (supported by INdAM-GNSAGA and within the context of the Laboratory Ypatia of Mathematical Sciences LIA LYSM AMU-CNRS-ECM-INdAM). The authors thank all these institutions and programs for the support and the excellent working conditions. The first author acknowledges the *MIUR Excellence Department Project* awarded to the Department of Mathematics, University of Rome Tor Vergata, CUP E83C18000100006. The work of the third author was partially supported by the grant 346300 for *IMPAN* from the Simons Foundation and the matching 2015-2019 Polish MNiSW fund (code: BCSim-2018-s09). Collaboration has also benefitted of funds Mission Sustainability 2017 - Fam Curves CUP E81-18000100005 (Tor Vergata University).

### INTRODUCTION

Given an effective divisor  $D \in |\mathcal{O}_{\mathbb{P}^n}(d)|$  and an integral (i.e., reduced and irreducible) projective curve  $\Gamma$  of degree m in  $\mathbb{P}^n$ , which is not contained in  $\operatorname{supp}(D)$ , let  $j(D, \Gamma)$  be the order of  $\Gamma \cap D$ . Assume D is very general and set

$$j(n,d,m):=\min\{j(D,\Gamma)\,|\,\Gamma\subset\mathbb{P}^n\text{ as above}\}\quad\text{and}\quad j(n,d):=\min_{m\geqslant 1}\{j(n,d,m)\}.$$

Similarly, with  $\Gamma$  and D as before, let  $i(D,\Gamma)$  stand for the *number of places* of  $\Gamma$  on D, that is, the number of centers of local branches of the curve  $\Gamma$  on D. Then, set

$$i(n,d,m):=\min\{i(D,\Gamma)\,|\,\Gamma\subset\mathbb{P}^n\text{ as above}\}\quad\text{and}\quad i(n,d):=\min_{m\geqslant 1}\{i(n,d,m)\}.$$

The problem of computing j(n, d) and i(n, d) has been considered in [17, 88, 89] (basically devoted to n = 2 case) and [18] (where the case  $n \ge 2$  is considered). The relations of this with the famous Kobayashi problem on hyperbolicity of the complement of a very general hypersurface in  $\mathbb{P}^n$  is well known and we do not dwell on this here (see, e.g., [18]).

Geng Xu ([88, Thm. 1]) proved that

$$j(2,d) = d-2$$
, for any  $d \ge 3$ ,

where the equality is attained either by a bitangent line or by an inflectional tangent line of D, i.e. the minimum is achieved by m = 1. Moreover, for d = 3, he also proved in [89, Corollary] that, for any integer  $m \ge 1$ , the number of rational curves of degree m which meet set-theoretically a given (arbitrary) smooth plane cubic curve D at exactly one point is finite and positive. Therefore, for d = 3 the minimum j(2,3) = 1 is achieved by any integer  $m \ge 1$ .

Xi Chen ([17, Thm. 1.2]) proved that, for d > m, one has

$$j(2, d, m) \ge \min\left\{dm - \frac{m(m+3)}{2}, \ 2dm - 2m^2 - 2\right\}.$$

Furthermore (cf. [17, Cor. 1.1]), for  $d \ge \max\{\frac{3m}{2} - 1, 3\}$  one has

$$j(2, d, m) = dm - \dim(|\mathcal{O}_D(m)|) = dm - \frac{m(m+3)}{2}$$
.

In addition, he conjectured (see [17, Conj. 1.1]) that

 $j(2, d, m) = dm - \dim(|\mathcal{O}_D(m)|)$  if  $d > \max\{m, 2\}.$ 

In arbitrary dimension  $n \ge 2$ , Xi Chen ([18, Thm 1.7]) proved that, for D very general and  $\Gamma$  as above, one has

(0.1) 
$$2g - 2 + i(D, \Gamma) \ge (d - 2n)m,$$

where g is the geometric genus of  $\Gamma$ , i.e., the arithmetic genus of its normalization.

In this paper we obtain a new inequality of type (0.1), although only in the case n = 2 (see Theorem 3.1). Indeed, let D be a very general curve of degree  $d = 2\ell - \varepsilon$  in  $\mathbb{P}^2$ , with  $\varepsilon \in \{0, 1\}$ . Let  $\Gamma$  be an integral curve in  $\mathbb{P}^2$  of geometric genus g and degree  $m, \Gamma \neq D$ , and let  $\nu : C \to \Gamma$  be the normalization. Let  $\delta(D, \Gamma)$  be the degree of the *reduction modulo* 2 of the divisor  $\nu^*(D)$  on C (cf. § 2.1). In Theorem 3.1 we prove that

(0.2) 
$$4g + \delta(D, \Gamma) \ge m(d + 2\varepsilon - 8) + 5.$$

Note that  $\delta(D, \Gamma) \leq i(D, \Gamma)$ , and the equality holds if and only if at any place p of  $\Gamma$  on D, the local intersection multiplicity of D and  $\Gamma$  at p is odd. This happens, for instance, if  $\Gamma$  intersects D transversely. In the latter case  $\delta(D, \Gamma) = i(D, \Gamma) = md$  and both (0.1) and (0.2) are uninteresting. On the other hand, (0.1) and (0.2) become interesting when  $\delta(D, \Gamma)$  and  $i(D, \Gamma)$  are small. Though the difference between the two quantities is a priori unpredictable, one may expect that, generally speaking,  $\delta(D, \Gamma)$  is strictly smaller than  $i(D, \Gamma)$ . Unfortunately, the genus g works against us in (0.2); however, for g = 0, 1 and d even, (0.2) is better than (0.1). Further related problems have been recently considered in [19, 64, 65].

As a final remark, note that (0.2) is more useful than (0.1) if one looks, as we do in this paper, at the geometric genera of curves contained in a double plane  $X_d$ , that is, a cyclic double cover of  $\mathbb{P}^2$  branched along a very general plane curve D of even degree d. For instance, letting  $q = 0, \delta(D, \Gamma) = 0, 2$  and d even, we are looking actually for rational curves on  $X_d$ . By (0.2) we see that such a rational curve over  $\Gamma$  might exist, as expected, only for  $d \leq 6$  (for low m one has even smaller bounds on d). The case d = 6 corresponds to a K3 surface, which always contains infinitely many rational curves. In contrast, it follows from (0.2) that the double planes with very general branching curves of even degree  $d \ge 8$  ( $d \ge 10$ , respectively) do not carry any rational curve (any rational or elliptic curve, respectively, hence are algebraically hyperbolic). For d = 8 and d = 10 these double planes are Horikawa surfaces  $H_8$  and  $H_{10}$ , that is, their Chern numbers satisfy  $c_2 = 5c_1^2 + 36$  (in other words,  $(c_1^2, c_2)$  lies on the Noether line). The algebraic hyperbolicity of  $H_{10}$  was established first by X. Roulleau and E. Rousseau ([75]). J. Liu ([65]) showed that some of the Horikawa surfaces  $H_{10}$  are even Kobayashi hyperbolic, whereas there is no hyperbolic  $H_8$ . Indeed, the Horikawa double planes  $H_8$  carrying elliptic curves are dense in the set of all such surfaces, while the Kobayashi hyperbolicity is open in the Hausdorff topology.

The proof of Theorem 3.1 presented in §3 follows, with minor variations due to the different setting, the basic ideas exploited in [21] (and later in [22]). These are based on a smart use of the theory of *focal loci*, see e.g. [20]. For the reader's convenience, we recall in §1 the basic notions and results of this theory. We apply this technique to families of double covers of  $\mathbb{P}^2$  branched along a very general plane curve D or along D plus a general line, according to whether the degree of D is even (see § 2.3 and § 3.2.1) or odd (see § 2.4 and § 3.2.2).

In the last §4 we provide a short survey on genera of subvarieties in projective varieties, with accent on projective hypersurfaces.

## 1. Focal loci

For the reader's convenience, we recall here some basic notions from [20, 21].

Let X be a smooth projective variety of dimension n+1. Assume we have a flat, projective family  $\mathcal{D} \xrightarrow{p} \mathcal{B}$  of effective divisors on X over a smooth, irreducible, quasiprojective base  $\mathcal{B}$ , with irreducible general fiber. Up to shrinking  $\mathcal{B}$  to a suitable Zariski dense, open subset, we may suppose that for any closed point  $b \in \mathcal{B}$ the fiber  $D_b$  of  $\mathcal{D} \xrightarrow{p} \mathcal{B}$  over b is irreducible. Assume we have a commutative diagram




where  $\mathbf{q} : \mathfrak{C} \to \mathcal{B}$  is a flat projective family such that, for all  $b \in \mathcal{B}$ , the fiber  $\Gamma_b$  over b is a reduced curve of *geometric genus* g, and where i is an inclusion: so, for any  $b \in \mathcal{B}$ , one has  $\Gamma_b \subset D_b$  via the inclusion  $i_b$ .

By a result of Tessier (see [80, Théorème 1]), there is a simultaneous normalization



such that  $\mathcal{C}$  is smooth and, for every  $b \in \mathcal{B}$ , the fiber  $C_b$  of  $q : \mathcal{C} \to \mathcal{B}$  is the normalization  $\nu_b : C_b \to \Gamma_b$  of  $\Gamma_b$ . For any  $b \in \mathcal{B}$ , the curve  $C_b$  is smooth of *(arithmetic) genus g.* 

Composing with the inclusion  $\mathcal{D} \stackrel{\mathcal{I}}{\hookrightarrow} \mathcal{B} \times X$ , we get the commutative diagram (1.3)



where  $pr_i$  is the projection onto the *i*th factor, for i = 1, 2. We set

$$s := j \circ i \circ \nu \colon \mathcal{C} \to \mathcal{B} \times X$$

and let  $\mathcal{N} := \mathcal{N}_s$  be the normal sheaf to s, defined by the exact sequence

$$0 \longrightarrow \mathcal{T}_{\mathcal{C}} \xrightarrow{ds} s^*(\mathcal{T}_{\mathcal{B} \times X}) \longrightarrow \mathcal{N} \longrightarrow 0,$$

where  $\mathcal{T}_Y$  stands for the tangent sheaf of a smooth variety Y. For  $b \in \mathcal{B}$  we set

$$N_b := \mathcal{N}|_{C_b} = \mathcal{N} \otimes \mathcal{O}_{C_b}$$
 and  $s_b = s|_{C_b} : C_b \to \{b\} \times X = X$ .

In addition, we let

$$\varphi := \operatorname{pr}_2 \circ s : \mathcal{C} \to X$$
.

Then  $\varphi_b = \varphi|_{C_b}$  coincides with  $s_b$  for any  $b \in \mathcal{B}$ , that is,

$$\varphi_b = s_b: \ C_b \xrightarrow{\nu_b} \Gamma_b \xrightarrow{(i_b)} D_b \xrightarrow{(\operatorname{pr}_2 \circ j)_b} X$$

As in  $[21, \S 2]$ , we set

(1.4) 
$$z(\mathcal{C}) := \dim (\varphi(\mathcal{C})),$$

so that  $z(\mathcal{C}) \leq n+1 = \dim(X)$ . If  $z(\mathcal{C}) = n+1$  one says that  $\mathfrak{C} \xrightarrow{\mathsf{q}} \mathcal{B}$ , or  $\mathcal{C} \xrightarrow{q} \mathcal{B}$ , is a *covering family*.

Proposition 1.1 (See [20, Prop. 1.4 and p. 98]). In the above setting, we have:

(a) for any  $b \in \mathcal{B}$ , the sheaf  $N_b$  fits into the exact sequence

$$0 \longrightarrow \mathcal{T}_{C_b} \xrightarrow{ds_b} s_b^*(\mathcal{T}_X) \longrightarrow N_b \longrightarrow 0$$

and  $\mathcal{C} \xrightarrow{q} \mathcal{B}$  induces on  $C_b$  a characteristic map

$$\chi_b: T_{\mathcal{B},b} \otimes \mathcal{O}_{C_b} \longrightarrow N_b$$
,

where  $T_{\mathcal{B},b}$  denotes the tangent space to  $\mathcal{B}$  at b;

(b) if  $b \in \mathcal{B}$  and  $x \in C_b$  are general points, then

$$\dim(N_{b,x}) = \dim(s_b^*(\mathcal{T}_X)_x) - \dim(\mathcal{T}_{C_b,x}) = n \quad \text{and} \quad \operatorname{rk}(\chi_{b,x}) = z(\mathcal{C}) - 1.$$

Definition 1.2 (See [20, Def. (1.5)]). Given  $b \in \mathcal{B}$ , the focal set of  $\mathcal{C} \xrightarrow{q} \mathcal{B}$  on  $C_b$  is the closed subscheme  $\Phi_b$  of  $C_b$  defined as

$$\Phi_b := \{ x \in C_b | \operatorname{rk}(\chi_{b,x}) < z(\mathcal{C}) - 1 \}.$$

If  $b \in \mathcal{B}$  is general, then  $\Phi_b$  is a proper subscheme of  $C_b$ . The points in  $\Phi_b$  are called focal points of  $\mathcal{C} \xrightarrow{q} \mathcal{B}$  on  $C_b$ . We denote by  $\Phi_b^{\mathrm{sm}}$  the open subset of  $\Phi_b$  consisting of the points  $x \in \Phi_b$  which map to smooth points of  $\Gamma_b$  via the normalization morphism  $\nu_b \colon C_b \to \Gamma_b$ .

Proposition 1.3 ([21, Prop. 2.3 and Prop. 2.4]). Let  $\mathcal{C} \longrightarrow \mathcal{B}$  be a covering family. Then the following hold.

- (i) Suppose that for  $x \in C_b$  the point s(x) is smooth in both  $\Gamma_b$  and  $D_b$ . Assume also that s(x) is a fundamental point of the family  $\mathcal{D} \xrightarrow{p} \mathcal{B}$ , i.e. it is a base point of the family. Then  $x \in \Phi_b^{\mathrm{sm}}$ .
- (ii) For a general point  $b \in \mathcal{B}$  one has

(1.5) 
$$\deg(\Phi_b^{\rm sm}) \leqslant 2g - 2 - K_X \cdot \Gamma_b \,.$$

#### 2. Double planes

In this section we collect useful material for the proof of our main result. The result itself is stated and proven in §3. The contents of this section, which suffice for our applications, can be easily adapted to the higher dimensional case.

2.1. The  $\delta$ -invariant. Let C be any smooth, irreducible, projective curve, and let  $\Delta = \sum_{i} m_i p_i$  be an effective divisor on C. We set  $\Delta_2 := \sum_{i} \overline{m}_i p_i$ , where  $\overline{m}_i \in \{0, 1\}$  is the residue of the integer  $m_i$  modulo 2. We also set  $\delta_2(\Delta) := \deg(\Delta_2)$ . For any smooth curve  $D \subset \mathbb{P}^2$  and any integral curve  $\Gamma \subset \mathbb{P}^2$ ,  $\Gamma \neq D$ , with

normalization  $\nu \colon C \to \Gamma$ , we set

(2.1) 
$$\delta(D,\Gamma) := \delta_2(\nu^*(D)).$$

We notice that

$$\delta(D,\Gamma) \leqslant i(D,\Gamma).$$

2.2. Basics on a certain weighted projective 3-space. For any positive integer  $\ell$ , we denote by  $\mathcal{L}_{\ell}$  the linear system  $|\mathcal{O}_{\mathbb{P}^2}(\ell)|$  of plane curves of degree  $\ell$ , and by  $\mathcal{U}_{\ell}$  its open dense subset of points corresponding to smooth curves. We let  $N_{\ell} = \dim(\mathcal{L}_{\ell}) = \frac{\ell(\ell+3)}{2}$ . We denote by  $\mathcal{D}_{\ell} \to \mathcal{L}_{\ell}$  the universal curve, and we use the same notation  $\mathcal{D}_{\ell} \to \mathcal{U}_{\ell}$  for its restriction to  $\mathcal{U}_{\ell}$ .

The linear system  $\mathcal{L}_{\ell}$  determines the  $\ell$ th Veronese embedding  $\mathbb{P}^2 \stackrel{v_{\ell}}{\hookrightarrow} \mathbb{P}^{N_{\ell}}$ , whose image is the  $\ell$ -Veronese surface  $V_{\ell}$  in  $\mathbb{P}^{N_{\ell}}$ . Let  $[x] = [x_0, x_1, x_2]$  be homogeneous coordinates in  $\mathbb{P}^2$ , and let

 $[x^{I}]$ , where  $I = (i_{0}, i_{1}, i_{2})$  is a multiindex such that  $|I| = i_{0} + i_{1} + i_{2} = \ell$ ,

be homogeneous coordinates in  $\mathbb{P}^{N_\ell}.$  In these coordinates the Veronese map is given by

$$\mathbb{P}^2 \ni [x] \xrightarrow{v_\ell} [x^I]_{|I|=\ell} \in \mathbb{P}^{N_\ell}, \text{ where } x^I := x_0^{i_0} x_1^{i_1} x_2^{i_2}.$$

We equip the weighted projective 3-space  $\mathbb{P}(1, 1, 1, \ell)$  with weighted homogeneous coordinates  $[x, z] := [x_0, x_1, x_2, z]$ , where  $x_0, x_1, x_2$  [resp. z] have weight 1 [resp. has weight  $\ell$ ]. We introduce as well coordinates  $[x^I, z]_{|I|=\ell}$  in  $\mathbb{P}^{N_{\ell}+1}$  and embed  $\mathbb{P}^{N_{\ell}}$  in  $\mathbb{P}^{N_{\ell}+1}$  as the hyperplane  $\Pi$  with equation z = 0. Then  $\mathbb{P}(1, 1, 1, \ell)$  can be identified with the cone  $W_{\ell} \subset \mathbb{P}^{N_{\ell}+1}$  over the *l*-Veronese surface  $V_{\ell}$  with vertex  $P = [0, \ldots, 0, 1]$ . Blowing P up yields a minimal resolution

$$\rho: Z_{\ell} \to W_{\ell} \cong \mathbb{P}(1, 1, 1, \ell)$$

with exceptional divisor  $E \cong V_{\ell} \cong \mathbb{P}^2$ . The projection from P induces a  $\mathbb{P}^1$ -bundle structure

$$\pi\colon Z_\ell\to V_\ell\cong\mathbb{P}^2\,.$$

Let f be the class of a fiber of  $\pi$ . One has

$$Z_{\ell} \cong \mathbb{P}(\mathcal{O}_{\mathbb{P}^2}(\ell) \oplus \mathcal{O}_{\mathbb{P}^2}) \text{ and } \mathcal{O}_{Z_{\ell}}(1) = \rho^*(\mathcal{O}_{W_{\ell}}(1))$$

For every integer m, we set

$$\mathcal{O}_{\ell}(m) := \pi^*(\mathcal{O}_{\mathbb{P}^2}(m)) \text{ and } \mathcal{L}_{\ell}(m) := |\mathcal{O}_{\ell}(m)|.$$

Note that

$$\mathcal{O}_{Z_{\ell}}(1) \cong \mathcal{O}_{\ell}(\ell) \otimes \mathcal{O}_{Z_{\ell}}(E).$$

(2.3) Since

(2.2)

$$E \cong \mathbb{P}^2$$
,  $\mathcal{O}_{Z_\ell}(1) \otimes \mathcal{O}_E \cong \mathcal{O}_E$ , and  $\mathcal{O}_\ell(\ell) \otimes \mathcal{O}_E \cong \mathcal{O}_{\mathbb{P}^2}(\ell)$ ,

we deduce

(2.4) 
$$\mathcal{O}_{Z_{\ell}}(E) \otimes \mathcal{O}_{E} \cong \mathcal{O}_{\mathbb{P}^{2}}(-\ell).$$

Finally, we denote by  $K_{\ell}$  the canonical sheaf of  $Z_{\ell}$ .

Lemma 2.1. One has

$$K_{\ell} \cong \mathcal{O}_{\ell}(\ell-3) \otimes \mathcal{O}_{Z_{\ell}}(-2) \cong \mathcal{O}_{\ell}(-\ell-3) \otimes \mathcal{O}_{Z_{\ell}}(-2E)$$

*Proof.* The Picard group  $\operatorname{Pic}(Z_{\ell})$  is freely generated by the classes  $\mathcal{O}_{\ell}(1)$  and  $\mathcal{O}_{Z_{\ell}}(E)$ , and also by  $\mathcal{O}_{\ell}(1)$  and  $\mathcal{O}_{Z_{\ell}}(1)$ , see (2.3). Let H [resp. L] be a general member of  $|\mathcal{O}_{Z_{\ell}}(1)|$  [resp. of  $\mathcal{L}_{\ell}(1)$ ]. Write  $K_{\ell} \sim \alpha H + \beta L$ , where  $\alpha, \beta \in \mathbb{Z}$ . From the relations  $K_{\ell} \cdot f = -2$ ,  $H \cdot f = 1$ , and  $L \cdot f = 0$  one gets  $\alpha = -2$ .

By adjunction formula and (2.4) we obtain

$$\mathcal{O}_{\mathbb{P}^2}(-3) \cong K_E = (K_\ell + E)|_E \cong (-2H + \beta L + E)|_E \cong \mathcal{O}_{\mathbb{P}^2}(\beta - \ell).$$

So,  $\beta = \ell - 3$ , as desired.

Finally, let  $\mathbb{G}_{\ell}$  be the group of all automorphisms of  $\mathbb{P}(1, 1, 1, \ell)$  which stabilize the divisor with equation z = 0. This group is naturally isomorphic to the automorphism group of the pair  $(W_{\ell}, V_{\ell})$ , i.e. automorphisms of  $W_{\ell}$  stabilizing  $V_{\ell}$ , where  $V_{\ell}$  is cut out on  $W_{\ell}$  by  $\Pi$ . In turn, the latter group is isomorphic to the automorphism group of the pair  $(Z_{\ell}, \rho^*(V_{\ell}))$ . One has the exact sequence

(2.5) 
$$1 \to \mathbb{C}^* \to \mathbb{G}_\ell \to \mathrm{PGL}(3, \mathbb{C}) \to 1.$$

2.3. The even degree case. Let D be a smooth curve in  $\mathbb{P}^2$  of even degree  $d = 2\ell \ge 2$  which, in the homogeneous coordinate system fixed in §2.2, is given by equation  $f(x_0, x_1, x_2) = 0$ , where f is a homogeneous polynomial of degree d. Viewed as a hypersurface of  $V_\ell$ , D is cut out on  $V_\ell$  by a quadric with equation  $Q(x^I)_{|I|=\ell} = 0$ , where Q is a homogeneous polynomial of degree 2 in the variables  $\{x^I\}_{|I|=\ell}$ .

The double plane associated with D is the double cover  $\psi : \mathbb{D}^* \to \mathbb{P}^2$  branched along D. It can be embedded in  $\mathbb{P}(1, 1, 1, \ell)$  as a hypersurface  $\mathbb{D}_a^*$  defined by a (weighted homogeneous) equation of the form  $az^2 = f(x_0, x_1, x_2)$ , for any  $a \in \mathbb{C}^*$ . Under the identification of  $\mathbb{P}(1, 1, 1, \ell)$  with  $W_\ell$ , we see that  $\mathbb{D}_a^*$  is cut out on  $W_\ell$  by a quadric in  $\mathbb{P}^{N_\ell + 1}$  of the form  $az^2 = Q(x^I)_{|I| = \ell}$ .

Consider the sublinear system  $\mathcal{Q}_{\ell}$  of  $|\mathcal{O}_{W_{\ell}}(2)|$  of surfaces cut out on  $W_{\ell}$  by the quadrics of  $\mathbb{P}^{N_{\ell}+1}$  with equation of the form  $az^2 = Q(x^I)_{|I|=\ell}$ .

When  $a \neq 0$ , the quadrics in question are such that their polar hyperplane with respect to P has equation z = 0. When a = 0, such a quadric is singular at P, it represents the cone with vertex at P over the quadric in  $\Pi = \{z = 0\} \cong \mathbb{P}^{N_{\ell}}$  with equation  $Q(x^{I})_{|I|=\ell} = 0$  and it cuts out on  $W_{\ell}$  a cone, with vertex at P, over a quadric section of  $V_{\ell}$ .

In particular,  $\mathcal{Q}_{\ell}$  contains the codimension 1 sublinear system  $\mathcal{Q}_{\ell}^{c}$  of all such cones with vertex at P over a quadric section of  $V_{\ell}$ , thus  $\dim(\mathcal{Q}_{\ell}) = N_{d} + 1$ . Moreover  $\mathcal{Q}_{\ell}$  is stable under the action of  $\mathbb{G}_{\ell}$  on  $W_{\ell}$ .

We set  $\widetilde{\mathcal{Q}}_{\ell} := \rho^*(\mathcal{Q}_{\ell})$ , which is a sublinear system of  $|\mathcal{O}_{Z_{\ell}}(2)|$ . Note that  $\widetilde{\mathcal{Q}}_{\ell}$  contains the sublinear system  $\widetilde{\mathcal{Q}}_{\ell}^c = \rho^*(\mathcal{Q}_{\ell}^c)$  of all divisors of the form 2*E* plus a divisor in  $\mathcal{L}_{\ell}(d)$ .

We denote by  $\mathcal{Q}_{\ell}^*$  the dense open subset of  $\mathcal{Q}_{\ell}$  of points corresponding to smooth surfaces. Since no surface in  $\mathcal{Q}_{\ell}^*$  passes through P, we may and will identify  $\mathcal{Q}_{\ell}^*$ with its pull-back via  $\rho$  on  $Z_{\ell}$ , which is a dense open subset of  $\widetilde{\mathcal{Q}}_{\ell}$  sitting in the complement of  $\widetilde{\mathcal{Q}}_{\ell}^c$ . We denote by  $\mathcal{D}_{\ell}^* \to \mathcal{Q}_{\ell}^*$  the universal family.

A surface  $\mathbb{D}^* \in \mathcal{Q}_{\ell}^*$  cuts out on  $V_{\ell}$  a smooth curve  $D \in \mathcal{U}_{\ell}$  and conversely; indeed the projection from P realizes  $\mathbb{D}^*$  as the double cover of  $\mathbb{P}^2$  branched along D. This yields a surjective morphism

$$\mathcal{Q}_{\ell}^* \ni \mathbb{D}^* \xrightarrow{\beta} \mathbb{D}^* \cap V_{\ell} := D \in \mathcal{U}_d$$
,

which sends the double plane  $\mathbb{D}^*$  to its branching divisor D. This morphism is equivariant under the actions of  $\mathbb{G}_{\ell}$  on both  $\mathcal{Q}_{\ell}^*$  and  $\mathcal{U}_d$ , where  $\mathbb{G}_{\ell}$  acts on  $\mathcal{U}_d$  via the natural action of the quotient group PGL(3,  $\mathbb{C}$ ), see (2.5). The morphism  $\beta$  is not injective, its fibers being isomorphic to  $\mathbb{C}^*$ .

As an immediate consequence of Lemma 2.1, we have:

Lemma 2.2. Let D be a smooth curve in  $\mathbb{P}^2$  of even degree  $d = 2\ell \ge 2$ , let  $\psi : \mathbb{D}^* \to \mathbb{P}^2$  be the double cover branched along D, let  $\Gamma \subset \mathbb{P}^2$  be a projective curve of

degree m not containing D, and let  $\Gamma^*$  be its pull–back via  $\psi$  considered as a curve in  $Z_\ell.$  One has

(2.6) 
$$K_{\ell} \cdot \Gamma^* = -m(d+6).$$

In the setting of Lemma 2.2, consider the diagram

$$(2.7) \qquad \qquad \begin{array}{c} C^* \xrightarrow{\nu^*} & \Gamma^* \\ \psi' \downarrow & & \downarrow \psi \\ C \xrightarrow{\nu} & \Gamma \end{array}$$

where  $\nu$  and  $\nu^*$  are the normalization morphisms and  $\psi$  and  $\psi'$  have degree 2 (to ease notation, here we have identified  $\psi$  with its restriction to  $\Gamma^*$ ).

Let  $\delta := \delta(D, \Gamma)$ . It could be that  $C^*$  splits into two components both isomorphic to C; in this case  $\delta = 0$ . If  $\delta = 0$  and the genus of C is zero, then  $C^*$  certainly splits. Suppose that  $C^*$  is irreducible, and let g and  $g^*$  be the geometric genera of  $\Gamma$  and  $\Gamma^*$  (i.e. the arithmetic genera of C and  $C^*$ , respectively). Since  $\psi'$  has exactly  $\delta$  ramification points, the Riemann-Hurwitz formula yields

(2.8) 
$$2(g^* - 1) = 4(g - 1) + \delta.$$

2.4. The odd degree case. Fix a line  $h \in |\mathcal{O}_{\mathbb{P}^2}(1)|$ , and let D be a smooth curve in  $\mathbb{P}^2$  of degree  $d = 2\ell - 1 \ge 1$ , which intersects h transversely. We denote by  $\mathcal{U}_d^h$ the open subset of  $\mathcal{U}_d$  consisting of such curves.

For each  $D \in \mathcal{U}_d^h$ , we consider the reducible curve of degree  $d + 1 = 2\ell$ 

$$\Delta := D + h \in |\mathcal{O}_{\mathbb{P}^2}(d+1)|$$

and the double cover  $\psi \colon \mathbb{D}^* \to \mathbb{P}^2$ , branched along  $\Delta$ . The difference with the even degree case is that  $\mathbb{D}^*$  is no longer smooth, but it has double points at the d points in  $D \cap h$ . In any event, as in the even degree case, we can consider the set  $\mathcal{Q}^*_{\ell;h} \subset |\mathcal{O}_{Z_\ell}(2)|$  of all such surfaces  $\mathbb{D}^*$ , with its universal family  $\mathcal{D}^*_{\ell;h} \to \mathcal{Q}^*_{\ell;h}$  which parametrizes all double planes  $\psi \colon \mathbb{D}^* \to \mathbb{P}^2$  as above. We still have the morphism

$$\beta: \mathcal{Q}^*_{\ell:h} \to \mathcal{U}^h_d$$

associating to  $\mathbb{D}^*$  the branching divisor  $\Delta$  of  $\varphi \colon \mathbb{D}^* \to \mathbb{P}^2$  minus h.

The group acting here is no longer the full group  $\mathbb{G}_{\ell}$  but its subgroup  $\mathbb{G}_{\ell;h}$  which fits in the exact sequence

$$1 \to \mathbb{C}^* \to \mathbb{G}_{\ell;h} \to \operatorname{Aff}(2,\mathbb{C}) \to 1$$

where  $\operatorname{Aff}(2,\mathbb{C})$  is the *affine group* of all projective transformations in PGL(3,  $\mathbb{C}$ ) stabilizing h.

Keeping the setting and notation of §2.3, Lemma 2.2 still holds, as well as diagram (2.7). If  $\Gamma$  intersects h at m distinct points which are off D, then the double cover  $\psi' : C^* \to C$  has  $\delta + m \ge m > 0$  ramification points, where  $\delta = \delta(D, \Gamma)$  as above. In particular,  $C^*$  is irreducible, and (2.8) is replaced by

(2.9) 
$$2(g^* - 1) = 4(g - 1) + \delta + m.$$

#### 3. The main result

In this section we prove the following:

**Theorem 3.1.** Let  $\delta \ge 0$  be an integer such that, for a very general curve D in  $\mathbb{P}^2$ of degree  $d = 2\ell - \varepsilon$ , where  $\varepsilon \in \{0, 1\}$ , there exists an integral curve  $\Gamma \subset \mathbb{P}^2$ ,  $\Gamma \neq D$ , of geometric genus g and degree m with  $\delta(D, \Gamma) = \delta$ . Then

$$(3.1) 4g + \delta \ge m(d + 2\varepsilon - 8) + 5.$$

The proof of Theorem 3.1 will be done in §3.2. First we need some more preliminaries, which we collect in the next subsection. We keep all notation and conventions introduced so far.

3.1. Constructing appropriate families. Fix integers  $m \ge 1$  and  $g \ge 0$ . Let  $\mathcal{H}$  be the locally closed subset of  $\mathcal{L}_m$ , whose points correspond to integral curves  $\Gamma \subset \mathbb{P}^2$  of degree m and geometric genus g;  $\mathcal{H}$  is a quasiprojective variety. We let  $\mathcal{U} \to \mathcal{H}$  be the universal curve.

3.1.1. The even degree case. Fix an even positive integer  $d = 2\ell$  and a non-negative integer  $\delta$ . Consider the locally closed subset  $\mathcal{I}$  of  $\mathcal{H} \times \mathcal{Q}_{\ell}^*$  of pairs  $(\Gamma, \mathbb{D}^*)$  such that  $\Gamma$  does not coincide with the branch curve D of  $\psi : \mathbb{D}^* \to \mathbb{P}^2$  and  $\delta(D, \Gamma) = \delta$ . Remember that we may equivalently interpret  $\mathbb{D}^*$  as a surface in  $W_{\ell}$  or in  $Z_{\ell}$ . Each irreducible component of  $\mathcal{I}$  is fixed by the obvious action of  $\mathbb{G}_{\ell}$  on  $\mathcal{H} \times \mathcal{Q}_{\ell}^*$ .

For any  $(\Gamma, \mathbb{D}^*) \in \mathcal{I}$ , the pull-back  $\Gamma^* \subset \mathbb{D}^*$  of  $\Gamma$  via  $\psi$  is a reduced curve in  $Z_{\ell}$ . Hence there is a morphism  $\mu : \mathcal{I} \to \mathcal{K}$ , where  $\mathcal{K}$  is the Hilbert scheme of curves of  $Z_{\ell}$ . We let  $\mathcal{V} \to \mathcal{K}$  be the corresponding universal family. The map  $\mu$  is equivariant under the actions of  $\mathbb{G}_{\ell}$  on both  $\mathcal{I}$  and  $\mathcal{K}$ .

Let  $\pi_1: \mathcal{I} \to \mathcal{H}$  and  $\pi_2: \mathcal{I} \to \mathcal{Q}_{\ell}^*$  be the two projections. Under the hypotheses of Theorem 3.1 and with notation as in § 2.3, the following holds.

Lemma 3.2. There exists an irreducible component I of  $\mathcal{I}$  which dominates  $\mathcal{Q}_{\ell}$  via  $\pi_2$ . Hence I dominates also  $\mathcal{U}_d \subset \mathcal{L}_d$  via  $\beta \circ \pi_2$ .

Given I as in Lemma 3.2, we choose an irreducible, smooth subvariety  $\mathcal{B}$  of I, such that  $\pi_2$  restricts to an étale morphism of  $\mathcal{B}$  onto its image, which is dense in  $\mathcal{Q}_{\ell}$ . To place our objects in the context of §1, consider the universal family  $\mathcal{D}_{\ell}^* \to \mathcal{Q}_{\ell}^*$  (cf. § 2.3) of double planes  $\mathbb{D}^*$  [resp.  $\mathcal{V} \to \mathcal{K}$  of curves  $\Gamma^* \subset \mathbb{D}^*$ ]. Up to possibly shrinking  $\mathcal{B}$  and performing an étale cover of it, the morphisms  $\mathcal{B} \xrightarrow{\pi_2} \mathcal{Q}_{\ell}^*$ and  $\mathcal{B} \xrightarrow{\mu} \mathcal{K}$  give rise to families

$$\mathcal{D} := \pi_2^*(\mathcal{I}\!\mathcal{D}_\ell^*) \xrightarrow{p} \mathcal{B} \text{ and } \mathfrak{C} := \mu^*(\mathcal{V}) \xrightarrow{\mathfrak{q}} \mathcal{B}.$$

over  $\mathcal{B}$  fitting in diagram (1.1). We may assume that there exists a simultaneous normalization  $\nu$  and a family  $\mathcal{C} \xrightarrow{q} \mathcal{B}$  as in (1.2), with  $\mathcal{C}$  smooth fitting in (1.3), where  $X = Z_{\ell}$ .

3.1.2. The odd degree case. Fix now an odd positive integer  $d = 2\ell - 1$  and a nonnegative integer  $\delta$ , and fix a line h in  $\mathbb{P}^2$ . We consider the locally closed subset  $\mathcal{I}$ of  $\mathcal{H} \times \mathcal{Q}_{\ell;h}$  consisting of pairs  $(\Gamma, \mathbb{D}^*) \in \mathcal{H} \times \mathcal{Q}_{\ell}^*$  such that  $\Gamma$  is not contained in the branch divisor  $\Delta$  of  $\psi : \mathbb{D}^* \to \mathbb{P}^2$ ,  $\delta(D, \Gamma) = \delta$ , and  $\Gamma$  intersects h at m distinct points which are off D. For any point  $(\Gamma, \mathbb{D}^*) \in \mathcal{I}$ , the pull-back  $\Gamma^* \subset \mathbb{D}^*$  of  $\Gamma$  via  $\psi$  is an integral curve in  $Z_{\ell}$ . So, we still have the morphisms  $\mu : \mathcal{I} \to \mathcal{K}, \pi_1 : \mathcal{I} \to \mathcal{H}$  and  $\pi_2 : \mathcal{I} \to \mathcal{Q}^*_{\ell;h}$ equivariant under actions of  $\mathbb{G}_{\ell;h}$ .

As before, we have the following

Lemma 3.3. There exists an irreducible component I of  $\mathcal{I}$  which dominates  $\mathcal{Q}^*_{\ell;h}$  via  $\pi_2$ .

As in the even case, given I as in Lemma 3.3, we may construct a smooth  $\mathcal{B}$  having an étale, dominant morphism to  $\mathcal{Q}_{\ell;h}$ , together with families

$$\mathcal{D} := \pi_2^*(\mathcal{I}\!\mathcal{D}_{\ell,h}^*) \xrightarrow{p} \mathcal{B}, \quad \mathfrak{C} := \mu^*(\mathcal{V}) \xrightarrow{\mathfrak{q}} \mathcal{B},$$

fitting in diagram (1.1). Consider a simultaneous normalization  $\nu$  and a family  $\mathcal{C} \xrightarrow{q} \mathcal{B}$  as in (1.2), with  $\mathcal{C}$  smooth. In view of Lemmata 3.2 and 3.3, the constructed families fit in diagram (1.3), with  $X = Z_{\ell}$ .

In both cases, the next lemma allows to apply Proposition 1.3 in our setting.

Lemma 3.4. For any d > 0,  $\mathcal{C} \xrightarrow{q} \mathcal{B}$  is a covering family, i.e.  $z(\mathcal{C}) = 3$ .

*Proof.* By the discussion in §3.1, for d even  $\varphi(\mathcal{C})$  is stable under the action of  $\mathbb{G}_{\ell}$  on  $Z_{\ell}$ , which is transitive; for d odd  $\varphi(\mathcal{C})$  is stable under the action of  $\mathbb{G}_{\ell;h}$ , which is transitive on the dense open subset of  $Z_{\ell}$  whose complement is  $\pi^{-1}(h) \cup E$ . This proves the assertion.

3.2. **Proof of Theorem 3.1.** Our proof follows the one of Theorem (1.2) in [21]. First we recall the following useful fact.

Lemma 3.5 (See [21, Lemma (3.1)]). Let  $g: V \to W$  be a linear map of finite dimensional vector spaces. Suppose that  $\dim(g(V)) > k$ . Let  $\{V_i\}_{i \in I}$  be a family of vector subspaces of V, such that  $\bigcup_{i \in I} V_i$  spans V, and for any pair  $(i, j) \in I \times I$ , with  $i \neq j$ , there is a finite sequence  $i_1 = i, i_2, \ldots, i_{t-1}, i_t = j$  of distinct elements of I with  $\dim(g(V_{i_h} \cap V_{i_{h+1}})) \ge k$ , for all  $h \in \{1, \ldots, t-1\}$ . Then there is an index  $i \in I$  such that  $\dim(g(V_i)) > k$ .

3.2.1. The even degree case. We need to construct a suitable subfamily of  $\mathfrak{C} \to \mathcal{B}$  with the covering property.

Fix a general point  $b_0 \in \mathcal{B}$ , and let  $\Gamma_0^*$  and  $\mathbb{D}_0^*$  be the corresponding elements of the families  $\mathfrak{C} \to \mathcal{B}$  and  $\mathcal{D} \to \mathcal{B}$ , respectively.

Let  $\mathcal{L}$  be the open subset of the linear system  $\mathcal{L}_{\ell}(d-1)$  as in (2.2) consisting of the surfaces  $F \in \mathcal{L}_{\ell}(d-1)$  which do not contain  $\Gamma_0^*$ . A general such surface Fmeets  $\Gamma_0^*$  transversely. By genericity, we may suppose that all surfaces F defined by the pull-back via  $\pi : \mathbb{Z}_{\ell} \to V_{\ell} \cong \mathbb{P}^2$  of degree d-1 monomials in the variables  $x_0, x_1, x_2$  belong to  $\mathcal{L}$ . For a given  $F \in \mathcal{L}$ , we denote by  $\mathcal{B}_F$  the subvariety of  $\mathcal{B}$ parameterizing all double planes in  $\mathcal{D} \to \mathcal{B}$  containing the complete intersection curve of F and  $\mathbb{D}_0^*$ . In addition, for a general point  $\xi \in \Gamma_0^*$  we let  $\mathcal{B}_{F,\xi}$  denote the subvariety of  $\mathcal{B}_F$  parameterizing all surfaces in  $\mathcal{D} \to \mathcal{B}_F$  which pass through  $\xi$ .

Lemma 3.6. For  $F \in \mathcal{L}$  and  $\xi \in \Gamma_0^*$  as above one has

 $\dim(\mathcal{B}_F) = 3$  and  $\dim(\mathcal{B}_{F,\xi}) = 2$ .

Furthermore,  $b_0$  is a smooth point of both  $\mathcal{B}_F$  and  $\mathcal{B}_{F,\xi}$ .

10

*Proof.* Consider the sublinear system  $\Lambda_F$  of  $\mathcal{Q}_{\ell} = \rho^*(\mathcal{Q}_{\ell})$  on  $Z_{\ell}$  consisting of all surfaces containing the complete intersection curve  $F \cap \mathbb{D}_0^*$ . Imposing to the surfaces in  $\Lambda_F$  the condition to contain a general point of F, the divisor F + 2E splits off, and the residual surface sits in  $\mathcal{L}_{\ell}(1)$ . Hence  $\Lambda_F$  contains a codimension 1 sublinear system consisting of surfaces of the form 2E + F + L, with L varying in  $\mathcal{L}_{\ell}(1)$ , which has dimension 2. Hence  $\dim(\Lambda_F) = 3$ . Since  $\mathcal{B}$  dominates  $\mathcal{Q}_{\ell}$  via  $\pi_2$ , which is finite on  $\mathcal{B}$ , and  $\mathcal{B}_F$  is the inverse image of  $\Lambda_F$ , one has dim $(\mathcal{B}_F)$  =  $\dim(\Lambda_F) = 3$ . The proof is similar for  $\mathcal{B}_{F,\xi}$ . The final assertion follows by the genericity assumptions. 

We denote by  $T_0$  the tangent space to  $\mathcal{B}$  at  $b_0$ , and by  $T_F$  and  $T_{F,\xi}$  the 3 and 2-dimensional subspaces of  $T_0$  tangent to  $\mathcal{B}_F$  and to  $\mathcal{B}_{F,\xi}$  at  $b_0$ , respectively.

Lemma 3.7. One has:

- (a)  $\bigcup_{F \in \mathcal{L}} T_F$  spans  $T_0$ ; (b) given  $F \in \mathcal{L}$ , the union  $\bigcup_{\xi \in \Gamma_0^*} T_{F,\xi}$  spans  $T_F$ .

*Proof.* (a) Since  $\pi_2$  is étale on  $\mathcal{B}$ ,  $T_0$  is isomorphic to the tangent space to  $\mathcal{Q}_{\ell}$  at  $\mathbb{D}_0^*$ . Remember that, by §2.3, the double plane  $\mathbb{D}_0^*$ , considered in  $W_\ell$ , is cut out by a quadric with equation  $z^2 = Q(x^I)_{|I|=\ell}$ . So  $T_0$  can be identified with the vector space of homogeneous quadratic polynomials of the form  $az^2 - G(x^I)_{|I|=\ell}$  modulo the one-dimensional linear space spanned by  $z^2 - Q(x^I)_{|I|=\ell}$  and by the linear space of quadratic polynomials in  $\{x^I\}_{|I|=\ell}$  defining  $V_{\ell}$ . <sup>1</sup> Hence  $T_0$  can be identified with the vector space of quadratic polynomials in  $\{x^I\}_{|I|=\ell}$ , modulo the vector space of quadratic polynomials in  $\{x^I\}_{|I|=\ell}$  defining  $V_{\ell}$ . This, in turn, can be identified with the vector space  $S_d$  of homogeneous polynomials of degree d in  $x_0, x_1, x_2$ .

Now  $T_F$  can be identified with the vector subspace  $S_d(f)$  of  $T_0 \cong S_d$  of polynomials of the form fh, where f is a fixed homogeneous polynomial of degree d-1(determined by F), and h is any linear form. By assumption on  $\mathcal{L}, \bigcup_{F \in \mathcal{L}} T_F$  contains all monomials of degree d, which do span  $S_d$ .

(b) Given F,  $T_{F,\xi}$  can be identified with the vector space of homogeneous polynomials of the form fh, where h vanishes at  $\pi(\xi) \in \mathbb{P}^2$ . These polynomials do span  $T_F \cong S_d(f).$ 

Next we consider the restrictions

$$\mathcal{D}_F \xrightarrow{p} \mathcal{B}_F, \quad \mathfrak{C}_F \xrightarrow{\mathfrak{q}} \mathcal{B}_F, \text{ and } \mathcal{D}_{F,\xi} \xrightarrow{p} \mathcal{B}_{F,\xi}, \quad \mathfrak{C}_{F,\xi} \xrightarrow{\mathfrak{q}} \mathcal{B}_{F,\xi},$$

respectively, of the families

$$\mathcal{D} \xrightarrow{p} \mathcal{B} \text{ and } \mathfrak{C} \xrightarrow{\mathfrak{q}} \mathcal{B}$$

Proposition 3.8. For general  $F \in \mathcal{L}$  and  $\xi \in \Gamma_0^*$ , the families

$$\mathfrak{C}_F \xrightarrow{\mathfrak{q}} \mathcal{B}_F$$
 and  $\mathfrak{C}_{F,\xi} \xrightarrow{\mathfrak{q}} \mathcal{B}_{F,\xi}$ 

have the covering property.

<sup>&</sup>lt;sup>1</sup>An explanation is in order. Consider a vector space V and a nonzero vector  $v \in V$ , along with the associated projective space  $\mathbb{P}(V)$  and the corresponding point  $[v] \in \mathbb{P}(V)$ . Then the tangent space  $T_{[v]}\mathbb{P}(V)$  can be canonically identified with  $\operatorname{Hom}(\langle v \rangle, V/\langle v \rangle) \cong V/\langle v \rangle$ .

*Proof.* We prove the assertion for  $\mathfrak{C}_F \xrightarrow{\mathfrak{q}} \mathcal{B}_F$ . The proof for  $\mathfrak{C}_{F,\xi} \xrightarrow{\mathfrak{q}} \mathcal{B}_{F,\xi}$  is similar (and analogous to the proof of the corresponding statement in [21, Theorem (1.2)]), hence it can be left to the reader.

Let  $\mathbb{M}$  be the set of all monomials of degree d-1 in  $x_0, x_1, x_2$ . Consider the family  $\{F_M\}_{M \in \mathbb{M}}$ , where  $F_M \in \mathcal{L}$  is defined by the pull-back via  $\pi \colon \mathbb{Z}_{\ell} \to V_{\ell} \cong \mathbb{P}^2$ of the monomial M. Take two monomials M', M'' which differ only in degree 1, i.e., their lowest common multiple U has degree d. Then  $\mathcal{B}_{F_{M'}} \cap \mathcal{B}_{F_{M''}}$  contains the pull-back via  $\pi_2$  of an open, dense subset of the pencil  $\langle \mathbb{D}_t^* \rangle$  spanned by  $\mathbb{D}_0^*$ and  $F_U$ , where  $F_U$  is the pull-back via  $\pi$  of the monomial U. The base locus of this pencil does not contain  $\Gamma_0^*$ . Therefore,  $\Gamma_0^*$  varies in a non-trivial one-parameter family  $\langle \Gamma_t^* \rangle$  together with members  $\mathbb{D}_t^*$  varying in the pencil  $\langle \mathbb{D}_t^* \rangle$ .

Next we apply Lemma 3.5 with

- $V = T_0;$
- $W = H^0(\Gamma_0^*, N_{\Gamma_0^*|Z_\ell});$
- the linear map g induced by the characteristic map (see Proposition 1.1 (a));
- the family of subspaces  $\{V_i\}_{i \in I}$  given by  $\{T_{F_M}\}_{M \in \mathbb{M}}$ .

For each pair of monomials M', M'', there is a sequence of monomials  $M_1 = M', M_2, \ldots, M_{t-1}, M_t = M''$ , such that for all  $i = 1, \ldots, t-1$ , the lowest common multiple of  $M_i$  and  $M_{i+1}$  has degree d. The above argument implies that  $g(T_{F_{M_i}} \cap T_{F_{M_{i+1}}})$  has dimension at least 1, for all  $i = 1, \ldots, m-1$ . Furthermore, one has  $\dim(g(T_0)) \ge 2$ , because  $\mathfrak{C} \to \mathcal{B}$  is a covering family (see (b) of Proposition 1.1 and Lemma 3.4). By Lemma 3.5 there is a monomial  $M \in \mathbb{M}$  such that  $\dim(g(T_{F_M})) \ge 2$ ; by virtue of Lemma 3.6, this implies that  $\mathfrak{C}_{F_M} \to \mathfrak{B}_{F_M}$  is a covering family. This proves the assertion.

To finish the proof of Theorem 3.1 in this case, consider the covering family  $\mathfrak{C}_{F,\xi} \xrightarrow{\mathsf{q}} \mathcal{B}_{F,\xi}$ , with  $F \in \mathcal{L}$  and  $\xi \in \Gamma_0^*$  general. Using (1.5), (2.6), and (2.8), for  $b = (\Gamma_b^*, \mathbb{D}_b^*) \in \mathcal{B}_{F,\xi}$  general (see §3.1.1) we deduce

(3.2) 
$$\deg(\Phi_b^{\rm sm}) \leqslant 4(g-1) + \delta + 2m(\ell+3) = 4(g-1) + \delta + m(d+6).$$

On the other hand, by construction and by (a) of Proposition 1.3,

(3.3) 
$$\deg(\Phi_b^{\mathrm{sm}}) \ge 1 + \deg(\Gamma_b^* \cap F) = 1 + 2(d-1)m.$$

Comparing (3.2) and (3.3) gives (3.1).

3.2.2. The odd degree case. The proof runs exactly as in the case of even d, so we will be brief and leave the details to the reader.

Fix again  $b_0 \in \mathcal{B}$ ,  $\Gamma_0^*$  and  $\mathbb{D}_0^*$  as in the even degree case. Following what we did in §3.1.2, we replace  $D_b$  by  $D_b + h$ , where  $h \subset \mathbb{P}^2$  is a general line. In the present setting we let  $\mathcal{L}$  be the open subset of  $\mathcal{L}_{\ell}(d)$  consisting of the surfaces  $F \in \mathcal{L}_{\ell}(d)$ which do not contain  $\Gamma_0^*$ . Again we may assume that all surfaces F defined by the pull-back via  $\pi$  of degree d monomials in the variables  $x_0, x_1, x_2$  belong to  $\mathcal{L}$ . Given  $F \in \mathcal{L}$ , we define  $\mathcal{B}_F$  and  $\mathcal{B}_{F,\xi}$  as in the even degree case, and the analogue of Lemma 3.6 still holds. Then, with the usual meaning for  $T_0$ ,  $T_F$  and  $T_{F,\xi}$ , the analogue of Lemma 3.7 holds. Similarly as in Proposition 3.8, the covering property holds for the restricted families

$$\mathcal{D}_F \xrightarrow{p} \mathcal{B}_F, \quad \mathfrak{C}_F \xrightarrow{\mathfrak{q}} \mathcal{B}_F, \quad \text{and} \quad \mathcal{D}_{F,\xi} \xrightarrow{p} \mathcal{B}_{F,\xi}, \quad \mathfrak{C}_{F,\xi} \xrightarrow{\mathfrak{q}} \mathcal{B}_{F,\xi}.$$

We conclude finally as in the even degree case: (3.2) holds with no change, whereas (3.3) has to be replaced by

$$\deg(\Phi_b^{\rm sm}) \ge 1 + \deg(\Gamma_b \cap F) = 1 + 2dm,$$

and again, (3.1) follows.

## 4. Genera of subvarieties: a survey

As we mentioned in the Introduction, inequality (0.2) allows to bound the genera of curves in double planes from below. In this section we provide a brief survey on genera bounds for subvarieties in different type of varieties, and discuss several conjectures. All varieties are supposed to be projective, reduced, irreducible, and defined over  $\mathbb{C}$ . The geometric genus  $p_g(Y)$  of a variety Y is the geometric genus of a smooth model of Y.

Two important sources of interest in bounding genera are: the Clemens Conjecture on count of rational curves in Calabi-Yau varieties ([27]), and the celebrated Kobayashi Conjecture on hyperbolicity of general hypersurfaces in  $\mathbb{P}^n$  of sufficiently large degree ([59]). Recall ([43], [59]) that the Kobayashi hyperbolicity of a variety X implies the algebraic hyperbolicity, and in particular, absence of rational and elliptic curves in X. A part concerning the Clemens Conjecture started with the following theorem.

**Theorem 4.1.** (H. Clemens [26]) The geometric genera of curves in a very general hypersurface X of degree  $d \ge 2n-1$  in  $\mathbb{P}^n$  satisfy the inequality  $g \ge \frac{1}{2}(d-2n+1)+1$ .

This shows, in particular, the absence of rational curves in very general surfaces of degree  $d \ge 5$  in  $\mathbb{P}^3$ . One of the subsequent results in higher dimensions was

**Theorem 4.2.** (E. Ballico [3]) There is an effective bound  $\varphi(n)$  such that a very general hypersurface of degree  $d \ge \varphi(n)$  in  $\mathbb{P}^n$  is algebraically hyperbolic.

A better effective bound was provided by Geng Xu ([88]). For instance ([25]), it follows from the results in [88] that a general sextic threefold in  $\mathbb{P}^4$  is algebraically hyperbolic.

The Demailly algebraic hyperbolicity theorem states the following.

**Theorem 4.3.** (J.-P. Demailly [38]) For any hyperbolic subvariety  $X \subset \mathbb{P}^n$  there exists  $\varepsilon > 0$  such that, for any curve  $C \subset X$ , the geometric genus g of C is bounded below in terms of the degree:  $g \ge \varepsilon \deg(C) + 1$ . Therefore, the curves of bounded genera in X form bounded families.

Due to a recent proof of the Kobayashi Conjecture, Theorem 4.3 can be applied to general (in Zariski sense) hypersurfaces in  $\mathbb{P}^n$ .

**Theorem 4.4.** (D. Brotbek [12], Y.-T. Siu [79]) A general hypersurface of sufficiently large degree in  $\mathbb{P}^n$  is Kobayashi hyperbolic.

For effective estimates of degrees of hyperbolic hypersurfaces, see Y. Deng ([41, 42]); see also J.-P. Demailly [40] for a survey and a simplified proof.

L. Ein obtained some analogs of Clemens' estimate in higher dimensions.

**Theorem 4.5.** (L. Ein [44, 45]) Let M be a smooth projective variety of dimension  $n \ge 3$ , let  $L \to M$  be an ample line bundle, and let  $X \in |dL|$  be a very general member. Then for  $d \ge 2n - \ell$  any subvariety  $Y \subset X$  of dimension  $\ell$  has positive geometric genus, and for  $d \ge 2n - \ell + 1$ , Y is of general type.

In the case  $M = \mathbb{P}^n$  there is a sharper bound.

**Theorem 4.6.** (C. Voisin [81, 82]) Let X be a very general hypersurface of degree  $d \ge 2n - \ell - 1$  in  $\mathbb{P}^n$ ,  $n \ge 3$ . Then for  $d \ge 2n - \ell + 1$  any subvariety  $Y \subset X$  of dimension  $l \le n - 3$  has positive geometric genus, and for  $d \ge 2n - \ell$ , Y is of general type.

Sharper bounds are known also for certain toric varieties (A. Ikeda [53]). For  $M = \mathbb{P}^n$  a further improvement is due to G. Pacienza.

**Theorem 4.7.** (G. Pacienza [73]) For  $n \ge 6$  and for a very general hypersurface  $X \subset \mathbb{P}^n$  of degree  $d \ge 2n-2$ , any subvariety  $Y \subset X$  is of general type.

Geng Xu improved Ein's theorem as follows.

**Theorem 4.8.** (G. Xu [92]) Let X be a very general complete intersection of  $m \leq n-3$  hypersurfaces of degrees  $d_1, \ldots, d_m$  in  $\mathbb{P}^n$ , where  $d_i \geq 2 \forall i$ , and let  $Y \subset X$  be a reduced and irreducible divisor. Let  $d = d_1 + \cdots + d_m$ . Then  $p_g(Y) \geq n-1$  if  $d \geq n+2$ , and Y is of general type if d > n+2.

See also Geng Xu ([91]), C. Chang and Z. Ran ([15]), L. Chiantini, A.-F. Lopez, and Z. Ran ([22]), H. Clemens and Z. Ran ([28]), S.S.-T. Lu and Y. Miyaoka ([66]), and L.-C. Wang ([86, 87]). The results in [86] include some classes of divisors in Calabi-Yau hypersurfaces of degree d = n + 1 in  $\mathbb{P}^n$ .

Let us mention several sporadic results. See also, e.g., R. Beheshti ([6]), M. Bernardara ([8]), L. Bonavero and A. Hoering ([10]), T.D. Browning and P. Vishe ([13]), I. Coskun and E. Reidl ([31]), O. Debarre ([37]), K. Furukawa ([48, 49]), J. Harris, M. Roth, and J. Starr ([51]), J. Kollar ([60]). Concerning the Clemens Conjecture on rational curves in quintic threefolds and Mirrow Symmetry, see, e.g., M. Kontsevich ([61]), A. Libgober and J. Teitelbaum ([63]), D.A. Cox and S. Katz ([35]), T. Coates and A. Givental ([29]) and the references therein.

- **Theorem 4.9.** (G. Pacienza [72], E. Riedl and D. Yang [74]) Let  $X \subset \mathbb{P}^n$ be a very general hypersurface of degree d. If either n = 6 and d = 2n - 3, or  $n \ge 7$  and  $\frac{3n+1}{2} \le d \le 2n - 3$ , then X contains lines but no other rational curves.
  - (D. Shin [78]) A general hypersurface of degree d > <sup>3</sup>/<sub>2</sub>n − 1 in P<sup>n</sup> does not contain any smooth conic; however (S. Katz [56]), a general quintic threefold in P<sup>4</sup> does.
  - (S. Katz [56], P. Nijsse [71], T. Johnsen and S. L. Kleiman [54], J. D'Almeida [36], E. Cotterill [32, 33], E. Ballico and C. Fontanari [5]; cf. also E. Ballico [4] and A. L. Knutsen [57, 58]) A general quintic threefold X in P<sup>4</sup> contains only finitely many rational curves of degree ≤ 12, and each rational curve C of degree ≤ 11 either is smooth and embedded in X with a balanced normal bundle O(-1) ⊕ O(-1), or is a plane 6-nodal quintic.
  - (D. Shin [77], G. Mostad Hana and T. Johnsen [70], E. Cotterill [34]) A general hypersurface of degree 7 in  $\mathbb{P}^5$  does not contain any rational curve of degree  $d \in \{2, \ldots, 16\}$ .
  - (G. Ferrarese and D. Romagnoli [47]) The degree of an elliptic curve on a very general hypersurface X of degree 7 in P<sup>4</sup> is a multiple of 7.
  - (B. Wang [85]) A general hypersurface of degree 54 in P<sup>30</sup> does not contain any rational quartic curve.

• (B. Wang [84]) A very general hypersurface of degree  $d \ge 2n-1$  in  $\mathbb{P}^n$  does not contain any smooth elliptic curve.

For n = 3, Geng Xu replaced Clemens' initial genus bound in Theorem 4.1 by the optimal one. See also L. Chiantini and A.F. Lopez ([21]) for an alternative proof and some generalizations.

**Theorem 4.10.** (G. Xu [90]) The genera of curves on a very general surface of degree  $d \ge 5$  in  $\mathbb{P}^3$  satisfy the inequality  $g \ge \frac{1}{2}d(d-3)-2$ , and this bound is sharp. For  $d \ge 6$  this sharp bound can be achieved only by a tritangent hyperplane section.

Let Gaps(d) be the set of all non-negative integers which cannot be realized as geometric genera of irreducible curves on a very general surface of degree d in  $\mathbb{P}^3$ . This set is union of finitely many disjoint and separated integer intervals. By Xu's Theorem 4.10, the first gap interval is  $\text{Gaps}_0(d) = [0, d(d-3)/2 - 3]$ . For d = 5, this is the only gap interval. For  $d \ge 6$ , the next gap interval is  $\text{Gaps}_1(d) = [d(d - 3)/2 + 2, d^2 - 2d - 9]$  ([23]). One can show ([24]) that  $\max(\text{Gaps}(d)) = O(d^{8/3})$ . The latter is based on certain existence results. For arbitrary smooth (not necessarily general) surfaces in  $\mathbb{P}^3$ , we have the following existence result.

**Theorem 4.11.** ([16], [24]) There exists a function  $c(d) \sim d^3$  such that, for any smooth surface S in  $\mathbb{P}^3$  of degree d and any  $g \ge c(d)$ , S carries a reduced, irreducible nodal curve of geometric genus g, whose nodes can be prescribed generically on S.

To formulate an analog in higher dimensions, we recall the following notion. Let Y be an irreducible variety of dimension s. A singular point  $y \in Y$  is called an *ordinary singularity of multiplicity* m (m > 1), if the Zariski tangent space of Y at y has dimension s + 1, and the (affine) tangent cone to Y at y is a cone with vertex y over a smooth hypersurface of degree m in  $\mathbb{P}^s$ .

The next result was first established by J.A. Chen ([16]) for curves in *n*-dimensional varieties, and then in [24] for subvarieties of arbitrary dimension  $s \leq n-1^{2}$ .

**Theorem 4.12.** ([16], [24]) Let X be an irreducible, smooth, projective variety of dimension n > 1, let L be a very ample divisor on X, and let  $s \in \{1, ..., n - 1\}$ . Then there is an integer  $p_{X,L,s}$  such that for any  $p \ge p_{X,L,s}$  one can find an irreducible complete intersection  $Y = D_1 \cap ... \cap D_{n-s} \subset X$  of dimension s with at most ordinary points of multiplicity s + 1 as singularities such that  $p_g(Y) = p$ , where  $D_i \in |L|$  for i = 1, ..., n - s - 1 are smooth and transversal and  $D_{n-s} \in |mL|$ for some  $m \ge 1$ . Moreover, for  $n \ge 3$  and s = 1 one can find a smooth curve Y in X of a given genus  $g(Y) = p \ge p_{X,L,1}$ .

Recall the famous:

**Green-Griffiths-Lang Conjecture.** ([50, 62]; see also [7], [39]) Let X be a projective variety of general type. Then there exists a proper closed subset  $Z \subset X$  such that any subvariety  $Y \subset X$  not of general type is contained in Z.

The following conjecture is inspired by the previous results and by the Green-Griffiths-Lang Conjecture in the surface case.

**Conjecture.** There exists a strictly growing function  $\varphi(d)$ , with natural values, such that the set of curves of geometric genus  $g \leq \varphi(d)$  in any smooth surface S of degree  $d \geq 5$  in  $\mathbb{P}^3$  is finite.

 $<sup>^{2}</sup>$ We are grateful to J.A. Chen for pointing out his nice paper [16] that we ignored when writing [24]. We apologize for our ignorance.

Notice ([24]) that for any  $g \ge 0$  and  $d \ge 1$  one can find a smooth surface  $S \subset \mathbb{P}^3$  of degree d carrying a nodal curve of genus g. Notice also that any smooth quartic surface in  $\mathbb{P}^3$  contains an infinite countable set of rational curves, hence the restriction  $d \ge 5$  is necessary. Let us mention a few facts supporting the conjecture. According to B. Segre ([76]) the number of lines on a smooth surface of degree  $d \ge 3$  does not exceed (d-2)(11d-6). The celebrated Bogomolov theorem ([9]) says that the number of rational and elliptic curves on a surface of general type with  $c_1^2 > c_2$  is finite. Moreover, due to Y. Miyaoka, this number admits a uniform estimate:

**Theorem 4.13.** (Y. Miyaoka [68]) Let S be a minimal smooth projective surface of general type satisfying the inequality for Chern numbers  $c_1^2 > c_2$ . Then the number of irreducible curves of genus 0 and 1 on S is bounded by a function of  $c_1$  and  $c_2$ .

Analogous facts are true under certain weaker assumptions on Chern numbers (Y. Miyaoka [69]), or on the singularities of rational and elliptic curves in S (S.S.-Y. Lu and Y. Miyaoka [67]). It is plausible that the number of curves of genus  $g \leq \varphi(d)$  on a smooth surface of degree d in  $\mathbb{P}^3$  can be uniformly bounded above by a function of d. The conjecture above is coherent with the following ones.

**Conjecture.** (C. Voisin [83]) Let  $X \subset \mathbb{P}^n$  be a very general hypersurface of degree  $d \ge n+2$ . Then the degrees of rational curves in X are bounded.

**Conjecture.** (P. Autissier, A. Chambert-Loir, and C. Gasbarri [2]) Let X be a smooth projective variety of general type with the canonical line bundle  $K_X$ . Then there exist real numbers A and B, and a proper Zariski closed subset  $Z \subset X$  such that for any curve C of geometric genus g in X not contained in Z, one has  $\deg_C(K_X) \leq A(2g-2) + B$ .

#### References

- C. Araujo and J. Kollár. Rational curves on varieties. In: Higher dimensional varieties and rational points (Budapest, 2001), 13–68. Bolyai Soc. Math. Stud. 12, Springer, Berlin, 2003.
- [2] P. Autissier, A. Chambert-Loir, and C. Gasbarri. On the canonical degrees of curves in varieties of general type. GAFA (Geom. Funct. Anal.) 22 (2012), 1051–1061.
- [3] E. Ballico. Algebraic hyperbolicity of generic high degree hypersurfaces. Arch. Math. (Basel) 63 (1994), 282–283.
- [4] E. Ballico. Non-existence of smooth rational curves of degree d = 13, 14, 15 contained in a general quintic hypersurface of P<sup>4</sup> and in some quadric hypersurface. Note Mat. 36 (2016), 77–88.
- [5] E. Ballico and C. Fontanari. Finiteness of rational curves of degree 12 on a general quintic threefold. Pure Appl. Math. Q. 11 (2015), 537–557.
- [6] R. Beheshti. Hypersurfaces with too many rational curves. Math. Ann. 360 (2014), 753–768.
- [7] G. Berczi. Towards the Green-Griffiths-Lang Conjecture via equivariant localization. arXiv:1509.03406 (2015), 26p.
- [8] M. Bernardara. Calabi-Yau complete intersections with infinitely many lines. Rend. Sem. Mat. Univ. Pol. Torino 66 (2008), 1–11.
- [9] F.A. Bogomolov. Families of curves on a surface of general type. Dokl. Akad. Nauk SSSR (5)236 (1977), 1041–1044.
- [10] L. Bonavero and A. Hoering. Counting conics in complete intersections. Acta Math. Vietnam. 35 (2010), 23–30.
- [11] D. Brotbek. Hyperbolicity related problems for complete intersection varieties. Compos. Math. 150 (2014), 369–395.
- [12] D. Brotbek. On the hyperbolicity of general hypersurfaces. Publ. Math., Inst. Hautes Étud. Sci. 126 (2017), 1–34.
- [13] T.D. Browning and P. Vishe. Rational curves on smooth hypersurfaces of low degree. Algebra Number Theory 11, No. 7 (2017), 1657–1675.

- [14] J. Cao and A. Hoering. Rational curves on compact Kaehler manifolds. Avalaible at: http://math.unice.fr/~hoering/articles/a27-rat-curves.pdf
- [15] M.C. Chang and Z. Ran. Divisors on some generic hypersurfaces. J. Diff. Geom. 38 (1993), 671–678.
- [16] J. A. Chen. On genera of smooth curves in higher dimensional varieties. Proc. Amer. Math. Soc. 125 (1997), 2221–2225.
- [17] X. Chen. On the intersection of two plane curves. Math. Res. Lett. 7(5-6) (2000), 631-642.
- [18] X. Chen. On algebraic hyperbolicity of log varieties. Commun. Contemp. Math. 6 (2004), 513–559.
- [19] X. Chen and Y. Zhu. A<sup>1</sup>-curves on affine complete intersections. arXiv:1610.03791 (2016), 1−9.
- [20] L. Chiantini and C. Ciliberto. A few remarks on the lifting problem. Asterisque 218 (1993), 95–109.
- [21] L. Chiantini and A. F. Lopez. Focal loci of families and the genus of curves on surfaces. Proc. Amer. Math. Soc. 127 (1999), 3451–3459.
- [22] L. Chiantini, A.-F. Lopez, and Z. Ran. Subvarieties of generic hypersurfaces in any variety. Math. Proc. Cambr. Philos. Soc. 130 (2001), 259–268.
- [23] C. Ciliberto, F. Flamini, and M. Zaidenberg. Genera of curves on a very general surface in P<sup>3</sup>. Int. Math. Res. Not. 22 (2015), 12177–12205.
- [24] C. Ciliberto, F. Flamini, and M. Zaidenberg. Gaps for geometric genera. Arch. Math. 106 (2016), 531–541.
- [25] C. Ciliberto and M. Zaidenberg. Scrolls and hyperbolicity. Intern. J. Math. 24 (2013), 38–63.
- [26] H. Clemens. Curves on generic hypersurfaces. Ann. Sci. E.N.S. 19 (1986), 629–636.
- [27] H. Clemens. Curves on higher-dimensional complex projective manifolds. In: Proceedings of the International Congress of Mathematicians, vol. 1, 2 (Berkley, CA, 1986), 634–640, AMS, Providence RI, 1987.
- [28] H. Clemens and Z. Ran. Twisted genus bounds for subvarieties of generic hypersurfaces. Amer. J. Math. 126 (2004), 89–120; Erratum: Amer. J. Math. 127 (2005), 241–242.
- [29] T. Coates and A. Givental. Quantum Riemann-Roch, Lefschetz and Serre. Ann. of Math. 165 (2007), 15–53.
- [30] I. Coskun and J. Starr. Rational Curves on Smooth Cubic Hypersurfaces. Int. Math. Res. Not. 24 (2009), 4626–4641.
- [31] I. Coscun and E. Reidl. Normal bundles of rational curves on complete intersections. arXiv:1705.08441 (2017), 23p.
- [32] E. Cotterill. Rational curves of degree 10 on a general quintic threefold. Commun. Algebra 33 (2005), 1833–1872.
- [33] E. Cotterill. Rational curves of degree 11 on a general quintic 3-fold. Quart. J. Math. 63 (2012), 539–568.
- [34] E. Cotterill. Rational curves of degree 16 on a general heptic fourfold. J. Pure Appl. Algebra 218 (2014), 121–129.
- [35] D.A. Cox and S. Katz. Mirror Symmetry and Algebraic Geometry. Mathematical Surveys and Monographs 68. Amer. Math. Soc. 1999.
- [36] J. D'Almeida. Courbes rationnelles de degré 11 sur une hypersurface quintique générale de P<sup>4</sup>. Bull. Sci. Math. 136 (2012), 899–903.
- [37] O. Debarre. Rational curves on hypersurfaces. Lecture notes for the II Latin American School of Algebraic Geometry and Applications, 1–12 June 2015. Cabo Frio, Brazil. January 4, 2016, 48p.
- [38] J. P. Demailly. Algebraic criteria for Kobayashi hyperbolic projective varieties and jet differentials. In: Algebraic Geometry. Proceedings of the Summer Research Institute, eds. J. Kollàr et al., Santa Cruz, CA, USA, July 9–29, 1995, Proceeding of Symposia in Pure Mathematics, Vol. 62 (pt. 2) (American Mathematical Society, Providence, RI, 1997), 285–360.
- [39] J. P. Demailly. Towards the Green-Griffiths-Lang Conjecture. Baklouti, Ali (ed.) et al., Analysis and geometry. MIMS-GGTM, Tunis, Tunisia, March 24–27, 2014. Proceedings of the international conference. In honour of Mohammed Salah Baouendi. Springer Proceedings in Mathematics and Statistics 127, 141–159 (2015).
- [40] J. P. Demailly. Recent results on the Kobayashi and Green-Griffiths-Lang conjectures. arXiv:1801.04765 (2018), 76p.
- [41] Y. Deng. Effectivity in the hyperbolicity-related problems. arXiv:1606.03831 (2016), 13p.

- [42] Y. Deng. On the Diverio-Trapani conjecture. arXiv:1703.07560 (2017), 31p.
- [43] S. Diverio and E. Rousseau. Hyperbolicity of Projective Hypersurfaces. IMPA Monographs, vol. 5, Springer, Switzerland, 2016.
- [44] L. Ein. Subvarieties of generic complete intersections. Invent. Math. 94 (1988), 163–169.
- [45] L. Ein. Subvarieties of generic complete intersections. II. Math. Ann. 289 (1991), 465–471.
- [46] D. Eklund. Curves on Heisenberg invariant quartic surfaces in projective 3-space. arXiv:1010.4058v3 (2018), 19p.
- [47] G. Ferrarese and D. Romagnoli. Elliptic curves on the general hypersurface of degree 7 of P<sup>4</sup>. Indag. Math. 91 (1988), 249–252.
- [48] K. Furukawa. Rational curves on hypersurfaces. J. Reine Angew. Math. 665 (2012), 157–188.
- [49] K. Furukawa. Dimension of the space of conics on Fano hypersurfaces. arXiv:1702.08890 (2017), 14p.
- [50] M. Green and Ph. Griffiths. Two applications of algebraic geometry to entire holomorphic mappings. The Chern Symposium 1979, 41–74. Proc. Internal. Sympos. Berkeley, CA, 1979. Springer-Verlag, New York 1980.
- [51] J. Harris, M. Roth, and J. Starr. Rational curves on hypersurfaces of low degree. J. reine angew. Math. 571 (2004), 73–106.
- [52] J. Harris and J. Starr. Rational curves on hypersurfaces of low degree, II. Comp. Math. 141 (2005), 35–92.
- [53] A. Ikeda. Subvarieties of generic hypersurfaces in a nonsingular projective toric variety. Math. Z. 263 (2009), 923–937.
- [54] T. Johnsen and S. L. Kleiman. Rational curves of degree at most 9 on a general quintic threefold. Comm. Algebra 24 (1996), 2721–2753.
- [55] T. Johnsen and S. L. Kleiman. Toward Clemens' conjecture in degrees between 10 and 24. Serdica Math. J. 23 (1997), 131–142.
- [56] S. Katz. On the finiteness of rational curves on quintic threefolds. Comp. Math. 60 (1986), 151–162.
- [57] A. L. Knutsen. On isolated smooth curves of low genera in Calabi-Yau complete intersection threefolds. Trans. Amer. Math. Soc. 384 (2012), 5243–5284.
- [58] A. L. Knutsen. Smooth, isolated curves in families of Calabi-Yau threefolds in homogeneous spaces. J. Korean Math. Soc. 50 (2013), 1033–1050.
- [59] S. Kobayashi. Hyperbolic manifolds and holomorphic mappings: An introduction. 2d ed., World Scientific, 2005.
- [60] J. Kollár. Rational curves on algebraic varieties. Ergebnisse der Mathematik und ihrer grenzgebiete (3), vol. 32, (1996), Springer, Berlin, 1996.
- [61] M. Kontsevich. Enumeration of rational curves via torus actions. In: R.H. Dijkgraaf et al. (eds)., 335–369. The Moduli Space of Curves. Progress in Mathematics 129, Birkhauser, 1995.
- [62] S. Lang. Hyperbolic and Diophantine analysis. Bull. Amer. Math. Soc. 14 (1986), 159–205.
- [63] A. Libgober and J. Teitelbaum. Lines on Calabi-Yau complete intersections, mirror symmetry, and Picard-Fuchs equations. Int. Math. Res. Not. 1993:1 (1993), 29–39.
- [64] Y. Liu. Hyperbolicity of cyclic covers and complements. arXiv:1601.07514 (2016), 1–17. To appear in: Trans. Amer. Math. Soc., DOI: https://doi.org/10.1090/tran/7097.
- [65] Y. Liu. Construction of hyperbolic Horikawa surfaces. arXiv:1702.08143 (2017), 18p. To appear in: Ann. Inst. Fourier.
- [66] S.S.-Y. Lu and Y. Miyaoka. Bounding codimension-one subvarieties and a general inequality between Chern numbers. Amer. J. Math. 119 (1997), 487–502.
- [67] S.S.-Y. Lu and Y. Miyaoka. Bounding curves in algebraic surfaces by genus and Chern numbers. Math. Res. Lett. 2 (1995), 663–676.
- [68] Y. Miyaoka. The orbibundle Miyaoka-Yau-Sakai inequality and an effective Bogomolov-McQuillan Theorem. Publ. RIMS, Kyoto Univ. 44 (2008), 403–417.
- [69] Y. Miyaoka. Counting lines and conics on a surface. Publ. RIMS, Kyoto Univ. 45 (2009), 919–923.
- [70] G. Mostad Hana and T. Johnsen. Rational curves on a general heptic fourfold. Bull. Belg. Math. Soc. Simon Stevin 16 (2009), 861–885.
- [71] P. Nijsse. Clemens' conjecture for octic and nonic curves. Indag. Math. 6(2) (1995), 213–221.
- [72] G. Pacienza. Rational curves on general projective hypersurfaces. J. Alg. Geom. 12 (2003), 245–267.

- [73] G. Pacienza. Subvarieties of general type on a general projective hypersurface. Trans. Amer. Math. Soc. 356 (2004), 2649–2661.
- [74] E. Riedl and D. Yang. Rational curves on general type hypersurfaces. arXiv:1605.01759 (2016), 9p.
- [75] X. Roulleau and E. Rousseau. On the hyperbolicity of surfaces of general type with small c<sub>1</sub><sup>2</sup>.
  J. Lond. Math. Soc. (2) 87 (2013), 453–477.
- [76] B. Segre. The maximum number of lines lying on a quartic surface. Quart. J. Math., Oxford Ser. 14 (1943), 86–96.
- [77] D. Shin. Rational curves on general hypersurfaces of degree 7 in P<sup>5</sup>. Osaka J. Math. 44 (2007), 1–10.
- [78] D. Shin. Conics on a general hypersurface in complex projective spaces. Bull. Korean Math. Soc. 50 (2013), 2071–2077.
- [79] Y.-T. Siu. Hyperbolicity of generic high-degree hypersurfaces in complex projective space. Invent. Math. 202 (2015), 1069–1166.
- [80] B. Teissier. Résolution simultanée : I Familles de courbes. Séminaire sur les singularités des surfaces (1976-1977), 1-10.
- [81] C. Voisin. On a conjecture of Clemens on rational curves on hypersurfaces. J. Differential Geom. 44 (1996), 200–214.
- [82] C. Voisin. A correction on "A conjecture of Clemens on rational curves on hypersurfaces". J. Differential Geom. 49 (1999), 601–611.
- [83] C. Voisin. On some problems of Kobayashi and Lang; algebraic approaches. Current Developments in Mathematics 2003, 53–125.
- [84] B. Wang. Genus of curves in generic hypersurfaces. arXiv:1110.0185v3 (2011), 14p.
- [85] B. Wang. First order deformations of pairs and non-existence of rational curves. Rocky Mountain J. Math. 46 (2016), 663–678.
- [86] L.-C. Wang. A remark on divisors of Calabi-Yau hypersurfaces. Asian J. Math. 4 (2000), 369–372.
- [87] L.-C. Wang. Divisors of generic hypersurfaces of general type. Taiwanese J. Math. 6 (2002), 507–513.
- [88] G. Xu. On the complement of a generic curve in the projective plane. Amer. J. Math. 118(3) (1996), 611–620.
- [89] G. Xu. On the intersection of rational curves with cubic plane curves. J. Algebraic Geom. 7 (1998), 1–13.
- [90] G. Xu. Subvarieties of general hypersurfaces in projective space. J. Differential Geom. 39 (1994), 139–172.
- [91] G. Xu. Divisors on hypersurfaces. Math. Z. 219 (1995), 581–589.
- [92] G. Xu. Divisors on generic complete intersections in projective space. Trans. Amer. Math. Soc. 348 (1996), 2725–2736.

Dipartimento di Matematica, Università degli Studi di Roma "Tor Vergata", Via della Ricerca Scientifica, 00133 Roma, Italy

*E-mail address*: cilibert@mat.uniroma2.it

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DEGLI STUDI DI ROMA "TOR VERGATA", VIA DELLA RICERCA SCIENTIFICA, 00133 ROMA, ITALY

E-mail address: flamini@mat.uniroma2.it

Université Grenoble Alpes, Institut Fourier, CS 40700, 38058 Grenoble cedex 09, France

E-mail address: Mikhail.Zaidenberg@univ-grenoble-alpes.fr