

Fabio GAVARINI, Paolo PAPI

*“Representations of the Brauer algebra  
and Littlewood’s restriction rules”*

**ABSTRACT**

Let  $G$  be either  $Sp(V)$  or  $O(V)$ . Using an action of the Brauer algebra, we describe the subspace  $T^k(V^{\otimes m}) \subseteq V^{\otimes m}$  of tensors of valence  $k$  as an induced representation. As an application, we recover a special case of Littlewood’s restriction rule, affording the decomposition of an irreducible  $GL(V)$ –module when restricted to  $G$ . Moreover we get an explicit realization of the irreducible representations of the Brauer algebra.

— — — — —

**REFERENCES**

- [Br] R. Brauer, *On algebras which are concerned with semisimple continuous groups*, Annals of Mathematics **38** (1937), 854–872.
- [D-P] C. De Concini, C. Procesi, *A characteristic free approach to invariant theory*, Advances in Mathematics **21** (1976), 330–354.
- [D-S] C. De Concini, E. Strickland, *Traceless tensors and the symmetric group*, Journal of Algebra **61** (1979), 112–128.
- [K] S. V. Kerov, *Realizations of representations of the Brauer semigroup*, Journal of Soviet Mathematics **47** (1989), 2503–2507.
- [L] D. E. Littlewood, *On invariant theory under restricted groups*, Phil. Trans. Roy. Soc. A **239** (1944), 387–417.
- [L-P] J-L. Loday, C. Procesi, *Homology of symplectic and orthogonal algebras*, Advances in Mathematics **21** (1976), 93–108.
- [W] H. Weyl, *The classical groups*, Princeton University Press, New York, 1946.
- [Wz] H. Wenzl, *On the structure of Brauer’s centralizer algebras*, Annals of Mathematics **188** (1988), 173–193.