

# DECAY OF CORRELATIONS FOR PIECEWISE EXPANDING MAPS

CARLANGLO LIVERANI

University of Rome *Tor Vergata*

ABSTRACT. This paper investigates the decay of correlations in a large class of non-Markov one-dimensional expanding maps. The method employed is a special version of a general approach recently proposed by the author. Explicit bounds on the rate of decay of correlations are obtained.

## CONTENT

0. Introduction .....	p. 2
1. Operators and Invariant Cones .....	p. 3
2. The Map and an invariant cone .....	p. 5
3. Decay of correlations .....	p. 7
4. General results concerning expanding maps .....	p. 12
Appendix I (The constants and an example) .....	p. 15
Appendix II (The main inequality) .....	p. 17
References .....	p. 17

---

I would like to thank M. Blank, C. Falcolini and S. Vaienti for helpful discussions. In addition, it is a pleasure to acknowledge the hospitality of the Centre de Physique Théorique of Marseille where part of this work was done.

Typeset by  $\mathcal{A}\mathcal{M}\mathcal{S}$ -TEX

## §0 INTRODUCTION.

In this paper we apply a new technique—introduced in [L] to study the decay of correlations in hyperbolic systems—to one-dimensional non-Markov expansive maps (see section 2 for a precise definition). A main feature of such a method is the possibility of obtaining explicit bounds on the rate of decay.

The standard approach to this problem<sup>1</sup> is to study the Perron–Frobenius (P–F) operator via Ionescu-Tulcea and Marinescu type spectral theorems [ITM]. Unfortunately, this suffices to prove exponential decay of correlations but does not provide any constructive bound on the rate of decay. One can hope to obtain some bounds using the theory of Ruelle zeta functions [Ru1], [Ru2], [Ru3] but it is not clear in which generality this can be accomplished.

Here, we will study the P-F operator as well, but from a different point of view: we will see that there exists a convex cone of functions that is mapped “strictly” inside itself by the P-F operator. We will then be able to take advantage of an idea by Garrett Birkhoff: he showed that one can associate an Hilbert metric to the above mentioned cone and that such a metric is contracted by the P-F operator. Once we obtain a contraction all the wanted consequences are the result of straightforward arguments. This will allow to obtain an explicit bound for the rate of decay.

While several bounds on the rate of decay are available for smooth (or, more generally, Markov) expanding maps [F], [Ry], [H], very little is known in the discontinuous case (more precisely in the non-Markov case).

The possibility to obtain bounds on the rate of decay, for non-Markov maps, was

---

<sup>1</sup>See [LY] where it is proven the existence of an invariant measure absolutely continuous with respect to Lebesgue and [HK] where the decay of correlations is investigated.

recently investigated by P. Collet [Co]. He used techniques developed for the study of markov chains [DS]. Here we show that the approach proposed in [L] is more direct, produces substantially better bounds for the class of maps studied in [Co], and we apply it to a larger class of maps.

If  $T$  is a map of the type under consideration (see §2) and  $\mu$  is the associated invariant probability measure, absolutely continuous with respect to the Lebesgue measure (i.e., there exists  $\phi \in L^1([0, 1])$  such that  $d\mu = \phi dx$ ), then I prove the following:

**Theorem 0.1.** *If  $\inf_{x \in [0, 1]} \phi(x) \geq \gamma > 0$  and the system  $(T, \mu)$  is mixing, then there exists  $b, K > 0, \Lambda \in (0, 1)$ , “constructive in  $T$ ,” such that, for each function  $f \in L^1([0, 1])$  and  $g \in BV$  (the space of function of bounded variation),  $\int_0^1 g = 1$ ,*

$$\left| \int_0^1 gf \circ T^n - \int_0^1 f d\mu \right| \leq K \Lambda^{-n} \|f\|_1 \left( 1 + b \bigvee_0^1 g \right).$$

In addition, it is possible to state a concrete and checkable condition that insures  $\inf \phi \geq \gamma > 0$  and shows that it is a rather general feature of expanding maps (see Lemma 4.2). Of course, the novelty of Theorem 0.1 consists in “constructive in  $T$ ,” which means that, given  $T$ , it is possible to state explicit, although involved, formulas for  $b, K$  and  $\Lambda$ . Such formulas are summarized, for the reader convenience, in Appendix I. I do not claim that the explicit  $b, K$  and  $\Lambda$  I produce here are optimal (in fact, I believe they are not), more work in this direction is called for. Lastly, it may be interesting to notice that the bounds are robust: they vary continuously in the  $C^2$  topology.

The paper is organized as follows: in section 1 I provide a brief review of some basic facts on which the subsequent arguments rest. Section 2 contains a detailed description of the class of maps under consideration, and it presents the main ingre-

dient in the proof of Theorem 0.1: a cone of functions invariant under the action of the Perron-Frobenius operator. In section 3 the bound on the decay of correlations is proven, but under apparently stronger hypotheses than the ones in the statement of Theorem 0.1. In section 4 I show that the hypotheses of Theorem 0.1 imply the assumptions used in section 3, whereby concluding the argument. As a byproduct, I also answer to some questions, concerning general properties of expanding maps, recently raised by P. Collet [Co]. Finally, the paper includes two appendices; in the first I summarize the formulas for the constants  $b$ ,  $K$  and  $\Lambda$  and discuss a concrete example, obtaining numerical values for such constants (whereby emphasizing the constructive nature of the present approach). In the second is proven an helpful inequality.<sup>2</sup>

## §1 OPERATORS AND INVARIANT CONVEX CONES.

This section illustrates some results in lattice theory originally due to Garrett Birkhoff [B2]. More details and the proofs of the following results can be found in [L].

Consider a topological vector space  $\mathbb{V}$  with a partial ordering “ $\succeq$ ,” that is a vector lattice.<sup>3</sup> We require the partial order to be “continuous,” i.e. given  $\{f_n\} \in \mathbb{V}$   $\lim_{n \rightarrow \infty} f_n = f$ , if  $f_n \succeq g$  for each  $n$ , then  $f \succeq g$ . We call such vector lattices “integrally closed.”<sup>4</sup>

---

<sup>2</sup>The inequality is far from new. I prove it, to facilitate the reader, because the form I use is slightly non standard.

<sup>3</sup>We are assuming the partial order to be well behaved with respect to the algebraic structure: for each  $f, g \in \mathbb{V}$   $f \succeq g \iff f - g \succeq 0$ ; for each  $f \in \mathbb{V}$ ,  $\lambda \in \mathbb{R}^+ \setminus \{0\}$   $f \succeq 0 \implies \lambda f \succeq 0$ ; for each  $f \in \mathbb{V}$   $f \succeq 0$  and  $f \preceq 0$  imply  $f = 0$  (antisymmetry of the order relation).

<sup>4</sup>To be precise, in the literature “integrally closed” is used in a weaker sense. First,  $\mathbb{V}$  does not need a topology. Second, it suffices that for  $\{\alpha_n\} \in \mathbb{R}$ ,  $\alpha_n \rightarrow \alpha$ ;  $f, g \in \mathbb{V}$ , if  $\alpha_n f \succeq g$ , then



We define the closed convex cone<sup>5</sup>  $\mathcal{C} = \{f \in \mathbb{V} \mid f \neq 0, f \succeq 0\}$  (hereafter, the term “closed cone”  $\mathcal{C}$  will mean that  $\mathcal{C} \cup \{0\}$  is closed), and the equivalence relation “ $\sim$ ”:  $f \sim g$  iff there exists  $\lambda \in \mathbb{R}^+ \setminus \{0\}$  such that  $f = \lambda g$ . If we call  $\tilde{\mathcal{C}}$  the quotient of  $\mathcal{C}$  with respect to  $\sim$ , then  $\tilde{\mathcal{C}}$  is a closed convex set. Conversely, given a closed convex cone  $\mathcal{C} \subset \mathbb{V}$ , enjoying the property  $\mathcal{C} \cap -\mathcal{C} = \emptyset$ , we can define an order relation by

$$f \preceq g \iff g - f \in \mathcal{C} \cup \{0\}.$$

Henceforth, each time that we specify a convex cone we will assume the corresponding order relation and vice versa.

It is then possible to define a projective metric  $\Theta$  (Hilbert metric),<sup>6</sup> in  $\mathcal{C}$ , by the construction:

$$\begin{aligned} \alpha(f, g) &= \sup\{\lambda \in \mathbb{R}^+ \mid \lambda f \preceq g\} \\ \beta(f, g) &= \inf\{\mu \in \mathbb{R}^+ \mid g \preceq \mu f\} \\ \Theta(f, g) &= \log \left[ \frac{\beta(f, g)}{\alpha(f, g)} \right] \end{aligned}$$

where we take  $\alpha = 0$  and  $\beta = \infty$  if the corresponding sets are empty.

The importance of the previous constructions is due, in our context, to the following theorem.

**Theorem 1.1.** *Let  $\mathbb{V}_1$ , and  $\mathbb{V}_2$  be two integrally closed vector lattices;  $T : \mathbb{V}_1 \rightarrow \mathbb{V}_2$  a linear map such that  $T(\mathcal{C}_1) \subset \mathcal{C}_2$ , for the two corresponding closed convex cones  $\mathcal{C}_1 \subset \mathbb{V}_1$  and  $\mathcal{C}_2 \subset \mathbb{V}_2$ . Let  $\Theta_i$  be the Hilbert metric corresponding to the cone  $\mathcal{C}_i$ .*

---

$\alpha f \succeq g$ . Here we will ignore these and other subtleties: our task is limited to a brief account of the results relevant to the present context.

<sup>5</sup>Here, by “cone,” we mean any set such that, if  $f$  belongs to the set, then  $\lambda f$  belongs to it as well, for each  $\lambda > 0$ .

<sup>6</sup>In fact, we define a semi-metric, since  $f \sim g \Rightarrow \Theta(f, g) = 0$ . The metric that we describe corresponds to the conventional Hilbert metric on  $\tilde{\mathcal{C}}$ .

Setting  $\Delta = \sup_{f, g \in T(\mathcal{C}_1)} \Theta_2(f, g)$  we have

$$\Theta_2(Tf, Tg) \leq \tanh\left(\frac{\Delta}{4}\right) \Theta_1(f, g) \quad \forall f, g \in \mathcal{C}_1$$

( $\tanh(\infty) \equiv 1$ ).

**Remark 1.2.** *If  $T(\mathcal{C}_1) \subset \mathcal{C}_2$  then it follows that  $\Theta_2(Tf, Tg) \leq \Theta_1(f, g)$ . However, a uniform rate of contraction depends on the diameter of the image being finite.*

In particular, if an operator maps a convex cone strictly inside itself (in the sense that the diameter of the image is finite), then it is a contraction in the Hilbert metric. This implies the existence of a “positive” eigenfunction (provided the cone is complete with respect to the Hilbert metric), and, with some additional work, the existence of a gap in the spectrum of  $T$  (see [B1] for details). The relevance of this theorem for the study of invariant measures and their ergodic properties is obvious.

It is natural to wonder about the strength of the Hilbert metric compared to other, more usual, metrics. While, in general, the answer depends on the cone, it is nevertheless possible to state an interesting result.

**Lemma 1.3.** *Let  $\|\cdot\|$  be a norm on the vector lattice  $\mathbb{V}$ , and suppose that, for each  $f, g \in \mathbb{V}$ ,*

$$-f \preceq g \preceq f \implies \|f\| \geq \|g\|.$$

*Then, given  $f, g \in \mathcal{C} \subset \mathbb{V}$  for which  $\|f\| = \|g\|$ ,*

$$\|f - g\| \leq \left(e^{\Theta(f, g)} - 1\right) \|f\|.$$

## §2 THE MAP AND AN INVARIANT CONE.

We consider a map  $T$  from the interval  $[0, 1]$  into itself.

We assume that there exists a finite partition (mod-0, with respect to the Lebesgue measure)  $\mathcal{A}_0$  of  $[0, 1]$  into open intervals such that, for each interval  $I \in \mathcal{A}_0$ , the map  $T$ , restricted to  $I$ , can be extended to a  $C^2$  map on an open interval containing the closure of  $I$ . In addition, we assume that  $|DT| \geq \lambda > 1$  (expansivity).<sup>7</sup>

The simplest examples, in this class of maps, occur when, for each  $I \in \mathcal{A}_0$ ,  $TI$  is equal, mod-0, to the union of elements of  $\mathcal{A}_0$ . This case is called “Markov case,” and it is well understood in the literature; in particular, explicit bounds on the rate of decay of correlations are available [F], [Ry], [H], [L]. If the above mentioned property fails, the map is called “non-Markov,” this is the case addressed here.

Thanks to the work of Lasota and Yorke [LY] it is known that any piecewise smooth expanding map has at least one invariant measure absolutely continuous with respect to Lebesgue.<sup>8</sup> We will call  $\phi$  the density, with respect to the Lebesgue measure, of such an invariant measure  $\mu$ , and assume that the dynamical systems  $(T, \mu)$  is mixing.

The results of [HK] and [BK] imply that the class of maps under consideration exhibits exponential decay of correlations. But, no explicit bound on the rate of decay is available in such a generality.

A first technical obstacle is that the set  $\{x \in [0, 1] \mid \phi(x) = 0\}$  may have positive

---

<sup>7</sup>The requirement  $\mathcal{A}_0$  finite is not essential but certainly several more extra conditions on the map should be introduced if  $\mathcal{A}_0$  is countable. For example, the condition  $\inf_{I \in \mathcal{A}_0} |TI| > 0$  is necessary for the main inequality (2.1) to make sense and to hold. Moreover, additional hypotheses would be needed to prove Lemma 3.5.

<sup>8</sup>See [BK] for a generalization of such a result.

Lebesgue measure. This is a concrete possibility and it is very easy to construct examples with this property. The simplest case of this behavior occurs when there exists an attracting set; clearly such a set supports all the invariant measures in the class under consideration (i.e., measures that are absolutely continuous with respect to the Lebesgue measure). We will assume that this is not the case. In principle, such an assumption does not imply any loss of generality: we can always study the restriction of the map to an invariant set.<sup>9</sup> Yet, some more complex situations may be possible. For example,  $\phi(x) \neq 0$  for each  $x \in [1, 0]$  but  $\inf_{x \in [0, 1]} \phi(x) = 0$ . As we will see in Lemma 4.1,  $\inf \phi = 0$  implies that some sets are visited very seldom, this makes much harder any attempt to estimate the rate of decay. To avoid such complications let's further reduce the class of maps under consideration.

On the one hand, we assume that  $|DT| \geq \lambda > 2$  (here no loss of generality is implied: this condition can always be satisfied replacing  $T$  by an appropriate power).

On the other hand we ask that there exists  $\gamma > 0$  such that  $\inf_{x \in [0, 1]} \phi(x) \geq \gamma$ .

To study the decay of correlations it is useful to introduce the Perron-Frobenius (P-F) operator  $\tilde{T}$ . Such operator is defined by the relation<sup>10</sup>

$$\int gf \circ T = \int f \tilde{T}g$$

for each  $f \in L^\infty([0, 1])$  and  $g \in L^1([0, 1])$ . A direct computation shows that

$$\tilde{T}g(x) = \sum_{y \in T^{-1}(x)} g(y) |D_y T|^{-1}.$$

We will show that the P-F operator leaves invariant a cone of functions; the decay of the correlations will then follow from the theory discussed in §1.

---

<sup>9</sup>In fact, there is nothing sacred about the interval  $[0, 1]$ .

<sup>10</sup>If not otherwise stated, all the integrals are between 0 and 1, and taken with respect to the Lebesgue measure.

Let  $BV$  be the space of functions of bounded variation on  $[0, 1]$ . The cones that we will use in this paper are:

$$\mathcal{C}_a = \left\{ g \in BV \mid g(x) \not\equiv 0; g(x) \geq 0 \forall x \in [0, 1]; \bigvee_0^1 g \leq a \int_0^1 g \right\},$$

for  $a > 0$ .<sup>11</sup>

The main ingredient for studying the maps under consideration is the following inequality (due to Lasota and Yorke, but see Appendix II for details) [LY]: for each  $g \in BV$

$$(2.1) \quad \bigvee_0^1 \tilde{T}g \leq 2\lambda^{-1} \bigvee_0^1 g + A \int_0^1 |g|$$

where  $A = \sup_{\xi \in [0, 1]} \frac{|D_\xi^2 T|}{|D_\xi T|^2} + 2 \sup_{I \in \mathcal{A}_0} \frac{\sup_{\xi \in I} |D_\xi T|^{-1}}{|I|}$  ( $\mathcal{A}_0$  and  $\lambda$  are defined at the beginning of the section).<sup>12</sup> The relevance of the cones  $\mathcal{C}_a$  and the inequality (2.1) are exemplified by the following result.

**Lemma 2.1.** *For each  $a > \frac{A}{1-2\lambda^{-1}}$ , there exists  $\sigma < 1$  such that*

$$\tilde{T}\mathcal{C}_a \subset \mathcal{C}_{\sigma a}.$$

*Proof.* For each  $g \in \mathcal{C}_a$ , inequality (2.1) yields

$$\bigvee_0^1 \tilde{T}g \leq 2\lambda^{-1} \bigvee_0^1 g + A \int_0^1 g \leq (2\lambda^{-1}a + A) \int_0^1 g.$$

The result follows by choosing  $\sigma = 2\lambda^{-1} + Aa^{-1}$ , and noticing that  $\int g = \int \tilde{T}g$ .

□

---

<sup>11</sup>By  $\bigvee_0^1 g$  we mean the variation of the function  $g$  in the interval  $[0, 1]$ .

<sup>12</sup>The alert reader has certainly noticed that we have defined  $\tilde{T}$  only on  $L^1([0, 1])$ , to define it on  $BV$  it is necessary to specify the value of  $\tilde{T}^n g$  at all points, that is also at the boundaries of the intervals belonging to the partition  $\mathcal{A}_0$ . In fact, we want to define all the powers of  $\tilde{T}$  as well, so we may be in trouble at countably many points (all the preimages of the boundaries of the partition). We will not worry about such points since they can be consistently ignored (loosely speaking, the value of  $\tilde{T}g$  at such points can be defined by taking left and/or right limits—see [HK] or [BK] for details).

### §3 DECAY OF CORRELATIONS.

In the previous section we have found a cone of functions that is left invariant by the P-F operator. This it is not quite enough to obtain a contraction in the corresponding Hilbert metric: the diameter of the image must also be investigated. The nature of the problem is elucidated by the following Lemma.

**Lemma 3.1.** *Calling  $\Theta_a$  the Hilbert metric associated to the cone  $\mathcal{C}_a$ , for each  $\nu < 1$ , and  $g \in \mathcal{C}_{\nu a}$*

$$\Theta_a(g, 1) \leq \ln \left[ \frac{\max \left\{ (1 + \nu) \int_0^1 g; \sup_{x \in [0, 1]} g(x) \right\}}{\min \left\{ (1 - \nu) \int_0^1 g; \inf_{x \in [0, 1]} g(x) \right\}} \right].$$

*Proof.* We have to find the set of  $\lambda$  and  $\mu$  such that  $\lambda 1 \preceq g \preceq \mu 1$ . Let us start with the first inequality; it is satisfied iff  $g - \lambda \in \mathcal{C}_a$ , i.e.

$$\lambda \leq g(x) \quad \forall x \in [0, 1]$$

and

$$\lambda \leq \int_0^1 g - a^{-1} \bigvee_0^1 g.$$

Consequently,

$$\alpha = \sup \lambda = \min \left\{ \inf_{x \in [0, 1]} g(x); \int_0^1 g - a^{-1} \bigvee_0^1 g \right\},$$

see §1 for a definition of  $\alpha$  and  $\beta$  in the present context. Since  $g \in \mathcal{C}_{\nu a}$ , it follows

$$\int_0^1 g - a^{-1} \bigvee_0^1 g \geq (1 - \nu) \int_0^1 g,$$

that is  $\alpha \geq \min\{(1 - \nu) \int g, \inf g\}$ . Analogously, one can compute  $\beta \leq \max\{(1 + \nu) \int g, \sup g\}$ .  $\square$

Note that, up to now, we do not have any control on the inf of a function belonging to our cones, therefore the above Lemma shows not only that more work is needed, but also in which direction to concentrate our efforts.

The first step is to notice that, if a function belongs to the cone  $\mathcal{C}_a$ , then it cannot be small too often. This is made precise by the following.

**Lemma 3.2.** *Given a partition, mod-0,  $\mathcal{P}$  of  $[0, 1]$ , if each  $p \in \mathcal{P}$  is a connected interval with Lebesgue measure less than  $\frac{1}{2a}$  (that is  $|p| < \frac{1}{2a}$ ), then, for each  $g \in \mathcal{C}_a$ , there exists  $p_0 \in \mathcal{P}$  such that*

$$g(x) \geq \frac{1}{2} \int_0^1 g \quad \forall x \in p_0.$$

*Proof.* Consider the set

$$\mathcal{P}_- = \left\{ p \in \mathcal{P} \mid \exists x_p \in p : g(x_p) < \frac{1}{2} \int_0^1 g \right\}.$$

Clearly the Lemma is proven if we show that  $\mathcal{P}_- \neq \mathcal{P}$ . Let us suppose the contrary.

For

$$\int_p g \leq |p| \left( g(x_p) + \bigvee_p g \right) < \frac{|p|}{2} \int_0^1 g + \frac{1}{2a} \bigvee_p g,$$

and remembering that  $g \in \mathcal{C}_a$ , we have

$$\int_0^1 g < \frac{1}{2} \int_0^1 g + \frac{1}{2a} \bigvee_0^1 g \leq \int_0^1 g$$

which is a contradiction.  $\square$

To continue, we define a particular class of partitions:

**Definition 3.3.** *For each  $n \in \mathbb{N}$ ,*

$$\mathcal{A}_n = \bigvee_{j=0}^n T^{-j} \mathcal{A}_0.$$

It is immediately clear that  $T^{n+1}$  is monotone and smooth on each element of the partition  $\mathcal{A}_n$  (in fact, this could be used as an alternative definition of  $\mathcal{A}_n$ ); moreover,  $\mathcal{A}_n$  consists of intervals with Lebesgue measure smaller than  $\lambda^{-n}$ .

We can then choose  $n_0$  such that all the elements of  $\mathcal{A}_{n_0}$  have measure less than  $\frac{1}{2a}$  (for example  $n_0 = \lceil \frac{\ln 2a}{\ln \lambda} \rceil + 1$  would do).

**Definition 3.4.** *We call a map “covering” if for each  $n \in \mathbb{N}$  there exists  $N(n)$  such that, for each  $I \in \mathcal{A}_n$ ,*<sup>13</sup>

$$T^{N(n)}I = [0, 1].$$

The above property corresponds to condition (H2) in [Co]. The importance of the notion of “covering” is emphasized by the following Lemma.

**Lemma 3.5.** *If the map  $T$  is covering, then for each  $a > \frac{A}{1-2\lambda^{-1}}$  there exists  $\Delta > 0$  such that*

$$\text{diam}(\tilde{T}^{N(n_0)}\mathcal{C}_a) \leq \Delta < \infty.$$

*Proof.* Let  $g \in \mathcal{C}_a$ , then, according to Lemma 3.2, there exists  $I_0 \in \mathcal{A}_{n_0}$  such that  $g(x) \geq \frac{1}{2} \int_0^1 g$  for each  $x \in I_0$ . By the covering property, for each  $x \in [0, 1]$  (apart, at most, finitely many points) there exists  $y \in I_0$  such that  $T^{N(n_0)}y = x$ ; hence

$$\left(\tilde{T}^{N(n_0)}g\right)(x) = \sum_{y \in T^{-N(n_0)}x} g(y)|D_y T^{N(n_0)}|^{-1} \geq \frac{\int_0^1 g}{2\|DT\|_\infty^{N(n_0)}}.$$

Lemma 2.1 implies  $\tilde{T}^{N(n_0)}\mathcal{C}_a \subset \mathcal{C}_{\sigma_1 a}$  with<sup>14</sup>

$$\sigma_1 = (2\lambda^{-1})^{N(n_0)} + \frac{1 - (2\lambda^{-1})^{N(n_0)}}{1 - 2\lambda^{-1}} Aa^{-1}.$$

---

<sup>13</sup>We have already remarked that what happens at the boundaries of the partitions is immaterial. In the same vein, each time that we write an equality between sets we always mean it apart a finite number of points.

<sup>14</sup>To obtain the following formula it is enough to notice that, iterating (2.1), for each  $k > 0$ , holds  $\bigvee_0^1 \tilde{T}^k g \leq (2\lambda^{-1})^k \bigvee_0^1 g + \frac{1 - (2\lambda^{-1})^k}{1 - 2\lambda^{-1}} A \int_0^1 g$ ; apply then Lemma 2.1 directly to  $\tilde{T}^{N(n_0)}$ .



Let  $\delta(g) = \frac{\inf \tilde{T}^{N(n_0)} g}{\int g}$ , up to now we have seen that  $\delta(g) \geq (2\|DT\|^{N(n_0)})^{-1}$  for

all  $g \in \mathcal{C}_a$ . Using Lemma 3.1, we can then estimate

$$\begin{aligned} \text{diam} \left( \tilde{T}^{N(n_0)} \mathcal{C}_a \right) &\leq 2 \sup_{g \in \mathcal{C}_a} \ln \left[ \frac{\max \left\{ (1 + \sigma_1) \int g; \inf \tilde{T}^{N(n_0)} g + \bigvee_0^1 \tilde{T}^{N(n_0)} g \right\}}{\min \left\{ (1 - \sigma_1) \int g; \inf \tilde{T}^{N(n_0)} g \right\}} \right] \\ &\leq 2 \sup_{g \in \mathcal{C}_a} \ln \left[ \frac{\max \{ (1 + \sigma_1); \delta(g) + a\sigma_1 \}}{\min \{ (1 - \sigma_1); \delta(g) \}} \right] \\ &\leq 2 \ln \left[ \frac{\max \{ (1 + \sigma_1); 1 + a\sigma_1 \}}{\min \left\{ (1 - \sigma_1); (2\|DT\|_\infty^{N(n_0)})^{-1} \right\}} \right] \equiv \Delta. \end{aligned}$$

□

The above Lemma, together with the results of §1, implies exponential decay of the correlations for covering maps.

**Theorem 3.6.** *If  $T$  is covering, then, for each  $f \in L^1([0, 1])$  and  $g \in BV$ ,  $\int_0^1 g = 1$ ,*

$$\left| \int_0^1 g f \circ T^n - \int_0^1 f d\mu \right| \leq K_n \Lambda^n \|f\|_1 \left( 1 + b \bigvee_0^1 g \right) \quad \forall n \in \mathbb{N},$$

with  $\Lambda = \tanh(\frac{\Delta}{4})^{\frac{1}{N(n_0)}}$ ,  $K_n = e^{\Delta \Lambda^{n-N(n_0)}} \Lambda^{-N(n_0)} \Delta \|\phi\|_\infty$  and  $b = (a - B)^{-1}$  ( $B = \frac{A}{1-2\lambda-1}$ ).<sup>15</sup>

*Proof.* Choose  $a > \frac{A}{1-2\lambda-1}$  and consider  $g \in \mathcal{C}_a$  normalized so that  $\int_0^1 g = 1$  (i.e.,  $g$  can be thought as the density of a measure). Then,

$$\left| \int_0^1 f(\tilde{T}^n g - \phi) dx \right| \leq \|f\|_1 \left\| \frac{\tilde{T}^n g}{\phi} - 1 \right\|_\infty \|\phi\|_\infty.$$

**Lemma 3.7.** *If  $\mathcal{C}_+ = \{g \in BV \mid g(x) \geq 0, \forall x \in [0, 1]\}$ , and  $\Theta_+$  is the corresponding Hilbert metric, then, for each  $g_1, g_2 \in \mathcal{C}_a$ ,*

$$\Theta_+(g_1, g_2) \leq \Theta(g_1, g_2)$$

<sup>15</sup>Note that, since  $\Lambda < 1$ ,  $\lim_{n \rightarrow \infty} K_n = \Lambda^{-N(n_0)} \Delta \|\phi\|_\infty \leq \Lambda^{-N(n_0)} \Delta (1 + a)$ .

*Proof.* Since  $\mathcal{C}_+ \supset \mathcal{C}_a$ , the identity is a map from  $BV$  to itself that maps  $\mathcal{C}_a$  into  $\mathcal{C}_+$ . The result follows then from Theorem 1.1.  $\square$

A simple computation yields

$$\Theta_+(g_1, g_2) = \ln \sup_{x, y \in [0, 1]} \frac{g_1(x)g_2(y)}{g_1(y)g_2(x)}.$$

Using the previous facts, and the trivial equality

$$\frac{(\tilde{T}^n g)(x)}{\phi(x)} = \frac{\tilde{T}^n g(x)\phi(y)}{\tilde{T}^n g(y)\phi(x)} \frac{\tilde{T}^n g(y)}{\phi(y)},$$

we have

$$e^{-\Theta_+(\tilde{T}^n g, \phi)} \frac{\tilde{T}^n g(y)}{\phi(y)} \leq \frac{\tilde{T}^n g(x)}{\phi(x)} \leq e^{\Theta_+(\tilde{T}^n g, \phi)} \frac{\tilde{T}^n g(y)}{\phi(y)}$$

for each  $x, y \in [0, 1]$ . Because  $\int_0^1 (\tilde{T}^n g - \phi) = 0$ , there must exist  $y_n^+, y_n^- \in [0, 1]$  such that  $\tilde{T}^n g(y_n^-) \leq \phi(y_n^-)$  and  $\tilde{T}^n g(y_n^+) \geq \phi(y_n^+)$ . Using the previous inequalities, with  $y = y_n^-$  and  $y = y_n^+$ , respectively, we obtain, for each  $x \in [0, 1]$ ,

$$e^{-\Theta_+(\tilde{T}^n g, \phi)} \leq \frac{\tilde{T}^n g(x)}{\phi(x)} \leq e^{\Theta_+(\tilde{T}^n g, \phi)}$$

and

$$\left\| \frac{\tilde{T}^n g}{\phi} - 1 \right\|_{\infty} \leq e^{\Theta_+(\tilde{T}^n g, \phi)} - 1 \leq e^{\Theta(\tilde{T}^n g, \phi)} - 1.$$

According to Theorem 1.1 and Lemma 3.5

$$\begin{aligned} \Theta(\tilde{T}^n g, \phi) &\leq \Theta([\tilde{T}^{N(n_0)}]^{[\frac{n}{N(n_0)}]} g, [\tilde{T}^{N(n_0)}]^{[\frac{n}{N(n_0)}]} \phi) \\ &\leq \tanh\left(\frac{\Delta}{4}\right)^{[\frac{n}{N(n_0)}]-1} \Theta(\tilde{T}^{N(n_0)} g, \phi). \end{aligned}$$

Hence,

$$\left\| \frac{\tilde{T}^n g}{\phi} - 1 \right\|_{\infty} \leq e^{\Lambda^{n-N(n_0)} \Theta(\tilde{T}^{N(n_0)} g, \phi)} - 1 \leq e^{\Lambda^{n-N(n_0)} \Delta \Delta \Lambda^{n-N(n_0)}}.$$

This estimate shows that for each  $f \in L^1([0, 1])$ ,  $g \in \mathcal{C}_a$ ,  $n \in \mathbb{N}$

$$\left| \int_0^1 f \circ T^n g dx - \int_0^1 f \phi dx \right| \leq K_n \|f\|_1 \Lambda^n$$

$$\Lambda = \tanh \left( \frac{\Delta}{4} \right)^{\frac{1}{N(n_0)}}; \quad K_n = e^{\Lambda^n - N(n_0) \Delta} \Delta \Lambda^{-N(n_0)} \|\phi\|_\infty.$$

Let us now consider  $g \in BV$ ,  $g \geq 0$ , and  $\int_0^1 g = 1$ . If  $\int_0^1 g \leq a$ , we have the above estimate, otherwise we define  $g_\rho = (g + \rho\phi)(1 + \rho)^{-1}$ , then  $\int_0^1 g_\rho = 1$  and

$$\int_0^1 g_\rho = \left[ \int_0^1 g + \rho \int_0^1 \phi \right] (1 + \rho)^{-1}.$$

Iterating (2.1) one obtains  $\int_0^1 \phi \leq \frac{A}{1-2\lambda-1} = B$ , then

$$\int_0^1 g_\rho \leq \left[ \int_0^1 g + \rho B \right] (1 + \rho)^{-1}$$

Choosing,

$$\rho = \frac{\int_0^1 g - a}{a - B}$$

we have  $g_\rho \in \mathcal{C}_a$ ; hence, the decay of correlations for  $g_\rho$  implies the decay of correlations for  $g$ . The result for arbitrary  $g \in BV$  follows, since any function can be written as the difference of two positive functions.  $\square$

In the next section we will see that, provided  $\inf \phi > 0$ , all the mixing maps are covering, whereby completing the proof of Theorem 0.1.

#### §4 GENERAL PROPERTIES OF EXPANDING MAPS.

In this section we address some questions concerning piecewise expanding maps posed by P. Collet in [Co]. We will see that the property that some image of any interval covers all  $[0, 1]$ , is a quite general feature of mixing maps. This shows that the results obtained in the paper apply to a wide class of maps.

We start by giving a checkable criterion for the hypotheses of Theorem 0.1.

**Definition 4.1.** We call a map “weakly-covering” if there exists  $N_0 \in \mathbb{N}$  such that, for each  $I \in \mathcal{A}_0$ ,

$$\bigcup_{j=0}^{N_0} T^j I = [0, 1].$$

This is a weaker version of (H1) in [Co]. The next Lemma shows that weak-covering is all is needed to insure that the first of the hypotheses of Theorem 0.1 holds.

**Lemma 4.2.** *If a map is weakly-covering, then there exists  $\gamma > 0$  such that  $\inf \phi \geq \gamma$ .*

*Proof.* A consequence of weak-covering and expansivity is that the property of being weakly-covering does not depend substantially on the partition.

**Sub-Lemma 4.3.** *If a map is weakly-covering, then, for each  $n \in \mathbb{N}$ , there exists  $N_0(n) \in \mathbb{N}$  such that, for each  $I \in \mathcal{A}_n$ ,*

$$\bigcup_{j=0}^{N_0(n)} T^j I = [0, 1].$$

*Proof.* Let  $I \in \mathcal{A}_n$ , then  $T^n$  is smooth on  $I$ , accordingly  $|T^n I| \geq \lambda^n |I|$ . If  $T^n I$  covers an element of  $\mathcal{A}_0$  then the Lemma is proven; if not, since  $T^n I$  is connected, it can intersect at most two elements of  $\mathcal{A}_0$ , so it is naturally broken in, at most, two pieces, let  $I_1$  be the larger of the two, clearly  $|I_1| \geq \lambda^n |I|/2$ .

We can then carry out a recursive argument: consider  $TI_1$ ; by construction it is connected, intersect it with the elements of  $\mathcal{A}_0$ , either it will cover one element or it will be divided in, at most, two sub-intervals, call  $I_2$  the larger one; consider  $TI_2$ , and so on. It follows that  $|I_k| \geq \left(\frac{\lambda}{2}\right)^{k-1} |I_1|$ , which implies that, eventually,  $I_k$  will cover an element of  $\mathcal{A}_0$ , this is all is needed to prove the Lemma.  $\square$

By definition  $\tilde{T}\phi = \phi$ , in addition, by Lemma 3.2 there exists  $I_0 \in \mathcal{A}_{n_0}$  such that  $\phi(x) \geq \frac{1}{2}$  for each  $x \in I_0$ . By Sub-Lemma 4.3, for each  $x \in [0, 1]$  there exists  $j \leq N_0(n_0)$  and  $y_* \in I_0$  such that  $T^j y_* = x$ . Hence,

$$\phi(x) = \tilde{T}^j \phi(x) = \sum_{y \in T^{-j}x} \phi(y) |D_y T^j|^{-1} \geq \phi(y_*) \|DT\|_\infty^{-j} \geq \frac{1}{2} \|DT\|_\infty^{-N_0(n_0)}.$$

□

The main result of this section is contained in the following Theorem.

**Theorem 4.4.** *If an expanding map is mixing and  $\inf \phi \geq \gamma > 0$ , then it is covering.*

*Proof.* For each interval  $I \subset [0, 1]$ , define

$$\chi_I(x) = \begin{cases} \frac{1}{|I|} & \text{for } x \in I \\ 0 & \text{for } x \notin I \end{cases}$$

then,  $\bigvee_0^1 \chi_I \leq \frac{2}{|I|}$  and  $\int_0^1 \chi_I = 1$ . Given any two intervals  $I, I'$ , the mixing property implies

$$\lim_{n \rightarrow \infty} \int_0^1 \chi_{I'} \chi_I \circ T^n = \int_0^1 \chi_I \phi \geq \gamma.$$

Consider some  $n_1 \in \mathbb{N}$  (to be chosen later), there exists  $N_* \in \mathbb{N}$  such that, for each

$I, I' \in \mathcal{A}_{n_1}$

$$\int_0^1 \chi_I \tilde{T}^n \chi_{I'} \geq \frac{\gamma}{2} \quad \forall n \geq N_*.$$

Choose  $I_0 \in \mathcal{A}_{n_1}$ , consider  $\tilde{T}^n \chi_{I_0}$ , from (2.1) follows

$$\bigvee_0^1 \tilde{T}^n \chi_{I_0} \leq (2\lambda^{-1})^n \frac{2}{|I_0|} + A \sum_{i=0}^{n-1} (2\lambda^{-1})^i \int_0^1 \chi_{I_0} \leq (2\lambda^{-1})^n \frac{2}{|I_0|} + \frac{A}{1 - 2\lambda^{-1}}.$$

Let  $b_n = \inf_{I \in \mathcal{A}_n} |I|$ , choose  $N_1 \geq N_*$  such that

$$\frac{2(2\lambda^{-1})^{N_1}}{b_{n_1}} \leq \frac{A}{1 - 2\lambda^{-1}}.$$

Then,

$$\bigvee_0^1 \tilde{T}^n \chi_{I_0} \leq \frac{2A}{1-2\lambda^{-1}} \quad \forall n \geq N_1.$$

Consider the set  $\mathcal{B}_- = \{I \in \mathcal{A}_{n_1} \mid \text{exists } x \in I : \tilde{T}^{N_1} \chi_{I_0}(x) < \frac{\gamma}{4}\}$ , and let  $L = \#\mathcal{B}_-$ .<sup>16</sup>

First of all, for each  $I \in \mathcal{A}_{n_1}$  there exists  $y \in I$  such that  $\tilde{T}^{N_1} \chi_{I_0}(y) \geq \frac{\gamma}{2}$ , if not

$$\int_0^1 \chi_I \tilde{T}^{N_1} \chi_{I_0} < \frac{\gamma}{2} \int_0^1 \chi_I = \frac{\gamma}{2}$$

contrary to our assumptions on  $N_1$ .

Consequently, for each  $I \in \mathcal{B}_-$ ,

$$\bigvee_I \tilde{T}^{N_1} \chi_{I_0} \geq \frac{\gamma}{4},$$

which implies

$$L \leq \frac{8A}{(1-2\lambda^{-1})\gamma} \equiv L_0.$$

But, if  $\#\{T^{-n}x\} \leq L_0$  we have

$$\gamma \leq \phi(x) = \sum_{y \in T^{-n}x} \phi(y) |D_y T^n|^{-1} \leq L_0 \|\phi\|_\infty \lambda^{-n}$$

which shows that, for  $N_2 = \left\lceil \frac{\ln L_0 \|\phi\|_\infty \gamma^{-1}}{\ln \lambda} \right\rceil + 1$ ,  $\#\{T^{-N_2}x\} > L_0$ .

Choose  $n_1 = N_2$ ; since  $T^{-n_1}x$  has at most one point in each element of  $\mathcal{A}_{n_1}$  and  $\#\{T^{-n_1}x\} > L_0$ , it follows that there exists  $y \in T^{-n_1}x$  such that  $y \in I \notin \mathcal{B}_-$ ; hence  $T^{n_1+N_1}I_0 = [0, 1]$ .

The statement is then proven by using the same reasoning employed in Sub-Lemma 4.3.  $\square$

Summarizing, if a map is weakly-covering and mixing then, in force of Lemma 4.2, we have  $\gamma > 0$  and, by Theorem 4.4, the map is covering. Hence, we can

---

<sup>16</sup>By  $\#$  we mean the cardinality of a set.

prove the exponential decay of correlations thanks to Theorem 3.6. We have proved Theorem 0.1.

## APPENDIX I (The constants and an example).

The constants  $\Lambda$ ,  $K$  and  $b$  in Theorem 0.1 can be chosen as

$$\begin{aligned}\Lambda &= \left[ \frac{1 - e^{-\frac{\Delta}{2}}}{1 + e^{-\frac{\Delta}{2}}} \right]^{\frac{1}{N^*}} < 1 \\ K &= e^{\Lambda^{-N^*}} \Lambda^{-N^*} \Delta(1 + a) \\ b &= \frac{1 - 2\lambda^{-1}}{A}.\end{aligned}$$

In the formulas appear several quantities that must be defined:  $\lambda = \inf_{\xi \in [0, 1]} |D_\xi T|$  is the minimum expansion rate;

$$A = \sup_{\xi \in [0, 1]} \frac{|D_\xi^2 T|}{|D_\xi T|^2} + 2 \sup_{I \in \mathcal{A}_0} \frac{\sup_{\xi \in I} |D_\xi T|^{-1}}{|I|}$$

is the constant that appears in the main inequality (2.1);

$$\Delta = 2 \ln \left[ \left( 3 + (2\lambda^{-1})^{N^*} \right) \left( \sup_{\xi \in [0, 1]} |D_\xi T| \right)^{N^*} \right]$$

is an estimate of the diameter of the image of the cone. In addition,

$$a = \max \left\{ 1, \frac{2A}{1 - 2\lambda^{-1}} \right\}$$

is the parameter that fixes the choice of the cone.<sup>17</sup>

The last quantity that need to be defined is  $N^*$ . Unfortunately the choice of  $N^*$  is not so simple. Recall that  $\mathcal{A}_n$  is a the coarser partition, in intervals, of  $[0, 1]$  such

---

<sup>17</sup>The reader may have noticed that  $a$  is never explicitly chosen in the body of the paper. In fact, all the estimates obtained depend on the choice of  $a$ , therefore the best strategy would be to chose  $a$  last by optimizing the bounds. Such an approach would lead to very complicate formulas, to avoid this here I make some choice, neither the best nor the worst.

that  $T^{n+1}$  is  $C^2$  on each interval; then, for each  $n \in \mathbb{N}$ , the number  $N(n)$  is defined as the smallest integer for which

$$T^{N(n)}I = [0, 1] \quad \forall I \in \mathcal{A}_n.$$

We choose  $n_0 = \left\lceil \frac{\ln 2a}{\ln \lambda} \right\rceil + 1$  and define

$$N^* = \max \left\{ N(n_0), \left\lceil \frac{\ln 2}{\ln \lambda - \ln 2} \right\rceil + 1 \right\}.$$

In section four we give abstract conditions insuring that, for each  $n \in \mathbb{N}$ ,  $N(n)$  exists finite, but I do not know any general bounds on  $N(n)$  so, in a concrete example, one must construct the partition  $\mathcal{A}_{n_0}$  and iterate its elements to find the value of  $N^*$ .

The above formulas are not the sharpest general bounds that can be obtain from the results contained in the paper: they represent a compromise between reasonable bounds and reasonable formulas. Nevertheless, it is important to remark that given a specific example it is possible to improve the estimates by following step by step the construction carried out in section three.

To be more concrete let us look at an example.

### Example

Consider the piecewise linear map

$$T(x) = \begin{cases} \frac{9}{2} \left( \frac{1}{9} - x \right) & x \in \left( 0, \frac{1}{9} \right) \\ \frac{9}{2} \left( x - \frac{1}{9} \right) & x \in \left( \frac{1}{9}, \frac{3}{9} \right) \\ \frac{9}{2} \left( \frac{5}{9} - x \right) & x \in \left( \frac{3}{9}, \frac{5}{9} \right) \\ \frac{9}{2} \left( x - \frac{5}{9} \right) & x \in \left( \frac{5}{9}, \frac{7}{9} \right) \\ \frac{9}{2} (1 - x) & x \in \left( \frac{7}{9}, 1 \right) \end{cases}$$



The partition on which  $T$  is defined is:

$$\mathcal{A}_0 = \left\{ \left(0, \frac{1}{9}\right); \left(\frac{1}{9}, \frac{1}{3}\right); \left(\frac{1}{3}, \frac{5}{9}\right); \left(\frac{5}{9}, \frac{7}{9}\right); \left(\frac{7}{9}, 1\right) \right\}$$

The map satisfies our assumptions since  $|DT| = \frac{9}{2} > 2$  and it is easy to check that it is covering (remember that, by Lemma 4.2 and Theorem 4.4, it suffice to check that  $T$  is weakly-covering).

The smaller element of the partition  $\mathcal{A}_0$  has size  $\frac{1}{9}$ , accordingly  $A = 4$  (see (2.1)). The choice of  $a$  is subject to the constraint  $a > \frac{A}{1-2\lambda^{-1}} = \frac{36}{5}$  (see Lemma 2.1), while  $n_0$  must satisfy  $\sup_{I \in \mathcal{A}_{n_0}} |I| \leq \frac{1}{2a}$ , it is then immediately clear that we must choose, at least,  $n_0 = 1$ .

The partition  $\mathcal{A}_1$  is

$$\mathcal{A}_1 = \left\{ \left(0, \frac{1}{27}\right); \left(\frac{1}{27}, \frac{7}{81}\right); \left(\frac{7}{81}, \frac{1}{9}\right); \left(\frac{1}{9}, \frac{11}{81}\right); \left(\frac{11}{81}, \frac{15}{81}\right); \dots \right\}$$

Clearly  $\inf_{I \in \mathcal{A}_1} |I| = \frac{2}{81}$ .

All the elements of  $\mathcal{A}_1$  have as image an element of  $\mathcal{A}_0$  that it is mapped on all  $[0, 1]$  apart for the ones that are mapped on  $(0, \frac{1}{9})$  and for  $(0, \frac{1}{27})$ . A direct computation shows that  $T^3(0, \frac{1}{27}) = [0, 1]$  and  $T^2(0, \frac{1}{9}) = [0, 1]$ , so  $N(1) = 3$ .

If we choose  $a = 7.25$ , then  $\sigma_1 \leq .994$  (see Lemma 3.5 for a definition of  $\sigma_1$ ).<sup>18</sup> Since  $a\sigma_1 \geq 1 + \sigma_1$  and  $1 - \sigma_1 \geq \frac{1}{2} \left(\frac{2}{9}\right)^3$ , the formula in the proof of Lemma 3.5 yields

$$\frac{\Delta}{2} \leq \ln \left[ 1 + 2a\sigma_1 \|DT\|^{N(1)} \right] \leq \ln 1314$$

$$\Lambda = \tanh\left(\frac{\Delta}{4}\right)^{\frac{1}{3}} \approx 1 - \frac{2}{3}e^{-\frac{\Delta}{2}} \leq 1 - \frac{1}{1971}$$

---

<sup>18</sup>Note that the general formulas at the beginning of the section would lead to the choice  $a = \frac{72}{5}$  and  $n_0 = 3$ . Since  $N(3) = 5$  it would follow  $\Lambda \leq 1 - \frac{1}{14000}$ ; we will see that a more careful approach can improve the estimate by almost two orders of magnitude.

moreover,  $\lim_{n \rightarrow \infty} K_n \leq 140$  and  $b \leq .3$ .

The above estimates are probably off by a couple of orders of magnitude, but they should not be considered too unsatisfactory: using the only other known rigorous estimates, [Co], one would have been lead to consider at least  $N(20)$  obtaining an estimate of  $(1 - \Lambda)^{-1}$  wrong by at least 20 orders of magnitude. In addition, the present estimate can certainly be improved: first of all it is easy to check that for each  $x \in [0, 1]$   $\#\{T^{-3}x\} \geq 2$ , this allows immediately to divide by two the number inside the log (cf. the proof of Lemma 3.5); second, some advantages can be obtained by using a different partition (instead of the dynamical one) and a different number (instead than  $\frac{1}{2}$ ) in Lemma 3.2.<sup>19</sup> Moreover, the result can conceivably be improved by choosing a cone of functions better adapted to the particular example at hand.

However, the problem of finding optimal bounds it is a business all in itself and not the one we have been concerned with in this paper.

## APPENDIX II (The main inequality).

Given  $I \in \mathcal{A}_0$ ,  $I = [a(I), b(I)]$ , and letting  $\chi_I$  be the characteristic function of  $I$ , we have

$$\begin{aligned} \bigvee_0^1 \tilde{T}g\chi_I &= \bigvee_I g|DT|^{-1} + |g(a(I))||D_{a(I)}T|^{-1} + |g(b(I))||D_{b(I)}T|^{-1} \\ &\leq \bigvee_I g|DT|^{-1} + \sup_{\xi \in I} |D_\xi T|^{-1} \left( \bigvee_I g + 2 \inf_{\xi \in I} |g(\xi)| \right) \\ &\leq 2\lambda^{-1} \bigvee_I g + \left( \sup_{\xi \in I} \frac{|D_\xi^2 T|}{|D_\xi T|^2} + \frac{2}{|I|} \sup_{\xi \in I} |D_\xi T|^{-1} \right) \int_I g. \end{aligned}$$

Since  $\bigvee_0^1 \tilde{T}g \leq \sum_{I \in \mathcal{A}_0} \bigvee_0^1 \tilde{T}g\chi_I$ , inequality (2.1) follows.

---

<sup>19</sup>Using these two ideas (i.e., the partition  $\{(0, \frac{1}{9}); (\frac{1}{9}, \frac{2}{9}); (\frac{2}{9}, \frac{3}{9}); \dots\}$  and  $\frac{7}{36}$  instead than  $\frac{1}{2}$ , which allows to set  $n_0 = 0$ ) it is already possible to obtain the improved estimate  $\Lambda \leq 1 - \frac{1}{377}$ .

## REFERENCES

- [BK] V. Baladi, G. Keller, *Zeta functions and transfer operators for piecewise monotone transformations*, Communications in Mathematical Physics **127** (1990), 459–477.
- [BY] M. Benedicks, L-S. Young, *Absolutely continuous invariant measures and random perturbations for certain one-dimensional maps*, Ergodic Theory and Dynamical Systems **12** (1992), 13–37.
- [B1] Garret Birkhoff, *Extension of Jentzsch's Theorem*, Transaction American Mathematical Society **85** (1957), 219–227.
- [B2] Garret Birkhoff, *Lattice Theory*, **25** 3rd ed., American Mathematical Society Colloquium Publications, Providence, Rhode Island, 1967.
- [DS] P. Diaconis, D. Stroock, *Geometric bounds for eigenvalues of Markov chains*, Annals of Applied Probability **1** (1991), 36–61.
- [Co] Collet, *An estimate of the decay of correlations for mixing non Markov expanding maps of the interval*, preprint (1984).
- [F] P. Ferrero, *Contribution a la theorie des etats d'equilibre en mecanique statistique*, Theses (1981).
- [FS1] P. Ferrero, B. Schmitt, *Produits aléatoires d'opérateurs matrices de transfert*, Probability Theory and Related Fields **79** (1988), 227–248.
- [FS2] P. Ferrero, B. Schmitt, *On the rate of convergence for some limit ratio theorem related to endomorphisms with a non regular invariant density*, preprint.
- [FS2] P. Ferrero, B. Schmitt, *Ruelle's Perron-Frobenius Theorem and Projective Metrics*, Random Fields, Esztergom (Hungary), Colloquia Mathematica Societas János Bolyai **27** (1979).
- [HK] F. Hofbauer, G. Keller, *Ergodic properties of invariant measures for piecewise monotone transformations*, Math. Zeit. **180** (1982), 119–140.
- [H] B.R. Hunt, *Estimating invariant measures and Lyapunov exponents*, preprint..
- [Ke] G. Keller, *Un théorème de la limite centrale pour une classe de transformations monotones par morceaux*, Comptes Rendus de l'Académie des Sciences, Série A **291** (1980), 155–158.
- [ITM] C. Ionescu Tulcea et Marinescu, Annals of Mathematics **52** (1950), 140–147.
- [L] C. Liverani, *Decay of correlations*, preprint.
- [LY] A. Lasota, J. Yorke, *On existence of invariant measures for piecewise monotonic transformations*, Trans. Amer. Math. Soc. **186** (1973), 481–487.
- [Ru1] D. Ruelle, *Thermodynamic formalism*, Addison-Wesley, New York, 1978.
- [Ru2] D. Ruelle, *The Thermodynamic Formalism for Expanding Maps*, Communications in Mathematical Physics **125** (1989), 239–262.
- [Ru3] D. Ruelle, *An extension of the theory of fredholm determinants*, IHES **72** (1990), 175–193.
- [Ry] Marek Rychlik, *Regularity of the metric entropy for expanding maps*, Transactions of the American Mathematical Society **315**, **2** (1989), 833–847.
- [Sa] H. Samelson, *On the Perron-Frobenius Theorem*, Michigan Mathematical Journal **4** (1956), 57–59.
- [Y] L-S. Young, *Decay of correlations for certain quadratic maps*, Communications in Mathematical Physics **146** (1992), 123–138.
- [Z] K. Ziemian, *On the Perron-Frobenius operator and limit theorems for some maps of an interval*, Ergodic Theory and Related Topics II, Proceedings, Teubner Textezür Math. **94**, 1987, pp. 206–211.

LIVERANI CARLANGLO, MATHEMATICS DEPARTMENT, UNIVERSITY OF ROME II, TOR VERGATA, 00133 ROME, ITALY.

*E-mail address:* liverani@mat.utovrm.it