

CHAPTER 1

The origins: Differential equations



s this book is about Dynamical Systems, let's start by defining the object of study. The concept of Dynamical System is a very general one and it appears in many branches of mathematics from discrete mathematics, number theory, probability, geometry and analysis and has wide applications in physics, chemistry, biology, economy and social sciences.

Probably the most general formulation of such a concept is the action of a group over an algebra. Given a group \mathbb{G} and an algebra \mathcal{A} the (left)-action of \mathbb{G} on \mathcal{A} is simply a map $f : \mathbb{G} \times \mathcal{A} \rightarrow \mathcal{A}$ such that

1. $f(gh, a) = f(g, f(h, a))$ for each $g, h \in \mathbb{G}$ and $a \in \mathcal{A}$;
2. $f(e, a) = a$ for every $a \in \mathcal{A}$, where e is the identity element of \mathbb{G} ;
3. $f(g, a + b) = f(g, a) + f(g, b)$ for each $g \in \mathbb{G}$ and $a, b \in \mathcal{A}$;
4. $f(g, ab) = f(g, a)f(g, b)$ for each $g \in \mathbb{G}$ and $a, b \in \mathcal{A}$;

In our discussion we will be mainly motivated by physics. In fact, we will consider only the cases in which $\mathbb{G} \in \{\mathbb{N}, \mathbb{Z}, \mathbb{R}_+, \mathbb{R}\}$ ¹ is interpreted as *time* and \mathcal{A} is given by an algebra of functions over some set X , interpreted as the *observables* of the system.² In addition, we will

¹Although even in physics other possibilities are very relevant, e.g. in the case of Statistical Mechanics it is natural to consider the action of the space translations, i.e. the groups $\{\mathbb{Z}^d, \mathbb{R}^d\}$ for some $d \in \mathbb{N}$, $d > 1$.

²Again other possibilities are relevant, e.g. the case of Quantum Mechanics (in the so called Heisenberg picture) where the algebra of the observable is non commutative and consists of the bounded operators over some Hilbert space.

restrict ourselves to situations where the action over the algebra is induced by an action over the set X (this is a map $f : \mathbb{G} \times X \rightarrow X$ that satisfies condition 1, 2 above).³ Indeed, given an action f of \mathbb{G} on X and an algebra \mathcal{A} of function on X one can define $\tilde{f}(g, a)(x) := a(f(g, x))$ for all $g \in \mathbb{G}$, $a \in \mathcal{A}$ and $x \in X$. It is then easy to verify that \tilde{f} satisfies conditions 1-4 above.

We will call discrete time Dynamical System the ones in which $\mathbb{G} \in \{\mathbb{N}, \mathbb{Z}\}$ and continuous time Dynamical Systems the ones in which $\mathbb{G} \in \{\mathbb{R}_+, \mathbb{R}\}$. Note that, in the first case, $f(n, x) = f(n - 1 + 1, x) = f(1, f(n - 1, x))$, thus defining $T : X \rightarrow X$ as $T(x) = f(1, x)$, holds $f(n, x) = T^n(x)$.⁴ Thus in such a case we can (and will) specify the Dynamical System by writing only (X, T) . In the case of continuous Dynamical Systems we will write $\phi_t(x) := f(t, x)$ and call ϕ_t a flow (if the group is \mathbb{R}) or a semi-flow (if the group is \mathbb{R}_+) and will specify the Dynamical System by writing (X, ϕ_t) . In fact, in this notes we will be interested only in Dynamical Systems with more structure i.e. *topological*, *measurable* or *smooth* Dynamical Systems. By topological Dynamical Systems we mean a triplet (X, \mathcal{T}, T) , where \mathcal{T} is a topology and T is continuous (if $B \in \mathcal{T}$, then $T^{-1}B \in \mathcal{T}$). By smooth we consider the case in which X has a differentiable structure and T is r -times differentiable for some $r \in \mathbb{N}$. Finally, a measurable Dynamical Systems is a quadruple (X, Σ, T, μ) where Σ is a σ -algebra, T is measurable (if $B \in \Sigma$, then $T^{-1}B \in \Sigma$) and μ is an invariant measure (for all $B \in \Sigma$, $\mu(T^{-1}B) = \mu(B)$).⁵

So far for general definitions that, to be honest, are not so inspiring. Indeed, what characterizes the modern Dynamical Systems is not so much the setting but rather the type of questions that are asked, first and foremost:

- **Which behaviors are visible in nature?** (stability and bifurcation theory).
- **What happens for very long times?** (statistics and asymp-

³Again relevant cases are not included, for example all Markov Process where the evolution is given by the action of some semigroup.

⁴Obviously $T^2(x) = T \circ T(x) = T(T(x))$, $T^3(x) = T \circ T \circ T(x) = T(T(T(x)))$ and so on.

⁵The definitions for continuous Dynamical Systems are the same with $\{\phi_t\}$ taking the place of T .

otic theory)

The rest of this book will deal in various ways with such questions.

The original motivation for the above setting and for these questions comes from the study of the motion which, after Newton, typically appears as solution of an *ordinary differential equation* (ODE). It is then natural to start with a brief reminder of basic ODE theory.⁶

In section 1.1 I will recall the theorem of existence and uniqueness of the solutions of an ODE. In addition, I will state the Gronwall inequality, a very useful inequality for estimating the growth rate of the solution of an ODE. Finally, a theorem yielding the smooth dependence of the solutions of an ODE from an external parameter or from the initial conditions is provided.

In section 1.2 is given a very brief account of linear equations with constant coefficients (by discussing the exponential of a matrix) and of Floquet theory. That is the study of the solutions of a linear equation with coefficients varying periodically in time. The basic result being that the asymptotic properties of the solutions can be understood just by looking at the solutions after one period.

Finally, section 1.3 discusses the possibility of qualitative understanding the behavior of the solutions of ODE that cannot be solved explicitly (essentially all the ODEs). The arguments are very naive and are intended only to convince the reader that a) something can be done; b) a more sophisticated theory needs to be developed in order to have a satisfactory picture.

⁶In fact, also the solutions of a partial differential equation (PDE) may give rise to a Dynamical System, yet the corresponding theory is typically harder to investigate.

1.1 Few basic facts about ODE: a reminder

We will be mainly interested in the initial Cauchy problem for ODE. That is, given a separable Banach space \mathcal{B} ,⁷ $V \in \mathcal{C}_{\text{loc}}^0(\mathcal{B} \times \mathbb{R}, \mathcal{B})$,⁸ and $x_0 \in \mathcal{B}$, find an open interval $0 \ni I \subset \mathbb{R}$ and $x \in \mathcal{C}^1(I, \mathcal{B})$ such that

$$\begin{aligned}\dot{x}(t) &= V(x(t), t) \\ x(0) &= x_0.\end{aligned}\tag{1.1.1}$$

In the next chapters I will be mainly interested in the case $\mathcal{B} = \mathbb{R}^d$, for some $d \in \mathbb{N}$. Thus, the reader uncomfortable with Banach spaces can safely substitute \mathbb{R}^d to \mathcal{B} in all the subsequent arguments. Yet, it is interesting that the theory can be developed for general Banach spaces at no extra cost. For simplicity, in the following we will always assume that all the Banach spaces are separable even if not explicitly mentioned. In essence, this is just a fancy way of saying that much of the following depends only on the Banach structure of \mathbb{R}^d , that is, for example, nowhere is used the fact that \mathbb{R}^d has a finite basis.

On the contrary, I will consider ODE on manifolds. Yet, ODE on manifolds can always be reduced to the case \mathbb{R}^d case. I will briefly comment on this issue in section 1.1.5.

The first problem that comes to mind is the following

Question 1 *Does the Cauchy problem (1.1.1) always admit a solution? If there exists a solution is it unique?*

To address such an issue it is convenient to consider the equation⁹

$$x(t) = x_0 + \int_0^t V(x(s), s) ds \tag{1.1.2}$$

⁷A Banach spaces is a complete normed vector spaces. This means that a Banach space is a vector space V , over \mathbb{R} or \mathbb{C} , equipped with a norm $\|\cdot\|$ such that every Cauchy sequence in V has a limit in V . By *separable* we mean that there exists a countable dense set. Check [RS80, Kat66] for more details or [DS88] for a lot more details.

⁸Given two Banach spaces $\mathcal{B}_1, \mathcal{B}_2$, an open set $U \subset \mathcal{B}_1$, and $q \in \mathbb{N}$ by $\mathcal{C}^q(U, \mathcal{B}_2)$ we mean the functions from U to \mathcal{B}_2 that are q time (Fréchet) differentiable and the q -th differentials are continuous (see Problem 1.19 for a very quick discussion of differentiation in Banach spaces). Such a vector space is itself a Banach space when equipped with the norm given by the sup of all its derivatives till the order q included. By $\mathcal{C}_{\text{loc}}^q(U, \mathcal{B}_2)$ we mean the vector space of the functions $f : U \rightarrow \mathcal{B}_2$ such that, for each $R > 0$, $f \in \mathcal{C}^q(U \cap B(0, R), \mathcal{B}_2)$ where $B(0, R) = \{x \in \mathcal{B}_1 : \|x\| \leq R\}$.

⁹The most convenient meaning of the integral of a function with values in a

Problem 1.1 Show that for each open interval $0 \ni I \subset \mathbb{R}$, if $x \in \mathcal{C}^1(I, \mathcal{B})$ is a solution of (1.1.1), then it is a solution of (1.1.2). Show that if $x \in \mathcal{C}^0(I, \mathcal{B})$ is a solution of (1.1.2) then $x \in \mathcal{C}^1(I, \mathcal{B})$ and is a solution of (1.1.1).

1.1.1 Existence and uniqueness

The issue of existence and uniqueness of the solutions of (1.1.1) can be solved by applying the classical Banach fixed point Theorem (see A.1.1), provided we make a stronger assumption on V .

Theorem 1.1.1 (Existence and Uniqueness theorem for ODE)

For each $V \in \mathcal{C}_{loc}^1(\mathcal{B} \times \mathbb{R}, \mathcal{B})$ and $x_0 \in \mathcal{B}$ there exists $\delta \in \mathbb{R}_+$ such that there exists a unique solution of (1.1.1) in $\mathcal{C}^1((-\delta, \delta), \mathcal{B})$.

PROOF. Let $\delta \in (0, 1)$. The reader can verify that the vector space $\mathcal{C}^0([-\delta, \delta], \mathcal{B})$, equipped with the norm $\|u\|_\infty := \sup_{t \in [-\delta, \delta]} \|u(t)\|$ is a Banach space.¹⁰ For each $u \in \mathcal{C}^0([-\delta, \delta], \mathcal{B})$ let us define the operator $K : \mathcal{C}^0([-\delta, \delta], \mathcal{B}) \rightarrow \mathcal{C}^0([-\delta, \delta], \mathcal{B})$ by

$$K(u)(t) := x_0 + \int_0^t V(u(s), s) ds.$$

For $L = \max\{2\|x_0\|, 1\}$ let $M = \sup_{|t| \leq 1, \|u\| \leq L} \{\|V(u, t)\| + \|\partial_u V(u, t)\|\}$, then for each $u \in B_L := \{u \in \mathcal{C}^0([-\delta, \delta], \mathcal{B}) : \|u\|_\infty \leq L\}$ holds $\|K(u(t))\| \leq \|x_0\| + \delta M \leq L$ provided we chose $\delta \leq (L - \|x_0\|)M^{-1}$. In addition, for each $u, v \in B_L$,

$$\|K(u) - K(v)\|_\infty \leq \delta M \|u - v\|_\infty \leq \frac{1}{2} \|u - v\|_\infty,$$

provided we chose $\delta \leq (2M)^{-1}$. We can then apply Theorem A.1.1 and obtain a unique solution of the equation $Ku = u$ in B_L . This shows

Banach space is the *Bochner sense*, which reduces to the usual Lebesgue integral in the case $\mathcal{B} = \mathbb{R}^d$, see [Yos95] for definition and properties. Yet, for our purposes the equivalent of the Riemannian integral suffices and it is defined in the obvious manner. See Problem 1.21 for details.

¹⁰It suffices to remember that the uniform limit of continuous functions is a continuous function.

the existence and uniqueness of the solution of (1.1.2). The Theorem follows then by remembering Problem 1.1. \square

Remark 1.1.2 *Note that in the proof of Theorem A.1.1 one can chose the same δ for an open set of initial condition.*

From now on we will ignore the difference between \mathcal{C}^r and $\mathcal{C}_{\text{loc}}^r$ and relay on the understanding of the reader.

Remark 1.1.3 *The hypotheses of the above Theorem can be easily weakened to the case of V locally Lipschitz in x and continuous in t , yet only continuity does not suffice for uniqueness as shown by the example*

$$\begin{aligned}\dot{x} &= \sqrt{x} \\ x(0) &= 0.\end{aligned}$$

which has the solutions $x(t) = 0$ and $x(t) = \frac{1}{4}t^2$.¹¹

Remark 1.1.4 *The introduction of δ in Theorem A.1.1 cannot be avoided as shown by the example*

$$\begin{aligned}\dot{x} &= x^2 \\ x(0) &= 1.\end{aligned}$$

Its solution $x(t) = (1 - t)^{-1}$ is not continuous, nor bounded, for $t = 1$.

We have seen a mechanism whereby the solution cannot be defined for all times, the next Lemma shows that, for \mathcal{C}^1 vector fields is the only mechanism.¹²

Lemma 1.1.5 *In the hypotheses of Theorem 1.1.1, if $x \in \mathcal{C}^1([0, \delta), \mathcal{B})$ is a solution of (1.1.1) and if there exists $M > 0$ such that $\sup_{t \in [0, \delta)} \|x(t)\| \leq M$, then there exists $\bar{\delta} > \delta$ and $\bar{x} \in \mathcal{C}^1([0, \bar{\delta}), \mathcal{B})$ that solves (1.1.1) (i.e. the solution can be extended for longer times).*

¹¹If \mathcal{B} is finite dimensional, then $V \in \mathcal{C}^0$ suffices for the existence of a solution. This follows by a direct application of Schauder fixed point Theorem to (1.1.2). For informations on such a fixed point theorem and fixed point theorems in general see [Zei86].

¹²I state the result for positive times, for negative times is the same.

PROOF. Let $\{t_n\}$ any sequence that converges to δ , then

$$\|x(t_n) - x(t_m)\| \leq \int_{t_n}^{t_m} \|V(x(s), s)\| ds \leq |t_n - t_m| \sup_{\|z\| \leq M} \sup_{s \in [0, \delta]} \|V(z, s)\|.$$

Thus $\{x(t_n)\}$ is a Cauchy sequence and admits a limit $x_* \in \mathcal{B}$. By the same argument it follows $x_* = \lim_{t \rightarrow \delta} x(t)$. We can then consider the equation

$$y(t) = x_* + \int_0^t V(y(s), s + \delta) ds.$$

By Theorem 1.1.1 there exists δ_1 and $y \in \mathcal{C}^1((-\delta_1, \delta_1), \mathcal{B})$ which satisfy the above equation. Let then $\bar{\delta} = \delta + \delta_1$ and define

$$\bar{x}(t) := \begin{cases} x(t) & \text{for all } t \in [0, \delta) \\ y(t - \delta) & \text{for all } t \in [\delta, \bar{\delta}). \end{cases}$$

Clearly $\bar{x} \in \mathcal{C}^0([0, \bar{\delta}), \mathcal{B})$ and, for $t \in [\delta, \bar{\delta})$ holds true

$$\begin{aligned} \bar{x}(t) &= x_* + \int_{\delta}^t V(\bar{x}(s), s) ds = x_0 + \int_0^{\delta} V(\bar{x}(s), s) ds + \int_{\delta}^t V(\bar{x}(s), s) ds \\ &= x_0 + \int_0^t V(\bar{x}(s), s) ds. \end{aligned}$$

Thus, again remembering Problem 1.1, the Lemma follows. \square

1.1.2 Gronwald inequality

We have seen that the escape (growth) to infinity is the only obstruction to enlarging the domain of the solution.¹³ To understand a bit better how the solution of an ODE can grow in time we will consider a simple linear case. Before doing so we need a technical but extremely useful Lemma.

Lemma 1.1.6 (Integral Gronwald inequality) *Let $L, T \in \mathbb{R}_+$ and $\xi, f \in \mathcal{C}^0([0, T], \mathbb{R})$. If, for all $t \in [0, T]$,*

$$\xi(t) \leq L \int_0^t \xi(s) ds + f(t),$$

¹³Of course, this is the case only for regular vector fields. For other possibilities think of the case of collision among planets.

then

$$\xi(t) \leq f(t) + L \int_0^t e^{L(t-s)} f(s) ds.$$

PROOF. Let us first consider the case in which $f \equiv 0$. In this case the Lemma asserts $\xi(t) \leq 0$. Indeed, since ξ is a continuous function there exists $t_* \in [0, (2L)^{-1}] \cap [0, T] =: I_1$ such that $\xi(t_*) = \sup_{t \in I_1} \xi(t)$. But then,

$$\xi(t_*) \leq L \int_0^{t_*} \xi(s) ds \leq \xi(t_*) L t_* \leq \frac{1}{2} \xi(t_*)$$

which implies $\xi(t_*) \leq 0$ and hence $\xi(t) \leq 0$ for each $t \in I_1$. If $I_1 = [0, T]$, then we are done, otherwise letting $t_1 := (2L)^{-1}$ we have

$$\xi(t) \leq L \int_{t_1}^t \xi(s) ds$$

and we can make the same argument as before in the interval $[t_1, 2t_1]$. Iterating we have $\xi(t) \leq 0$ for all $t \in [0, T]$.

To treat the general case we reduce it to the previous one. Let

$$\zeta(t) := \xi(t) - f(t) - L \int_0^t e^{L(t-s)} f(s) ds.$$

Then

$$\begin{aligned} \zeta(t) &\leq L \int_0^t \xi(s) ds - \int_0^t L e^{L(t-s)} f(s) ds \\ &= L \int_0^t \zeta(s) ds + L \int_0^t \left\{ f(s) ds + L \int_0^s e^{L(s-\tau)} f(\tau) d\tau \right\} \\ &\quad - \int_0^t L e^{L(t-s)} f(s) ds. \end{aligned}$$

Next, notice that

$$\begin{aligned} \int_0^t ds L \int_0^s e^{L(s-\tau)} f(\tau) d\tau &= L \int_0^t d\tau f(\tau) \int_\tau^t ds e^{L(s-\tau)} \\ &= \int_0^t f(s) \{e^{L(t-s)} - 1\} ds. \end{aligned}$$

Thus,

$$\zeta(t) \leq L \int_0^t \zeta(s) ds.$$

We have then reduced the problem to the previous case which implies that it must be $\zeta(t) \leq 0$ from which the Lemma follows. \square

We can now discuss the announced examples. Let $L(\mathcal{B}, \mathcal{B})$ be the Banach space of the linear bounded operators from \mathcal{B} to \mathcal{B} .¹⁴

Lemma 1.1.7 *For each $A \in C^1(\mathbb{R}, L(\mathcal{B}, \mathcal{B}))$, consider the Cauchy problem*

$$\begin{aligned} \dot{x}(t) &= A(t)x(t) \\ x(0) &= x_0. \end{aligned}$$

If $\|A(t)\| \leq L$ for all $0 \leq t \leq T \in \mathbb{R}$, then $\|x(t)\| \leq e^{Lt}\|x_0\|$ for all $0 \leq t \leq T$.

PROOF. If we write the equation in the equivalent integral form we have

$$\|x(t)\| \leq \|x_0\| + \int_0^t \|A(s)x(s)\| ds \leq \|x_0\| + L \int_0^t \|x(s)\| ds.$$

Setting $\xi(t) := \|x(t)\|$ and applying Lemma 1.1.6 the Lemma follows. \square

Corollary 1.1.8 *If in the previous Lemma $\|A(t)\| \leq L$ for all $t \in \mathbb{R}$, then the solution is defined on all \mathbb{R} .*

Problem 1.2 *Explain why Lemma 1.1.7 does not apply to the following setting: $\mathcal{B} = C^1(\mathbb{R}^n, \mathbb{R})$ and*

$$\dot{x}(t, z) = \alpha(z)\partial_z x(t, z),$$

for some $\alpha \in C^1(\mathbb{R}^n, \mathbb{R})$. Compare with Problem 1.25.

¹⁴The norm of $L \in L(\mathcal{B}, \mathcal{B})$ is given by $\|L\| := \sup_{\substack{v \in \mathcal{B} \\ \|v\|=1}} \|Lv\|$. If $\mathcal{B} = \mathbb{R}^d$, then $L(\mathcal{B}, \mathcal{B})$ is just the vector space of the $d \times d$ matrices.

1.1.3 Flows

In this section we analyze the case in which the vector field is time independent and grows at most linearly.

Lemma 1.1.9 *Given $V \in \mathcal{C}^1(\mathcal{B}, \mathcal{B})$, if there exists $L, M \geq 0$ such that $\|V(x)\| \leq L\|x\| + M$, then the solution of (1.1.1) exists for all times and for all initial conditions.*

PROOF. We argue by contradiction. For each $x_0 \in \mathcal{B}$ let $I(x_0) = [0, \delta(x_0))$ be the maximal interval in which the solution is defined. If $\delta(x_0) < \infty$, then for each $t \leq \delta(x_0)$

$$\|x(t)\| \leq \|x_0\| + L \int_0^t \|x(s)\| ds.$$

Thus Gronwald inequality implies

$$\|x(t)\| \leq \|x_0\| + Mt + e^{Lt} \{\|x_0\| + ML^{-1}\}$$

for $t \in [0, \delta(x_0))$. Then, by Lemma 1.1.5, the solution can be extended, contrary to the assumption that $[0, \delta(x_0))$ was the maximal interval. \square

For each $x_0 \in \mathcal{B}$ and $t \in \mathbb{R}$ let $x(t, x_0)$ be the solution of (1.1.1) at time t .

Lemma 1.1.10 *For each V as in Lemma 1.1.9, setting $\phi_t(x_0) := x(t, x_0)$, $\phi_{-t} = \phi_t^{-1}$ for $t \geq 0$, we have that (\mathcal{B}, ϕ_t) , $t \in \mathbb{R}$, is a Dynamical System.*

PROOF. All we need to prove is that ϕ_t is an action of \mathbb{R} on \mathcal{B} . First of all note that ϕ_t is indeed invertible. If not then there would be $x, x' \in \mathcal{B}$ such that $\phi_t(x) = \phi_t(x')$. But then the uniqueness of the solutions of the ODE implies $x = x'$. Moreover it is easy to check that $\phi_{-t}(x_0) = x(-t, x_0)$. Finally, $\phi_t(\phi_s(x)) = \phi_{t+s}(x)$. \square

Remark 1.1.11 *We have thus proved that a large class of vector fields gives rise to flows.*

1.1.4 Dependence on a parameter

Having established the existence and uniqueness of the solution, the next natural questions present itself.

Question 2 *How do the solutions depend on the initial condition? How do the solutions depend on a change of the vector field?*

To discuss such issues it is convenient to first analyze first the second question. More precisely, given $V \in \mathcal{C}^2(\mathcal{B} \times \mathbb{R} \times \mathbb{R}^d, \mathcal{B})$ we consider the Cauchy problem

$$\begin{aligned}\dot{x}(t) &= V(x(t), t, \lambda) \\ x(0) &= x_0.\end{aligned}\tag{1.1.3}$$

Clearly the solution will depend on the parameter λ . The question is then: calling $x(t, \lambda)$ the solution of (1.1.3), for a given $t \in \mathbb{R}$ what can we say about the function $x(t, \cdot)$?

A first, rough, information follows by a general results extending Theorem A.1.1.

Theorem 1.1.12 *Given Banach spaces $\mathcal{B}, \mathcal{B}_1$, a bounded closed set $A \subset \mathcal{B}$, an open set $U \in \mathcal{B}_1$ and a map $K : A \times U \rightarrow \mathcal{B}$. If*

- i) $K(A, \lambda) \subset A$, for each $\lambda \in U$;*
- ii) there exists $\sigma \in (0, 1)$ such that $\|K(v, \lambda) - K(w, \lambda)\| \leq \sigma \|v - w\|$ for each $v, w \in A$, $\lambda \in U$;*
- iii) there exists $L \geq 0$ such that $\|K(v, \lambda) - K(v, \lambda')\| \leq L \|\lambda - \lambda'\|$ for each $v \in A$, $\lambda, \lambda' \in U$;*

then for each $\lambda \in U$, there exists a unique $v(\lambda) \in A$ such that $K(v(\lambda), \lambda) = v(\lambda)$ and, for each $\lambda, \lambda' \in U$

$$\|v(\lambda) - v(\lambda')\| \leq (1 - \sigma)^{-1} L \|\lambda - \lambda'\|.\tag{1.1.4}$$

PROOF. The existence of $v(\lambda)$ follows by Theorem A.1.1, it thus suffices to prove (1.1.4).

$$\begin{aligned}\|v(\lambda) - v(\lambda')\| &= \|K(v(\lambda), \lambda) - K(v(\lambda'), \lambda')\| \\ &\leq \|K(v(\lambda), \lambda) - K(v(\lambda'), \lambda)\| + L \|\lambda - \lambda'\| \\ &\leq \sigma \|v(\lambda) - v(\lambda')\| + L \|\lambda - \lambda'\|\end{aligned}$$

which readily implies the result. \square

Problem 1.3 Define $K(u, \lambda)(t) = x_0 + \int_0^t V(u(s), s, \lambda) ds$, and apply the above Theorem to show that, for t small enough, the solutions $x(t, \cdot)$ of problem (1.1.3) are Lipschitz.

In fact, much more holds true. As an example of what can be done I provide a theorem which hypotheses are not optimal but suffice for all our future needs.

Theorem 1.1.13 (Smooth dependence on a parameter) *Given two Banach spaces $\mathcal{B}, \mathcal{B}_1$, let $V \in \mathcal{C}^2(\mathcal{B} \times \mathbb{R} \times \mathcal{B}_1, \mathcal{B})$, $A \subset \mathcal{B}$ and $U \subset \mathcal{B}_1$ open sets so that Theorem 1.1.1 applies for some fixed $\delta > 0$ for all $x_0 \in A$ and $\lambda \in U$. Let $X(t, x_0, \lambda)$ be the unique solution of (1.1.3). Then, for each $(x_0, t) \in A \times (-\delta, \delta)$, $X(t, x_0, \cdot) \in \mathcal{C}^1(U, \mathcal{B})$.*

PROOF. For each $x_0 \in A$ let us consider the ODE in $L(\mathcal{B}_1, \mathcal{B})$

$$\begin{aligned} \dot{\xi}(t) &= \partial_x V(X(t, x_0, \lambda), t, \lambda) \cdot \xi(t) + \partial_\lambda V(X(t, x_0, \lambda), t, \lambda) \\ \xi(0) &= 0. \end{aligned} \quad (1.1.5)$$

We claim that $\xi(t) = \partial_\lambda X(t, x_0, \lambda)$. To verify it suffices to prove that there exists $C > 0$ such that, for $h \in \mathcal{B}$ small enough, if $\zeta(t, h, \lambda) := X(t, x_0, \lambda + h) - X(t, x_0, \lambda) - \xi(t)h$, then $\|\zeta(t, h)\| \leq C\|h\|^2$. By Taylor formula we have

$$\begin{aligned} \dot{\zeta}(t, h) &= V(X(t, x_0, \lambda + h), t, \lambda + h) - V(X(t, x_0, \lambda), t, \lambda) \\ &\quad - \partial_x V(X(t, x_0, \lambda), t) \cdot \xi(t)h - \partial_\lambda V(X(t, x_0, \lambda), t, \lambda)h \\ &= \partial_x V(X(t, x_0, \lambda), t) \cdot \zeta(t, h) + R(t) \end{aligned} \quad (1.1.6)$$

where

$$\begin{aligned} \|R(t)\| &\leq C (\|X(t, x_0, \lambda + h) - X(t, x_0, \lambda)\|^2 + \|h\|^2) \\ &\leq 2C(\|\zeta(t, h)\|^2 + (1 + \|\xi(t)\|^2)\|h\|^2). \end{aligned}$$

with $C = \|V\|_{\mathcal{C}^2}$. Note that $\zeta(0) = 0$. We can then conclude by using Lemma 1.1.6. Indeed such a Lemma applied to (1.1.5) implies $\|\xi(t)\| \leq e^{C_1 t}$, for some $C_1 > 0$. Next, let $T > 0$ be the maximal time such that $\|\zeta(t, h)\| \leq 1/2$ and $e^{2C_1 T} \leq 2$. Then, for $t \leq T$, (1.1.6) yields

$$\|\zeta(t, h)\| \leq \int_0^t 2C\|\zeta(s)\| ds + 3\|h\|^2$$

and Lemma 1.1.6 implies the announced estimate. \square

Problem 1.4 Prove Theorem 1.1.13 under the weaker assumption $V \in \mathcal{C}^1$.

Corollary 1.1.14 (Smooth dependence on initial conditions) Let $V \in \mathcal{C}^2(\mathcal{B} \times \mathbb{R}, \mathcal{B})$, and let δ so that Theorem A.1.1 applies for all x_0 in some open set U . For each $x_0 \in U$ let $X(t, x_0)$ be the unique solution of (1.1.1), then, for each $t \in (-\delta, \delta)$, $X(t, \cdot) \in \mathcal{C}^1(U, \mathcal{B})$ and $\xi = \partial_{x_0} X$ is a solution of

$$\begin{aligned}\dot{\xi}(t) &= \partial_x V(X(t, x_0), t) \cdot \xi(t) \\ \xi(0) &= \mathbb{1}.\end{aligned}\tag{1.1.7}$$

PROOF. Set $z = x - x_0$ and consider the resulting equation

$$\begin{aligned}\dot{z} &= V(z + x_0, t) =: \bar{V}(z, t, x_0) \\ z(0) &= 0.\end{aligned}$$

One can then consider x_0 as an external parameter, applying Theorem 1.1.13 yields the result. \square

1.1.5 ODE on Manifolds—few words

Let us remind that a *topological manifold* is a second countable Hausdorff space which is locally homeomorphic to Euclidean space. A *chart* over a topological manifold M is a pair (U, ϕ) such that $U \subset M$ is an open set and $\phi : U \rightarrow \mathbb{R}^n$, for some $n \in \mathbb{N}$, is a homeomorphism of U and the open set $\Phi(U)$. An *atlas* on a topological manifold is a countable collection of charts $\{(U_\alpha, \phi_\alpha)\}$. We say that the atlas is \mathcal{C}^k if $\phi_\alpha \circ \phi_\beta^{-1}$ is \mathcal{C}^k when is defined. We say that two \mathcal{C}^k atlas are equivalent if their union is a \mathcal{C}^k atlas. A \mathcal{C}^k manifold is a topological manifold equipped with an equivalence class of \mathcal{C}^k atlas (often called a *differentiable structure*).

Although most often we will be concerned with manifolds embedded in some \mathbb{R}^d , also other possibilities will be relevant. Let us consider two examples.

Problem 1.5 Show that \mathbb{R}^d is a \mathcal{C}^∞ manifold.¹⁵

¹⁵ Note that, contrary to \mathcal{C}^k , \mathcal{C}^∞ is not a Banach space (there is no good norm). It is possible to give to it the structure of a Fréchet space [RS80], but we will refrain from such subtleties. We just consider $\mathcal{C}^\infty = \bigcap_{n \in \mathbb{N}} \mathcal{C}^n$ as a vector space.

Problem 1.6 Let $f \in \mathcal{C}^k(\mathbb{R}^d, \mathbb{R})$, and consider $M = \{(x, y) \in \mathbb{R}^d \times \mathbb{R} : y = f(x)\}$. Consider the atlas consisting of the chart (M, ϕ) where $\phi(x, y) = x$. This is a \mathcal{C}^∞ manifold.

Problem 1.7 Check that $\mathbb{T}^d = \mathbb{R}^d / \mathbb{Z}^d$ is a \mathcal{C}^∞ manifold.

Given two differentiable manifolds (a \mathcal{C}^k manifold with $k \geq 1$) M_1, M_2 and a map $f : M_1 \rightarrow M_2$ we say that $f \in \mathcal{C}^r(M_1, M_2)$, $r \leq k$, if for each atlas $\{(U_\alpha, \phi_\alpha)\}$ of M_1 and atlas $\{(V_\beta, \psi_\beta)\}$ of M_2 , holds true $\psi_\beta \circ f \circ \phi_\alpha^{-1} \in \mathcal{C}^r$ on their domain of definition.

Given a differentiable manifold M and $x \in M$, we say that two curves $\gamma_1, \gamma_2 \in \mathcal{C}^1((-1, 1), M)$, such that $\gamma_1(0) = \gamma_2(0) = x$, are equivalent at x if for each chart (U, ϕ) such that $x \in U$ holds true $(\phi \circ \gamma_1)'(0) = (\phi \circ \gamma_2)'(0)$. A *tangent vector* at x is an equivalence class of curves.

Problem 1.8 Show that if M is locally homeomorphic to \mathbb{R}^d , then the set of tangent vectors at any $x \in M$ form canonically a d dimensional vector space.¹⁶

We will use $\mathcal{T}_x M$ to designate the *tangent space* at x , that is the set of the tangent vectors at x . The tangent bundle is the disjoint union of the tangent spaces, i.e. $\mathcal{T}M = \cup_{x \in M} \{x\} \times \mathcal{T}_x M$. Finally, *vector field* is a section of the tangent bundle, i.e. $\tilde{V} : M \rightarrow \mathcal{T}M$ such that $\tilde{V}(x) = (x, V(x))$, $V(x) \in \mathcal{T}_x M$. From now on, with a slight abuse of notation, we will identify \tilde{V} with V . Also, given $f \in \mathcal{C}^1(M_1, M_2)$, since the image of a \mathcal{C}^1 curve is a \mathcal{C}^1 curve, we have naturally defined a map $f_* : \mathcal{T}M_1 \rightarrow \mathcal{T}M_2$.

Problem 1.9 If $f \in \mathcal{C}^1(\mathbb{R}^d, \mathbb{R}^n)$ discuss the relation between f_* and the derivative Df .

We have finally the language to define O.D.E. on manifolds, in fact the Cauchy problem is exactly given again by (1.1.1), only now V is a, possibly time dependent, \mathcal{C}^1 vector field.

¹⁶If (U, ϕ) is a chart containing x , and γ_1, γ_2 two curves, think of the curves $\gamma_\lambda(t) = \gamma_1(\lambda t)$ and $\phi^{-1}(\phi(\gamma_1(t)) + \phi(\gamma_2(t)) - \phi(x))$.

Problem 1.10 Suppose that x_0 belongs to some chart (U, ϕ) , show that the solution of

$$\begin{aligned}\dot{x} &= V(x, t) \\ x(0) &= x_0\end{aligned}$$

for a sufficiently small time can be obtained by the solution of an appropriate O.D.E. in $\phi(U)$.

Problem 1.11 Given a finite atlas $\{(U_\alpha, \phi_\alpha)\}$, show that there exists a smooth partition of unity subordinated to the atlas, that is a collection $\{\phi_\alpha\} \in C^\infty(M, \mathbb{R})$ such that $\sum_\alpha \phi_\alpha = 1$ and $\text{supp } \phi_\alpha \subset U_\alpha$.

Problem 1.12 Given a smooth vector field V consider

$$\begin{aligned}\dot{x} &= V(x) \\ x(0) &= x_0\end{aligned}\tag{1.1.8}$$

with $x_0 \in U_\alpha$ for some element of an atlas $\{(U_\alpha, \phi_\alpha)\}$. Let $z_\alpha(t)$ be the solution of

$$\begin{aligned}\dot{z}_\alpha &= (\phi_\alpha)_* V(z_\alpha) \\ z_\alpha(0) &= \phi_\alpha(x_0)\end{aligned}$$

and suppose that $\phi_\alpha^{-1}(z_\alpha(1)) \in U_\beta$. Consider then the solution of

$$\begin{aligned}\dot{z}_\beta &= (\phi_\beta)_* V(z_\beta) \\ z_\beta(1) &= \phi_\beta(\phi_\alpha^{-1}(z_\alpha(1))).\end{aligned}$$

Show that there exists $t_1 > 1$ such that

$$\begin{aligned}x(t) &= \phi_\alpha^{-1}(z_\alpha(t)) \quad \text{for } t \in [0, 1] \\ x(t) &= \phi_\beta^{-1}(z_\beta(t)) \quad \text{for } t \in (1, t_1)\end{aligned}$$

is a solution of (1.1.8) in the time interval $[0, t_1)$.

Remark 1.1.15 We have seen that the theory of ODE on manifolds can be reduced locally to the case of \mathbb{R}^d . Yet, the reader should be aware that the global properties of the solutions can be very different. We will comment at length on this issue later on.

1.2 Linear ODE and Floquet theory

Let us briefly discuss the simplest possible differential equation: the affine ones. We restrict ourselves to the case $\mathcal{B} = \mathbb{R}^d$ for some $d \in \mathbb{N}$ since we will use some spectral theory which is substantially more complex in the general case.

1.2.1 Linear equations

Consider

$$\begin{aligned}\dot{x} &= Ax \\ x(0) &= x_0.\end{aligned}\tag{1.2.9}$$

Problem 1.13 *Show, by induction, that for each $n \in \mathbb{N}$ the solution of (1.2.9) satisfies*

$$x(t) = \sum_{k=0}^n \frac{1}{k!} A^k t^k x_0 + \int_0^t dt_1 \int_0^{t_1} dt_2 \cdots \int_0^{t_{n-1}} dt_n A^{n+1} x(t_n).$$

Taking the limit for $n \rightarrow \infty$ in the above expression one readily obtains $x(t) = \sum_{n=0}^{\infty} \frac{1}{n!} A^n t^n x_0$. That this is a solution can be verified directly inserting this formula in (1.2.9) (and noticing that the series and the series obtained by deviating term by term are uniformly convergent). By the standard analytic functional calculus for matrices (and operators, see Appendix C) we can thus write $x(t) = e^{At} x_0$. The above discussion provides a general solution for all equations of the type 1.2.9.

In reality life it is not that simple: if one has a concrete matrix A and wants to compute e^{At} , this may be quite unpleasant. A general strategy, although not necessarily the simplest one, is to perform a linear change of variables $x = Uz$. Then $\dot{z} = U^{-1}AUz$, and U is chosen so that $\Lambda = U^{-1}AU$ is in Jordan normal form. Then

$$x(t) = Uz(t) = Ue^{\Lambda t} z_0 = Ue^{\Lambda t} U^{-1} x_0.$$

It suffices then to know how to take exponentials of Jordan blocks, and this can be computed by using the defining series.

Problem 1.14 Compute $e^{\Lambda t}$ for

$$\Lambda = \begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix} ; \quad \Lambda = \begin{pmatrix} a & 1 \\ 0 & a \end{pmatrix} ; \quad \Lambda = \begin{pmatrix} a & 1 & 0 \\ 0 & a & 1 \\ 0 & 0 & a \end{pmatrix}.$$

Another, equivalent, point of view is to look for solutions of the type $x(t) = e^{at}v$, substituting in the first of (1.2.9) one obtains $av = Av$. Thus, as we know already, each eigenvalue of A provides a solution of (1.2.9) (ignoring the initial condition). If there exists real eigenvectors $\{v_i\}_{i=1}^d$ which span all \mathbb{R}^d then one can write the general solution, depending on d parameters α_i , as $x(t) = \sum_{i=1}^d \alpha_i v_i e^{a_i t}$, where a_i is the eigenvalue associated to the eigenvector v_i . One can then satisfy the initial condition by solving $x_0 = \sum_{i=1}^d \alpha_i v_i$. The same can be done if the eigenvectors are complex, by working in \mathbb{C}^d instead then \mathbb{R}^d . If Jordan blocks are present one can look for solutions of the form $x(t) = \sum_{k=0}^p \frac{1}{(p-k)!} t^k e^{at} v_k$, compare this formula with your solution of Problem 1.14.

Remark 1.2.1 Note that if the matrix A does not have eigenvalues with zero real part, then (by spectral decomposition) one can write $\mathbb{R}^d = V_- \oplus V_+$, where $AV_{\pm} = V_{\pm}$ and A restricted to V_- has eigenvalues with negative real part while on V_+ has eigenvalues with positive real part. Hence if $x_0 \in V_-$ it will hold $\lim_{n \rightarrow \infty} x(t) = 0$, and if $x_0 \in V_+$ it will hold $\lim_{n \rightarrow \infty} \|x(t)\| = \infty$. Accordingly if $x_0 \notin V_-$ we can write it as $x_0 = x_- + x_+$, where $x_{\pm} \in V_{\pm}$. Hence $\lim_{n \rightarrow \infty} \|x(t)\| = \infty$ and the trajectory will escape to infinity while getting exponentially close to the subspace V_+ . This is our first long time result.

A slightly more complex situation is given by

$$\begin{aligned} \dot{x} &= Ax + b(t) \\ x(0) &= x_0, \end{aligned} \tag{1.2.10}$$

where $b \in C^0(\mathbb{R}, \mathbb{R}^d)$. The solution of (1.2.10) is given by¹⁷

$$x(t) = e^{At}x_0 + \int_0^t e^{A(t-s)}b(s)ds. \tag{1.2.11}$$

¹⁷To obtain it just look for a solution of the form $x(t) = e^{At}z(t)$ and deduce the differential equation for z .

1.2.2 Floquet theory

Let us consider the simplest case of a linear time dependent equation: the case when there exists T such that $A(t+T) = A(t)$. More precisely, let $\Phi(x_0, s, t)$ be the solution of the Cauchy problem¹⁸

$$\begin{aligned}\dot{x}(t) &= A(t)x(t) \\ x(s) &= x_0.\end{aligned}\tag{1.2.12}$$

Problem 1.15 *Verify the following facts for each $x_0, y_0 \in \mathcal{B}$ and for each $a, b, t, s, \tau \in \mathbb{R}$*

- $\Phi(ax_0 + by_0, s, t) = a\Phi(x_0, s, t) + b\Phi(y_0, s, t)$,
- $\Phi(x_0, s, t) = \Phi(\Phi(x_0, s, \tau), \tau, t)$,
- $\Phi(x_0, s + T, t + T) = \Phi(x_0, s, t)$.

By the first property of Problem 1.15 there exists $K \in \mathcal{C}^1(\mathbb{R}^2, L(\mathcal{B}, \mathcal{B}))$ such that $\Phi(x_0, s, t) = K(s, t)x_0$, the second property implies that $K(\tau, t)K(s, \tau) = K(s, t)$, the third that $K(s + T, t + T) = K(s, t)$. The next step is the first occurrence in this book of a very simply but very powerful idea to analyze dynamical systems: a Poincaré section. Essentially the idea consist in looking at the system only at specially selected moments in time. In this case it is convenient to look at $t \in \{nT\}_{n \in \mathbb{Z}}$. That is, we want to investigate $\Phi(x_0, 0, nT) =: F(x_0, n)$.

Lemma 1.2.2 *The couple (\mathbb{R}^d, F) is a discrete Dynamical System.*

PROOF. We have to show that F is an action of \mathbb{Z} on \mathbb{R}^d . Let $f(x_0) := F(x_0, 1)$.

$$\begin{aligned}F(x_0, n) &= \Phi(x_0, 0, nT) = \Phi(\Phi(x_0, 0, (n-1)T), (n-1)T, nT) \\ &= \Phi(\Phi(x_0, 0, (n-1)T), 0, T) = f(\Phi(x_0, 0, (n-1)T)) = f^n(x_0).\end{aligned}$$

In addition, note that the uniqueness of the solutions of the ODE implies that if $f(x_0) = 0$, then $x_0 = 0$. Now, by construction, $f(x_0) = K(0, T)x_0$, thus $K(0, T)$ is an invertible matrix. Hence $F(x_0, -n) = f^{-n}(x_0)$ for all $n \in \mathbb{N}$. \square

¹⁸The solution is well defined for all times by Lemma 1.1.9.

By using the functional calculus (see Problem C.19) one can define $B := T^{-1} \ln K(0, T)$, so $e^{BT} = K(0, T)$. Let us now consider $P(t) := K(0, t)e^{-Bt}$.

$$\begin{aligned} P(t+T) &= K(0, t+T)e^{-B(t+T)} = K(T, t+T)K(0, T)K(0, T)^{-1}e^{-Bt} \\ &= K(0, t)e^{-Bt} = P(t). \end{aligned}$$

We have just proven the following result.

Theorem 1.2.3 (Floquet theorem) *The solutions of the equation (1.2.12) can be written as $x(t) = P(t)e^{Bt}K(s, 0)x_0$ where $P(t+T) = P(t)$ is periodic.*

Note that the matrix B can be complex valued. This can be avoided at a little extra cost.

Problem 1.16 *Prove that the solutions of the equation (1.2.12) can be written as $x(t) = P(t)e^{Bt}x_0$ where B is real and $P(t+2T) = P(t)$ is periodic of period $2T$.*

Note that Theorem 1.2.3 implies that the long time behavior is completely contained in the eigenvalues of the matrix B often called *floquet exponents*.

Problem 1.17 *Find the solutions of*

$$\dot{x} = a(t)Ax$$

where $a \in C^0(\mathbb{R}, \mathbb{R})$ is periodic of period T and A is a fixed matrix.

Problem 1.18 *Given a fixed matrix A and a function at matrix values $B(t)$ of period T , consider the equation $\dot{x} = (A + \varepsilon B(t))x$, $\varepsilon \in \mathbb{R}$. Show that, for ε small enough, calling ν_i the Floquet exponents and setting $\lambda_i = e^{\nu_i}$ (often called Floquet multiplier), the λ_i are ε -close to the eigenvalues of A .*

1.3 Qualitative study of ODE

The previous discussion has shed some light on the behavior of linear ODE, unfortunately the interesting ODE are typically non linear.

Although some nonlinear ODE can be solved explicitly (see any ODE book for examples) typically this is not possible, hence the need of a qualitative theory. As for the qualitative study of functions this can be done quite naively in one dimension, while higher dimensions requires some non trivial theory. Let us see such a naive qualitative theory for ODE via few examples.

1.3.1 The one dimensional case

This situation is very similar to the study of the graph of a function of one variable. Indeed to draw the graph one studies the first derivative and here the first derivative is specified by the equation. Let us consider a couple of simple examples. Consider

$$\begin{aligned}\dot{x} &= e^{-x^2} + x - 2 = V(x) \\ x_0 &= 0.\end{aligned}$$

One cannot integrate the function $V(x)^{-1}$ (which would yield an explicit solution of the ODE), yet from the equation follows that there exists a close to 2 such that \dot{x} is negative if $x \leq a$ and positive otherwise. This implies that the solution starts to be decreasing and keeps decreasing forever.

Next, consider

$$\begin{aligned}\dot{x} &= 1 - 2tx \\ x_0 &= a.\end{aligned}$$

Such an equation cannot be solved by separation of variables, yet the above arguments still apply. In particular, for $t \geq 0$, we have $\dot{x}(t) < 0$ iff $x(t) > \frac{1}{2t}$. On the other hand if $x(t) > \frac{1}{2t}$ it will be so forever. In fact, consider $g(t) = x(t) - \frac{1}{2t}$, then $g'(t) = \dot{x}(t) + \frac{1}{2t^2}$. So if $g(t) = 0$, then $g'(t) > 0$ hence for $t' < t$ one has $g(t') < 0$. Thus the solution will increase until it will intersect the curve $\frac{1}{2t}$ and then it will start decreasing but always staying above such a curve.

maybe expand and put picture?

1.3.2 Autonomous equations in two dimensions

In this case the basic idea is to consider one component as a function of the other and in this way reduce to the previous case. Let us see

some examples.

Van Der Pol equation

Consider the equation

$$\begin{aligned}\dot{x} &= y - f(x) \\ \dot{y} &= -x\end{aligned}$$

with $f(x) = x^3 - x$. The idea is to forget the time dependence and concentrate only on the shape of the trajectories. Thus we can represent trajectories on the xy plane. Indeed, apart from the point $(0, 0)$, either \dot{x} or \dot{y} are different from zero. In the first case one can locally invert $x(t)$ and write $y(x) = y(t(x))$. When this is possible one obtains

$$\frac{dy}{dx} = \frac{x}{f(x) - y},$$

which can be studied as in the previous examples.

Volterra equation

$$\begin{aligned}\dot{x} &= ax - Ax^2 - \lambda xy \\ \dot{y} &= -dy + \lambda xy.\end{aligned}$$

Second order in one dimension

Consider the equation

$$\begin{aligned}\ddot{x} &= -\gamma\dot{x} + \frac{x^2}{1+x^4} \\ x(0) &= 0; \quad \dot{x}(0) = v.\end{aligned}$$

Setting $(z, w) = (x, \dot{x})$, we can write it as

$$\begin{aligned}\dot{z} &= w \\ \dot{w} &= -\gamma w + \frac{z^2}{1+z^4}\end{aligned}$$

which is the type discussed above.

Clearly if we consider still higher dimensional case the above naive approach cannot help us very much, hence the need of a more sophisticated theory.

Problems

- 1.19.** Given two Banach spaces $\mathcal{B}_1, \mathcal{B}_2$ and a function $f : \mathcal{B}_1 \rightarrow \mathcal{B}_2$ we can define the partial derivative at $x \in \mathcal{B}_1$ in the direction $v \in \mathcal{B}_1$ (Gâteaux derivative) by

$$\partial_v f(x) = \lim_{h \rightarrow 0} h^{-1} [f(x + hv) - f(x)],$$

if the limit exists. On the other hand we say that f is Fréchet differentiable at x if there exists $A \in L(\mathcal{B}_1, \mathcal{B}_2)$ (the space of the continuous linear operators from \mathcal{B}_1 to \mathcal{B}_2) such that

$$\lim_{h \rightarrow 0} \frac{\|f(x + h) - f(x) - Ah\|}{\|h\|} = 0,$$

and A is called the Fréchet differential at f at x (often written $Df(x)$). Show that if f is Fréchet differentiable at zero, then it is continuous and Gâteaux differentiable.

- 1.20.** Let $f \in \mathcal{C}^0(\mathcal{B}_0, \mathcal{B}_1)$ and $g \in \mathcal{C}^0(\mathcal{B}_1, \mathcal{B}_2)$ such that f is Fréchet differentiable at $x \in \mathcal{B}_0$ and g is Fréchet differentiable at $f(x) \in \mathcal{B}_1$. Show that $g \circ f \in \mathcal{C}^0(\mathcal{B}_0, \mathcal{B}_2)$ is Fréchet differentiable at x and that $D(g \circ f)(x) = Dg(f(x)) \cdot Df(x) \in L(\mathcal{B}_0, \mathcal{B}_2)$. Of course, this is nothing else than a glorified version of the *chain rule*.
- 1.21.** Given a compact interval $I \subset \mathbb{R}$, a Banach space \mathcal{B} , and a continuous function $f \in \mathcal{C}^0(I, \mathcal{B})$, shows that one can define the equivalent of the Riemannian integral.
- 1.22.** Prove the fundamental theorem of calculus in this setting. That is, for $f \in \mathcal{C}^1(\mathcal{B}_1, \mathcal{B}_2)$ let $Df(x) \in L(\mathcal{B}_1, \mathcal{B}_2)$ be the Fréchet differential at $x \in \mathcal{B}_1$, then for each $x, y \in \mathcal{B}_1$

$$f(y) = f(x) + \int_0^1 Df(x + t(y - x)) \cdot (y - x) dt.$$

- 1.23.** A more general strategy to define the integral (simpler, but weaker, than the Bochner one) is as follows. Consider an interval $I \subset \mathbb{R}$, a Banach space \mathcal{B} , and a bounded function $f : I \rightarrow \mathcal{B}$, such

that there exist a sequence of simple functions¹⁹ $\{s_n\}$ with the property

$$\lim_{n \rightarrow \infty} \int_I \|f(t) - s_n(t)\| dt = 0.$$

Shows that there exists $A \in \mathcal{B}$ such that for each $\ell \in \mathcal{B}'$ holds²⁰

$$\ell(A) = \int_I \ell(f(t)) dt.$$

- 1.24.** Study the solutions of the following equations for all possible initial conditions and $p \in \mathbb{N}$

$$\dot{x} = |x|^p$$

$$\dot{x} = x(\ln |x|)^p$$

- 1.25.** Let $K \in \mathcal{C}^1(\mathbb{R} \times [0, 1])$. Show that the equation

$$\begin{aligned} \partial_t u(t, s) &= \int_0^1 K(t + s, \tau) u(t, \tau)^2 d\tau \\ u(0, s) &= s^2. \end{aligned}$$

has a unique continuous solution for t small enough.

- 1.26.** Under the same hypotheses of Problem 1.18 show that if $\int_0^T B(s) ds = 0$ and the eigenvalues of A have all multiplicity one, then the Floquet multiplier differ from the eigenvalues of e^{AT} only of order ε^2 .

- 1.27.** Study the equation

$$(1 + x)y\dot{y} + (x + y^2) = 0.$$

- 1.28.** Study the equation (Bernoulli)

$$\dot{y} + p(x)y = q(x)y^n.$$

- 1.29.** Study the equation

$$\ddot{x} = -\gamma\dot{x} - x^3.$$

¹⁹That is s_n is a piecewise constant function.

²⁰Remember that \mathcal{B}' is the vector space of all the linear *bounded* function from \mathcal{B} to \mathbb{R} (or \mathbb{C} if the Banach space is over the field of complex numbers). When equipped with the norm $\|\ell\| := \sup_{\|v\| \leq 1} |\ell(v)|$ it is itself a Banach Space. See [RS80] for details.

Hints to solving the Problems

In this section, and in the parallel sections in later chapters, I give hints for the solution of some of the Problems.

It is a very good idea to try very hard to solve the problems *before* looking at the hints: it is difficult to appreciate the solution if one has no feeling for the problem. The only way I know to get a feeling for a problem and its difficulty is to *seriously* try to solve it.

Also, keep in mind that I suggest one way to proceed, often other ways are possible and, possibly, even better.

1.1 Start by showing that $x \in \mathcal{C}^1(I, \mathcal{B})$.

1.13 For $n = 0$ it is just (1.1.2). To verify it for any n it suffices to show that

$$\int_0^t dt_1 \int_0^{t_1} dt_2 \cdots \int_0^{t_{n-1}} dt_n 1 = \frac{t^n}{(n+1)!}.$$

This follows since the domain of integration is $D = \{x \in [0, t]^{n+1} : t_{n+1} \leq t_n \leq \cdots \leq t\}$. On the other hand, for each permutation σ of the set $\{1, \dots, n+1\}$ the sets $D_\sigma = \{x \in [0, t]^{n+1} : t_{\sigma_{n+1}} \leq t_{\sigma_n} \leq \cdots \leq t\}$ have the same measure, all the D_σ are disjoint and the union of all of them gives $[0, t]^{n+1}$.

1.16 First notice that if a matrix has no eigenvalues on the negative axis then the contour γ in C.3.2 can be taken symmetric around the real axis and, by using C.3.2 with the standard definition of \ln with a cut on the negative real axis, this defines $\ln K(0, T)$ with real entries (since the formula for his complex conjugate is the same). In general use the spectral decomposition to write $K(0, T) = C + D$ where $\sigma(C) \cap \mathbb{R}_- = \emptyset$ and $\sigma(D) \subset \mathbb{R}_-$. Then $\sigma(D^2) \subset \mathbb{R}_+$, hence $B = \frac{1}{T} \ln C + \frac{1}{2T} \ln D^2$ is real and $e^{2BT} = C^2 + D^2 = K(0, T)^2$. The rest of the argument is as before.

1.18 Show that the solution satisfies

$$x(t) = e^{At} x_0 + \varepsilon \int_0^t e^{A(t-s)} B(s) x(s) ds.$$

and apply the perturbation theory in Appendix C.

1.21 Let $I = [a, b]$. Since the function is continuous, it is uniformly continuous, hence for $\varepsilon > 0$ there exists $\delta > 0$ such that, for each partition $\xi = \{[x_0, x_1], \dots, [x_{n-1}, x_n]\}$, $x_0 = a, x_n = b, x_{n+1} - x_n \leq \delta$, holds $\sup_{z, y \in [x_{n+1}, x_n]} \|f(z) - f(y)\| \leq \varepsilon$. Accordingly, for each choice of $z_n, y_n \in [x_{n+1}, x_n]$ we have

$$\left\| \sum_{k=0}^{n-1} f(z_k)(x_{k+1} - x_k) - \sum_{k=0}^{n-1} f(y_k)(x_{k+1} - x_k) \right\| \leq \varepsilon.$$

By similar arguments one can compare the sum defined on one partition with the sum defined on a finer partition. Finally sum on different partitions can be compared with the sum on the coarser partition finer of both. This shows that all sufficiently fine partitions yield the same approximate value, hence one can consider the partitions $\xi_n = \{[a + i\frac{b-a}{n}, a + (i+1)\frac{b-a}{n}]\}_{i=0}^{n-1}$ and define

$$\int_I f(t)dt := \lim_{n \rightarrow \infty} \sum_{i=0}^{n-1} f(a + i\frac{b-a}{n}) \frac{b-a}{n}.$$

By the above discussion this is equivalent to the same limit taken along any other partition the diameter of which elements tend uniformly to zero.

1.23 Since f is bounded, i.e. exists $M > 0$ such that $\sup_{t \in I} \|f(t)\| \leq M$, for each $\ell \in \mathcal{B}'$ holds $|\int_I \ell(f(t))dt| \leq |I| M \|\ell\|$. This means that the integral defines an element \tilde{A} of the double dual $(\mathcal{B}')'$, which normally is larger than \mathcal{B} . On the other hand for $s_n = \sum_i \alpha_i \mathbb{1}_{I_i}$, where I_i are the interval of constancy of s_n , we can define $A_n = \sum_i \alpha_i |I_i|$. Clearly $\ell(A_n) = \int_I \ell(s_n(t))dt$. By hypothesis A_n is a Cauchy sequence, hence will have a limit $A \in \mathcal{B}$, since for each $\ell \in \mathcal{B}'$ holds

$$\ell(A) = \lim_{n \rightarrow \infty} \ell(A_n) = \lim_{n \rightarrow \infty} \int_I \ell(s_n(t))dt = \int_I \ell(f(t))dt = \ell(\tilde{A})$$

it follows $\tilde{A} = A \in \mathcal{B}$.

1.25 Consider the Banach space $\mathcal{B} = \mathcal{C}^0([0, 1], \mathbb{R})$. Then $u(t, \cdot) \in \mathcal{B}$ and one can apply Theorem 1.1.1.

1.26 By Problem 1.18 we know that the solution at time T is given by the matrix $D(\varepsilon) := e^{AT} \left[\mathbf{1} + \varepsilon \int_0^T e^{-As} B(s) e^{As} ds \right]$. By the results in Appendix C it follows that, for ε small enough, the eigenvalues of $D(\varepsilon)$ are still simple and analytic on ε . Thus, let $\lambda(\varepsilon)$ one of such eigenvalues and $\Pi(\varepsilon)$ the associated eigenprojector. We have $D(\varepsilon)\Pi(\varepsilon) = \lambda(\varepsilon)\Pi(\varepsilon)$. Differentiating yields $\dot{D}(\varepsilon)\Pi(\varepsilon) + D(\varepsilon)\dot{\Pi}(\varepsilon) = \dot{\lambda}(\varepsilon)\Pi(\varepsilon) + \lambda(\varepsilon)\dot{\Pi}(\varepsilon)$. Multiplying on the right by $\Pi(\varepsilon)$, since $\Pi(\varepsilon)D(\varepsilon) = D(\varepsilon)\Pi(\varepsilon)$, we have

$$\Pi(\varepsilon)\dot{D}(\varepsilon)\Pi(\varepsilon) = \dot{\lambda}(\varepsilon)\Pi(\varepsilon).$$

Since $\Pi(\varepsilon)v = \langle a(\varepsilon), v \rangle b(\varepsilon)$ for some vectors a, b analytic in ε , $\dot{\lambda}(\varepsilon) = \langle a(\varepsilon), \dot{D}(\varepsilon)b(\varepsilon) \rangle$. We can now apply such a general formula to our specific case:

$$\begin{aligned} \langle a(0), \dot{D}(0)b(0) \rangle &= \langle a(0), e^{AT} \int_0^T e^{-As} B(s) e^{As} b(0) ds \rangle \\ &= \langle a(0), e^{AT} \int_0^T e^{-As} B(s) e^{As} b(0) ds \rangle \\ &= \lambda(0) \int_0^T \langle a(0), B(s)b(0) \rangle ds = 0. \end{aligned}$$

Notes

I presented the results in a really condensed way and with no pretension to exhaust the theory of ODE. If one wants to have a better understanding of the field and some ideas of how an ODE can be solved in special cases better consult [HS74, Arn92, CL55].