

# Towards integrable perturbation of 2d CFT on de Sitter space

Christian D. Jäkel <sup>\*</sup>1 and Yoh Tanimoto<sup>†</sup>2

<sup>1</sup>Department of Applied Mathematics, University of São Paulo (USP),  
Rua de Matão 1010, CEP 05508-090 São Paulo, Brazil

<sup>2</sup>Dipartimento di Matematica, Università di Roma Tor Vergata,  
Via della Ricerca Scientifica 1, I-00133 Roma, Italy

## Abstract

We describe a procedure to deform the dynamics of a two-dimensional conformal net to possibly obtain a Haag-Kastler net on the de Sitter spacetime. The new dynamics is given by adding a primary field smeared on the time-zero circle to the Lorentz generators of the conformal net.

As an example, we take an extension of the chiral  $U(1)$ -current net by a charged field with conformal dimension  $d < \frac{1}{4}$ . We show that the perturbing operators are defined on a dense domain.

## 1 Introduction

The first interacting quantum field theories, the  $\mathcal{P}(\phi)_2$ -models, have been constructed by starting with the free field on the Minkowski space, defining an interaction term, perturbing the dynamics by it locally, finding the interacting vacuum and changing the Hilbert space [GJ72]. The  $\mathcal{P}(\phi)_2$ -models have been constructed also on the de Sitter space [FHK75], then recently formulated into the operator-algebraic framework [BJM23]. Perturbing the dynamics on the de Sitter space has the advantage that one may construct interacting models on the same Hilbert space, as one can avoid Haag's theorem [Wei11].

These procedures of perturbing the dynamics have been formulated more generally in [JM18]: One starts with a Haag-Kastler net on the de Sitter space (in the sense of [BB99]), chooses the new dynamics with an interacting vacuum. The above  $\mathcal{P}(\phi)_2$ -models fit in this programme. As the arguments do not depend particularly on the free fields, one may wish to find other examples. We propose such an example in this work, where the starting QFT is a two-dimensional conformal field theory and the perturbation is given by a primary field. Specifically, the conformal field theory is a two-dimensional extension of the chiral  $U(1)$ -current algebra, and we take the charge-carrying field as the interaction term. Such fields

---

<sup>\*</sup>jaekel@ime.usp.br

<sup>†</sup>hojt@mat.uniroma2.it

have been constructed recently as two-dimensional conformal Wightman fields [AGT]. Such a conformal field can be seen as a field on the de Sitter space through a conformal map.

The  $U(1)$ -current algebra is defined on the Hilbert space  $\mathcal{H}_0$ , and the two-dimensional extension contains two copies of it as the left and right chiral components. The chiral components have charged sectors  $\mathcal{H}_\alpha$  parametrized by  $\alpha \in \mathbb{R}$ . For a fixed  $\alpha \in \mathbb{R}$ , we take  $\bigoplus_{j \in \mathbb{Z}} \mathcal{H}_{j\alpha} \otimes \mathcal{H}_{j\alpha}$  as the Hilbert space<sup>1</sup>. There is a charged field  $Y_\alpha(z)$  that maps  $\mathcal{H}_\beta \otimes \mathcal{H}_\beta$  to  $\mathcal{H}_{\beta+\alpha} \otimes \mathcal{H}_{\beta+\alpha}$ . This is the basis of our perturbing field. We show that the symmetric field  $Y_\alpha(z) \otimes Y_\alpha(z^{-1}) + Y_\alpha(z)^* \otimes Y_\alpha(z^{-1})^*$  can be added to the Lorentz generators of the de Sitter space, and they still satisfy the Lorentz relations weakly.

From a physical point of view, perturbing the Hamiltonian of a CFT by a (relevant) field has been proposed to obtain massive integrable models in [Zam89]. Depending on the initial CFT and the perturbing field, various integrable models should be obtained. While our results are specific to the  $U(1)$ -current, the proof of (weak) Lorentz relations depends essentially on the fact that we take a primary field that is commutative at the time-zero circle. Therefore, the idea should generalize to many CFTs and primary fields.

This paper is organized as follows. In Section 2, we briefly recall the algebraic framework on the de Sitter spacetime, how a two-dimensional CFT can be considered on the de Sitter spacetime and the perturbation of the dynamics by a local field. In Section 3, a family of two-dimensional extensions of the  $U(1)$ -current net and their charged fields are reviewed. In Section 4, we make estimates of the charged fields restricted to the time-zero circle and show that the restriction defines operators if the charge  $\alpha$  satisfies  $|\alpha| < \frac{1}{\sqrt{2}}$ . Section 5 shows that the time-zero charged fields commute with each other. In Section 6, we show that the Lorentz generators perturbed by the charged field still satisfy the Lorentz relations weakly on a certain domain. In Section 7, we describe how this programme can be completed.

## 2 General strategy

### 2.1 Haag-Kastler nets on the de Sitter space

The two-dimensional de Sitter space  $dS^2$  is embedded in the ambient three-dimensional Minkowski space  $\mathbb{R}^{2+1}$  by the equation  $x_0^2 - x_1^2 - x_2^2 = r^2$ , where  $r > 0$ . The set  $dS^2$  is invariant under the (proper orthochronous) Lorentz group  $\mathcal{L}_+^\uparrow$  (the connected component of the stabilizer subgroup of the point  $(0, 0, 0)$  in the three-dimensional Poincaré group). On this space, the causal structure and the metric can be introduced by restricting those of the ambient Minkowski space. The region  $\{(x_0, x_1, x_2) \mid |x_0| < x_1\}$  is called the wedge in the  $x_1$ -direction. We denote its intersection with  $dS^2$  by  $W_1$ . Any image of  $W_1$  by a Lorentz transformation is called a **wedge** in  $dS^2$ . For any wedge  $W$  there is a one-parameter group  $\Lambda_W(t)$  of Lorentz boosts that fix  $W$ , which are referred to as the boosts associated with  $W$ .

The Haag-Kastler axioms, usually considered on the Minkowski space, can be also formulated on  $dS^2$  [BB99], where the spectrum condition is replaced by the geodesic KMS property as below. A **Haag-Kastler net on  $dS^2$**  is a triple  $(\mathcal{A}, U, \Omega)$ , where  $\mathcal{A}$  is a family of von Neumann algebras on a Hilbert space  $\mathcal{H}$  parametrized by open regions  $O \subset dS^2$ ,  $U$  is a

---

<sup>1</sup>From a general point of view, it is more natural to take  $\mathcal{H}_{j\alpha} \otimes \mathcal{H}_{-j\alpha}$  (cf. [LR95]). In this case, the resulting net is unitarily equivalent through the map  $J(z) \rightarrow -J(z)$ , which can be unitarily implemented.

unitary representation of  $\mathcal{L}_+^\uparrow$  on  $\mathcal{H}$  (continuous in the strong-operator topology) and  $\Omega$  is a vector in  $\mathcal{H}$ , such that

(HK1) **Isotony:**  $\mathcal{A}(O_1) \subset \mathcal{A}(O_2)$  for  $O_1 \subset O_2$ ;

(HK2) **Locality:** If  $O_1$  and  $O_2$  are spacelike separated, then  $\mathcal{A}(O_1) \subset \mathcal{A}(O_2)'$ ;

(HK3) **Lorentz covariance:**  $\mathcal{A}(gO) = \text{Ad } U(g)(\mathcal{A}(O))$  for  $g \in \mathcal{L}_+^\uparrow$ ;

(HK4) **Cyclicity:**  $\Omega$  is cyclic for each  $\mathcal{A}(O)$ ;

(HK5) **The geodesic KMS property:** For any wedge  $W$ , it holds that  $U(\Lambda_W(2\pi t)) = \Delta_W^{-it}$ , where  $\Delta_W^{it}$  is the modular group of the algebra  $\mathcal{A}(W)$  with respect to  $\Omega$ .

The geodesic KMS property is equivalently stated by saying that  $U(\Lambda_W(t))$  satisfies the KMS condition for  $\mathcal{A}(W)$  with temperature  $2\pi$  with respect to  $\Omega$  [BB99].

## 2.2 Two-dimensional conformal net on the de Sitter space

Our starting point is a two-dimensional conformal field theory. A two-dimensional conformal field theory on the Minkowski space is a theory that is (locally) covariant with respect not only to the Poincaré group but also to the universal covering  $\overline{\text{Möb}} \times \overline{\text{Möb}}$  of the Möbius group  $\text{Möb} \times \text{Möb}$  including special conformal transformations, and often further to  $\text{Diff}_+(S^1) \times \text{Diff}_+(S^1)$ . It is known that any two-dimensional conformal (Haag-Kastler) net, *a priori* defined on the Minkowski space  $\mathbb{R}^{1+1}$ , extends to the Einstein cylinder [KL04][MT19, Theorem A.5], and then restricted to a part of it that is conformally equivalent to the de Sitter space  $\text{dS}^2$  [GL03]. Therefore, a two-dimensional CFT can be considered as a QFT on  $\text{dS}^2$  in this natural sense. Let us briefly review how this is done.

In our framework, a conformal Haag-Kastler net can be described as follows. First, we consider  $\mathbb{R}^{1+1}$  as the product of two lightrays. Each lightray  $\mathbb{R}$  has  $S^1$  as the one-point compactification, and the group  $\text{Diff}_+(S^1)$  acts on it. Therefore,  $\overline{\text{Diff}_+(S^1)} \times \overline{\text{Diff}_+(S^1)}$  acts on  $\mathbb{R}^{1+1}$  locally in the sense of [BGL93]. Furthermore, by spacelike locality, this action factors through the subgroup  $\mathfrak{R} := \{R_{2n\pi} \times R_{-2n\pi} : n \in \mathbb{Z}\}$  where  $R_t \in \overline{\text{Diff}_+(S^1)}$  is the lift of the rotation by  $t$  [KL04, Proposition 2.1] (see also [MT19, Theorem A.5]). We denote this group by  $\mathcal{C}$ . The Minkowski space  $\mathbb{R}^{1+1}$  is conformally equivalent to the product  $I_{2\pi} \times I_{2\pi}$  of open intervals of length  $2\pi$ , and through the local action of  $\mathcal{C}$ , the Haag-Kastler net can be extended to  $\mathbb{R} \times \mathbb{R}$  quotiented by the action of  $\mathfrak{R}$ , where  $\mathbb{R}$  is the universal covering of  $S^1$ . This space is conformally equivalent to the Einstein cylinder  $\mathcal{E} = S^1 \times \mathbb{R}$  (the product structure is different from the previous one). We say that two regions  $O_1, O_2$  are spacelike separated if there is a diamond obtained by shifting the Minkowski space  $I_{2\pi} \times I_{2\pi}$ , which includes  $O_1, O_2$  such that  $O_1$  and  $O_2$  are spacelike separated there.

To be precise, the axioms for conformal nets on  $\mathcal{E}$  are the following: Let  $\mathcal{A}$  be a family of von Neumann algebras on  $\mathcal{H}$  parametrized by open regions in  $\mathcal{E}$ , let  $U$  be a unitary projective representation of the group  $\mathcal{C}$  on  $\mathcal{H}$  (note that the restriction of  $U$  to the subgroup  $\overline{\text{Möb}} \times \overline{\text{Möb}} / \mathfrak{R}$  can be actually made into a true representation of  $\overline{\text{Möb}} \times \overline{\text{Möb}}$  [Bar54, Theorem 7.1], and the generators of one-parameter subgroups in  $\overline{\text{Möb}} \times \overline{\text{Möb}}$  are uniquely defined) and let  $\Omega$  be a vector in  $\mathcal{H}$  such that

- (CN1) **Isotony:**  $\mathcal{A}(O_1) \subset \mathcal{A}(O_2)$  for  $O_1 \subset O_2$ ;
- (CN2) **Locality:** If  $O_1$  and  $O_2$  are spacelike separated, then  $\mathcal{A}(O_1) \subset \mathcal{A}(O_2)'$ ;
- (CN3) **Conformal covariance:**  $\mathcal{A}(\gamma O) = \text{Ad } U(\gamma)(\mathcal{A}(O))$  for  $\gamma \in \mathcal{C}$ , and if  $O$  is disjoint from  $\text{supp } \gamma$ , then  $\text{Ad } U(\gamma)(x) = x$  for  $x \in \mathcal{A}(O)$ ;
- (CN4) **Positive energy:** The generators of the chiral rotation subgroups  $R_t \times \iota, \iota \times R_t$ , where  $\iota \in \overline{\text{Diff}}_+(S^1)$  is the unit element, are positive;
- (CN5) **Vacuum:**  $\Omega$  is cyclic for each  $\mathcal{A}(O)$  and is a unique (up to scalar) vector such that  $U(\gamma)\Omega = \Omega$  for  $\gamma \in \overline{\text{Möb}} \times \overline{\text{Möb}}$  in the sense above.

In a conformal net, the Bisognano-Wichmann property holds automatically [BGL93].

On the Einstein cylinder  $\mathcal{E}$ , the vertical strip of length  $\pi$  is conformally equivalent to the de Sitter space [GL03] (for each radius  $r$  there are different de Sitter space, but we do not specify it, because the only point is that any of such de Sitter space is conformally equivalent to the same part of the cylinder), see Figure 1. The time-zero circle  $S^1$  ( $x_0 = 0$ ) in the de Sitter space is the (compactified) time-zero line ( $a_0 = 0$ ) on the cylinder, and space rotations act on it. Other Lorentz transformations are contained in the conformal group  $\mathcal{C}$ . Indeed, the spacelike rotations  $R_t \times R_{-t}$  and Lorentz boosts (that is the product of lightlike dilations with opposite sign) generate the three-dimensional Lie group  $\text{SO}(2, 1)$ , the  $(2+1)$ -dimensional Lorentz group. Therefore, by restricting a conformal net to the de Sitter space, it satisfies the axioms (HK1-5): the geodesic KMS property is satisfied because of the Bisognano-Wichmann property.




Figure 1: The Minkowski space  $M_0$  (cf. [AGT, Figure 1]) and the de Sitter space  $dS^2$  conformally embedded in  $\mathbb{R}^2$ . The cylinder is obtained by identifying the dotted lines. The dark grey region is a wedge  $W$  and the light grey region is a double cone.

In general, a two-dimensional conformal net  $\mathcal{A}$  contains **chiral components**, that are observables living on the lightrays and invariant under the action of  $\iota \times \overline{\text{Diff}}_+(S^1)$  ( $\overline{\text{Diff}}_+(S^1) \times \iota$ , respectively). They can be regarded as Haag-Kastler nets on the lightrays  $\mathbb{R}$ , extend to  $S^1$  by conformal covariance and are called **conformal nets on  $S^1$** . More precisely, a triple  $(\mathcal{A}_0, U_0, \Omega_0)$  where  $\mathcal{A}_0$  is a family of von Neumann algebras on a Hilbert space

$\mathcal{H}_0$  parametrized by open connected nonempty non dense intervals in  $S^1$ ,  $U_0$  is a unitary projective representation of  $\text{Diff}_+(S^1)$  and  $\Omega_0$  is a vector in  $\mathcal{H}_0$ , is called a **conformal net on  $S^1$**  if it satisfies

- (CNS1) **Isotony:**  $\mathcal{A}_0(I_1) \subset \mathcal{A}_0(I_2)$  for  $I_1 \subset I_2$ ;
- (CNS2) **Locality:** If  $I_1$  and  $I_2$  are disjoint, then  $\mathcal{A}_0(I_1) \subset \mathcal{A}_0(I_2)'$ ;
- (CNS3) **Conformal covariance:**  $\mathcal{A}_0(\gamma I) = \text{Ad } U_0(\gamma)(\mathcal{A}_0(I))$  for  $\gamma \in \text{Diff}_+(S^1)$ , and if  $I$  is disjoint from  $\text{supp } \gamma$ , then  $\text{Ad } U_0(\gamma)(x) = x$  for  $x \in \mathcal{A}_0(I)$ ;
- (CNS4) **Positive energy:** The generators of the rotation subgroup in  $\text{Diff}_+(S^1)$  is positive;
- (CNS5) **Vacuum:**  $\Omega_0$  is cyclic for each  $\mathcal{A}_0(O)$  and is a unique (up to a scalar) vector such that  $U_0(\gamma)\Omega_0 = \Omega_0$  for  $\gamma \in \text{Möb}$ ;

A two-dimensional conformal net  $\mathcal{A}$  contains both left and right chiral components, indeed, the operators  $U(\gamma_0 \times \iota), U(\iota \times \gamma_0)$  are such elements. In addition, left and right chiral components commute with each other [Reh00].

The positive-energy representation  $U_0$  is associated with a positive-energy representation of the Virasoro algebra  $\{L_m\}$  [Car04, Appendix]:

$$[L_m, L_n] = (m - n)L_{m+n} + \frac{c}{12}m(m^2 - 1)\delta_{m,-n},$$

for a certain value  $c > 0$  (by an abuse of notations, we use the symbols  $\{L_m\}$  both for abstract Lie algebra elements and for unbounded operators satisfying the above relations). The self-adjoint operators  $\mathfrak{t} = \frac{1}{2}L_0 - \frac{1}{4}(L_1 + L_{-1}), \mathfrak{d} = \frac{1}{2}(L_{-1} - L_1)$  are the generators of translations and dilations of  $\mathbb{R} \subset S^1$  (in the sense that  $\mathbb{R}$  is embedded in  $S^1$  by the stereographic projection), respectively [Wei05, Appendix A].

The two-dimensional conformal group  $\mathcal{C}$  has the tensor product  $\{L_m \otimes \mathbb{1}, \mathbb{1} \otimes L_n\}$  as the Lie algebra. The Lorentz boosts are generated by  $\mathfrak{d} \otimes \mathbb{1} - \mathbb{1} \otimes \mathfrak{d}$ , while the spacelike rotations are generated by  $L_0 \otimes \mathbb{1} - \mathbb{1} \otimes L_0$ .

For  $z \in S^1$ , one can consider the operator-valued distribution  $T(z) = \sum_n L_n z^{-n}$ , called the Virasoro field (the convention of the exponent is the one such that  $T(z)^* = T(z)$  [CTW19], different from  $L(z) = \sum_n L_n z^{-n-2}$  in vertex algebras [Kac98]). The two-dimensional stress-energy tensor  $T^{\mu\nu}$  has four components, and the fields  $T(z) \otimes \mathbb{1} + \mathbb{1} \otimes T(z^{-1})$  and  $T(z) \otimes \mathbb{1} - \mathbb{1} \otimes T(z^{-1})$  correspond to the components of the stress-energy tensor  $T^{01}$  and  $T^{00}$  [Reh].

## 2.3 Perturbation by a local field

As we saw above, a two-dimensional conformal net can be considered as a Haag-Kastler net on  $dS^2$  and hence as a starting point for a new construction in the sense of [JM18]. We take a starting Haag-Kastler net  $\mathcal{A}$  on  $dS^2$ , a unitary representation  $U$  of the  $(2+1)$ -dimensional Lorentz group and a vacuum vector  $\Omega$ . We call “time-zero wedges” wedge regions whose end points reside on  $S^1$ .

The general strategy of [JM18, Theorem 4.1] goes as follows:

- Assume there is a rotation-invariant vector  $\Omega$  cyclic for  $\mathcal{A}(W_1)$  in the natural positive cone  $\mathcal{P}(\mathcal{A}(W_1), \Omega_0)$  associated to the pair  $\mathcal{A}(W_1)$  and the free vacuum vector  $\Omega_0$ . The rotation-invariance of  $\Omega$  implies that  $\Omega \in \mathcal{P}(\mathcal{A}(W), \Omega_0)$  for all time-zero wedges  $W$ .
- Assume there exists a new representation  $\tilde{U}$  of the Lorentz group such that
  - its restriction to the rotation subgroup coincides with that of  $U$ ;
  - its (interacting) Lorentz boost associated to the wedge  $W_1$  is implemented by the modular group for the pair  $\mathcal{A}(W_1)$  and (the interacting vacuum vector)  $\Omega$ ;
  - it satisfies the finite speed of light condition [JM18, Definition 3.3], which roughly says that the action of  $\tilde{U}$  on the time-zero algebras  $\mathcal{A}(O_I)$ , with  $O_I$  a double cone given by the intersection of time-zero wedges, preserves locality.

The last two assumptions should be satisfied automatically if the new representation  $\tilde{U}$  is generated from a local field, as below.

- The Lorentz covariance of the new net is given by  $\tilde{U}$ . For any wedge region  $W$ ,  $\tilde{\mathcal{A}}(W) = \text{Ad } \tilde{U}(g)(\mathcal{A}(W_1))$ , where  $g$  is such that  $gW_1 = W$  (this is well-defined by finite speed of propagation, in particular, the time-zero wedge algebra is preserved by the new boosts with respect to the same wedge). Any double cone is written as  $O = \bigcap_{W \supset O} W$ , and accordingly we define  $\tilde{\mathcal{A}}(O) = \bigcap_{W \supset O} \tilde{\mathcal{A}}(W)$ . This satisfies locality again by finite speed of propagation.

Then  $(\tilde{\mathcal{A}}, \tilde{U}, \tilde{\Omega})$  is a new Haag-Kastler net on  $\text{dS}^2$ . This construction avoids Haag's theorem [Wei11], because the spacetime is compact and there is no dilation covariance that pushes a double cone to infinity.

Assume that the net  $\mathcal{A}$  is generated by a conformal Wightman field  $\psi$  and it has a well-defined restriction to the  $a_0 = 0$  circle, which we denote by  $\psi(0, \theta)$ . Let  $\mathbf{e}_n(\theta) = e^{in\theta}$ . The field  $\psi(0, \theta)$  smeared by a test function  $f$  on the circle is denoted by  $\psi(0, f)$ . We find interesting candidates for such  $\tilde{U}$  by adding the generators  $\psi(0, \mathbf{e}_1), \psi(0, \mathbf{e}_0), \psi(0, \mathbf{e}_{-1})$  to  $L_1 \otimes \mathbb{1} + \mathbb{1} \otimes L_{-1}, L_0 \otimes \mathbb{1} - \mathbb{1} \otimes L_0, L_{-1} \otimes \mathbb{1} + \mathbb{1} \otimes L_1$  of the Lorentz group  $\mathcal{L}_+^\uparrow$ .

Below we take concrete examples of two-dimensional CFTs and a candidate for  $\tilde{U}$  using the charged fields in it.

## 3 U(1)-current and two-dimensional extension

### 3.1 Chiral components

To make the programme concrete, we take an explicit example of two-dimensional CFT, whose chiral component is the U(1)-current net on  $S^1$  [BMT88]. It is generated by the current (the derivative of the massless free field)  $J(z) = \sum_n J_n z^{-n-1}$ , where  $z \in S^1$ , and its Fourier coefficients  $J_n$  satisfy the commutation relations

$$[J_m, J_n] = n\delta_{m, -n}.$$

There is a representation of this algebra with a unique **vacuum vector**  $\Omega_0$  such that  $J_n \Omega_0 = 0$  for  $n \geq 0$ , and equipped with an inner product  $\langle \cdot, \cdot \rangle$  with respect to which it holds that  $J_n^* = J_{-n}$ . The Hilbert space  $\mathcal{H}_0$  is spanned by the vectors of the form

$$J_{-n_1} \cdots J_{-n_k} \Omega_0,$$

where  $n_1 \geq n_2 \geq \cdots \geq n_k$ . We denote by  $\mathcal{H}_0^{\text{fin}}$  the linear span of these vectors.

This current can be smeared by a smooth function  $f$  and gives an unbounded operator  $J(f) = \sum_n f_n J_n$ , where  $f_n = \frac{1}{2\pi} \int e^{-in\theta} f(\theta) d\theta$  are the Fourier components. The exponential  $W(f) = e^{iJ(f)}$  is called a Weyl operator. One can construct a conformal net on  $S^1$  by  $\mathcal{A}_0(I) = \{e^{iJ(f)} : \text{supp } f \subset I\}''$ . A representation of the Virasoro algebra  $\{L_n\}$  is given by the Sugawara formula  $L_n = \frac{1}{2} \sum_k : J_{n-k} J_k :$ , and it integrates to a projective unitary representation  $U_0$  of  $\text{Diff}_+(S^1)$ . With this  $U_0$ ,  $(\mathcal{A}_0, U_0, \Omega_0)$  is called the **U(1)-current net**.

A **representation** of a conformal net  $\mathcal{A}_0$  on  $S^1$  is a family of isomorphisms  $\{\rho_I\}$  of local algebras  $\{\mathcal{A}_0(I)\}$  to von Neuman algebras on a certain Hilbert space  $\mathcal{H}_\rho$  that satisfy the compatibility condition

$$\rho_{I_1}(x) = \rho_{I_2}(x) \quad \text{for } x \in \mathcal{A}(I_1), \quad I_1 \subset I_2.$$

For each  $\alpha \in \mathbb{R}$ , the U(1)-current net admits a representation  $\rho_\alpha$ . This representation  $\rho_\alpha$  can be realized on the same Hilbert space  $\mathcal{H}_0$  as the vacuum representation, but to distinguish them we denote it by  $\mathcal{H}_\alpha$  and the lowest weight vector by  $\Omega_\alpha$ . The assignment  $W(f) \mapsto \rho_\alpha(W(f)) = e^{i\alpha \int f(\theta) d\theta} W(f)$  gives the representation. This is an **automorphism** of each local algebra  $\mathcal{A}_0(I)$ . In terms of generators, it amounts to replace  $J_0$  (which acts as the 0 operator in the vacuum representation) by the scalar  $\alpha$ . We denote by  $\mathcal{H}_\alpha^{\text{fin}}$  the subspace spanned by vectors of the form  $J_{-n_1} \cdots J_{-n_k} \Omega_\alpha$ .

Two such automorphisms  $\rho_{\alpha_1}, \rho_{\alpha_2}$  can be composed, and yield a new automorphism (representation)  $\rho_{\alpha_1 + \alpha_2}$ . This composition law is called the fusion rule for the U(1)-current. For a fixed  $\alpha \in \mathbb{R}$ , the family  $\{\rho_{j\alpha}\}_{j \in \mathbb{Z}}$  on  $\{\mathcal{H}_{j\alpha}\}$  is closed under fusion, indeed  $j_1\alpha + j_2\alpha = (j_1 + j_2)\alpha$ .

On these representations spaces  $\mathcal{H}_\alpha$ , there are also unbounded operators  $\{J_{\alpha,n}, L_{\alpha,n}\}$  satisfying the same algebraic relations. In particular,  $J_{\alpha,n}$  restrict to the representation on  $\mathcal{H}_\alpha$ . Furthermore, the Sugawara formula holds:  $L_{\alpha,n} = \frac{1}{2} \sum_k : J_{\alpha,n-k} J_{\alpha,k} :$ .

## 3.2 Charged primary fields

Recall [AGT] that we use formal series  $\sum_{s \in \mathbb{R}} A_s z^s$  for a family of operators  $\{A_s\}_{s \in \mathbb{R}}$  (actually the formal series is just the parametrized family  $\{A_s\}$  itself, but certain operations on them are implicit).

Let us fix  $\alpha \in \mathbb{R}$  with  $|\alpha| \leq 1$ . We constructed a formal series  $Y_\alpha$  acting on the dense domain in the Hilbert space  $\hat{\mathcal{H}} = \bigoplus_{j \in \mathbb{Z}} \mathcal{H}_{j\alpha}$ . On this space, operators  $\hat{J}_n = \bigoplus_{j \in \mathbb{Z}} J_{j\alpha,n}$ ,  $\hat{L}_n = \bigoplus_{j \in \mathbb{Z}} L_{j\alpha,n}$  can be defined naturally and satisfy the same commutation relations. The latter generates a projective unitary representation  $\hat{U}_0$  of  $\overline{\text{Diff}_+(S^1)}$ . Let  $d = \frac{\alpha^2}{2}$ . We know from [AGT] (see [TZ12] and [TL97] for the original references) that  $Y_\alpha$  satisfies

$$[\hat{L}_m, Y_{\alpha,s}] = ((d-1)m - s) Y_{\alpha,m+s}, \quad (1)$$

$$\|Y_{\alpha,s}\| \leq 1. \quad (2)$$

By (2), we can smear the field by a test function supported in  $\mathbb{R}$  and obtain  $Y_\alpha(f)$ . By (1), it is conformally covariant with respect to  $\hat{U}$  with the conformal dimension  $d$ . If we construct the formal series  $Y_\alpha(z) = \sum_{s \in \mathbb{R}} Y_{\alpha,s} z^{-s-d}$ , it is not local with itself, but satisfies a braiding relation.

More explicitly, these are constructed as follows: Each coefficient  $Y_{\alpha,s}$  is a map  $\mathcal{H}_\beta \rightarrow \mathcal{H}_{\beta+\alpha}$  (on each  $\mathcal{H}_\beta$ , only  $Y_{\alpha,s}$  with  $s \in \mathbb{Z} + \alpha\beta - d$  are non-zero). Let  $c_\alpha$  be the unitary charge shift operator  $\mathcal{H}_\beta \rightarrow \mathcal{H}_{\beta+\alpha}$  defined by  $c_\alpha \hat{J}_{-n_1} \cdots \hat{J}_{-n_k} \Omega_\beta = \hat{J}_{-n_1} \cdots \hat{J}_{-n_k} \Omega_{\beta+\alpha}$ ,  $n_j > 0$ . We can regard  $c_\alpha$  as an operator on  $\hat{\mathcal{H}} = \bigoplus_{j \in \mathbb{Z}} \mathcal{H}_{j\alpha}$ . Following [TZ12], we define

$$E^\pm(\alpha, z) = \exp \left( \mp \sum_{n>0} \frac{\alpha \hat{J}_{\pm n}}{n} z^{\mp n} \right),$$

$$Y_\alpha(z) = c_\alpha E^-(\alpha, z) E^+(\alpha, z) z^{\alpha J_0}, \quad (3)$$

where  $z^{\alpha J_0}$  means  $z^{\alpha\beta}$  on  $\mathcal{H}_\beta$ .

From this construction, it is easy to see the commutation relation  $[\hat{J}_m, Y_\alpha(z)] = \alpha Y_\alpha(z) z^m$ , or equivalently,

$$[\hat{J}_m, Y_{\alpha,s}] = \alpha Y_{\alpha, m+s}. \quad (4)$$

This implies that  $Y_\alpha(z)$  is relatively local to  $J(w)$ .

### 3.3 Two-dimensional Wightman field

From two copies of a conformal nets  $\mathcal{A}_0$  on  $S^1$ , one can construct a two-dimensional conformal net by tensor product:  $\mathcal{A}_0(I_+ \times I_-) := \mathcal{A}_0(I_+) \otimes \mathcal{A}_0(I_-)$ . The unitary representation of  $\mathcal{C}$  is given by  $U_0(\gamma_+) \otimes U_0(\gamma_-)$  and the vacuum by  $\Omega_0 \otimes \Omega_0$ . For a fixed  $\alpha \in \mathbb{R}$ , there is an extension of this net on the space  $\bigoplus_{j \in \mathbb{Z}} \mathcal{H}_{j\alpha} \otimes \mathcal{H}_{j\alpha}$  [MTW18, AGT]: on each direct summand  $\mathcal{H}_{j\alpha} \otimes \mathcal{H}_{j\alpha}$ , the tensor product net  $\mathcal{A}_0$  acts by the representation  $\rho_{j\alpha} \otimes \rho_{j\alpha}$ .

Furthermore, let  $Y_{\alpha,s}$  as above on  $\hat{\mathcal{H}} = \bigoplus_{j \in \mathbb{Z}} \mathcal{H}_{j\alpha}$ . The components  $Y_{\alpha,s} \otimes \mathbb{1}$  of the charged field  $Y_\alpha(z) \otimes \mathbb{1}$  of the left chiral component acts on  $\hat{\mathcal{H}} \otimes \hat{\mathcal{H}}$  trivially on the right chiral component. Similarly, the components  $\mathbb{1} \otimes Y_{\alpha,s}$  of the charged field  $\mathbb{1} \otimes Y_\alpha(z)$  of the right chiral component acts on  $\hat{\mathcal{H}} \otimes \hat{\mathcal{H}}$ , trivially on the left chiral component.

We consider the combined charged field

$$\tilde{\psi}(w, z) = Y_\alpha(w) \otimes Y_\alpha(z) + (Y_\alpha(w) \otimes Y_\alpha(z))^*,$$

which is a formal series whose coefficients are operators acting on  $\hat{\mathcal{H}} \otimes \hat{\mathcal{H}}$ . Actually, as the components of  $Y_\alpha(w) \otimes Y_\alpha(z)$  raise (respectively the components of  $(Y_\alpha(w) \otimes Y_\alpha(z))^*$  lower) the left and right charges by  $\alpha$  at the same time, they restrict to  $\tilde{\mathcal{H}} = \bigoplus_{j \in \mathbb{Z}} \mathcal{H}_{j\alpha} \otimes \mathcal{H}_{j\alpha}$ . As we take  $\alpha < \frac{1}{\sqrt{2}} < 1$ , by [AGT, Theorem 5.9], the field  $\tilde{\psi}(w, z)$  is a two-dimensional conformal Wightman field that generates a two-dimensional Haag-Kastler net. Let us call this net  $\tilde{\mathcal{A}}$ .



## 4 Estimates of the charged fields

As we wish to perturb the conformal net by a charged field, we are interested in the time-zero restriction of  $\psi(w, z) = Y_\alpha(w) \otimes Y_\alpha(z) + (Y_\alpha(w) \otimes Y_\alpha(z))^*$ . This amounts to take  $z = w^{-1}$ . However, it is *a priori* unclear whether this is possible, because taking  $z = w^{-1}$  should give a formal series of  $z$  alone, but each component is an infinite sum of components of  $\tilde{\psi}(w, z)$ :

$$Y_\alpha(w) \otimes Y_\alpha(z) = \sum_{s \in \mathbb{R}} Y_{\alpha,s} w^{-s-d} \otimes \sum_{t \in \mathbb{R}} Y_{\alpha,t} z^{-t-d}$$

Therefore, by substituting  $z = w^{-1}$ ,

$$\begin{aligned} Y_\alpha(w) \otimes Y_\alpha(w^{-1}) &= \sum_{s \in \mathbb{R}} Y_{\alpha,s} w^{-s-d} \otimes \sum_{t \in \mathbb{R}} Y_{\alpha,t} w^{t+d} \\ &= \sum_{s \in \mathbb{R}} \sum_{t \in \mathbb{R}} Y_{\alpha,s+t} \otimes Y_{\alpha,t} w^s \end{aligned}$$

and we have to make sure that the sum  $\sum_{t \in \mathbb{R}} Y_{\alpha,s+t} \otimes Y_{\alpha,t}$  (the sum is countable on each  $\mathcal{H}_{j\alpha} \otimes \mathcal{H}_{j\alpha}$ ) gives a finite result on a certain dense domain. As the domain we take  $\bigoplus_{j,\text{alg}} \mathcal{H}_{j\alpha}^{\text{fin}} \otimes_{\text{alg}} \mathcal{H}_{j\alpha}^{\text{fin}}$  where  $\bigoplus_{j,\text{alg}}$  denotes the algebraic direct sum and  $\otimes_{\text{alg}}$  denotes the algebraic tensor product. We show that the above sum is convergent if  $|\alpha| < \frac{1}{\sqrt{2}}$ .

For this purpose, we need a general result on primary fields with conformal weight  $d$  [CKLW18, Appendix B (141)]. This is proven for fields with integer  $d$ , but it is straightforward to generalize it (because one only needs the primarity). It states that

$$\|Y_{\alpha,-n-d}\Omega\|^2 = \binom{2d+n-1}{n} = \frac{\Gamma(2d+n)}{\Gamma(n+1)\Gamma(2d)} \sim n^{2d-1}, \quad (5)$$

where the last asymptotic follows from the Stirling's approximation of the Gamma function  $\Gamma(x)$  with complex variable [WW21, 12.33].

We first observe that, for  $|\alpha| \geq \frac{1}{\sqrt{2}}$ ,  $d = \frac{\alpha^2}{2} \geq \frac{1}{4}$ , thus  $\sum_{n \in \mathbb{Z}} Y_{\alpha,-n} \otimes Y_{\alpha,s-n}$  does not converge on  $\Omega_0 \otimes \Omega_0$ . Indeed,  $\|Y_{\alpha,-n-d} \otimes Y_{\alpha,s-n-d} \cdot \Omega_0 \otimes \Omega_0\|^2 = \binom{2d+n-1}{n} \binom{2d+n-s-1}{n-s} \sim n^{4d-2}$  (for a fixed  $s$ , as  $n \rightarrow \infty$ ) and these vectors are orthogonal to each other, hence the sum  $\sum_{n \in \mathbb{Z}} Y_{\alpha,-n-d} \otimes Y_{\alpha,s-n-d}$  diverges on  $\Omega_0 \otimes \Omega_0$ . This means that, if  $|\alpha| \geq \frac{1}{\sqrt{2}}$ , there is no hope to define an operator of the form  $\sum_n Y_{\alpha,-n-d} \otimes Y_{\alpha,s-n-d}$  on a domain containing  $\Omega_0 \otimes \Omega_0$ .

On the other hand, if  $d < \frac{1}{4}$ , there is a hope to carry through the general programme of Section 2.3.

**Theorem 4.1.** *Let  $|\alpha| < \frac{1}{\sqrt{2}}$ . Then the coefficients of  $\tilde{\psi}(w, w^{-1})$  of  $w$  is a convergent sum on any vector in  $\bigoplus_{j,\text{alg}} \mathcal{H}_{j\alpha}^{\text{fin}} \otimes_{\text{alg}} \mathcal{H}_{j\alpha}^{\text{fin}}$ .*

*Proof.* We note first that a general vector in one tensor component is a linear combination of  $c_{j\alpha} \hat{J}_{-m_1} \cdots \hat{J}_{-m_k} \Omega_0$ , where  $m_\ell > 0$  and  $c_{j\alpha}$  commutes with  $\hat{J}_m$ ,  $m \neq 0$ . Therefore, it is enough to prove the convergence on vectors in  $\mathcal{H}_0^{\text{fin}} \otimes_{\text{alg}} \mathcal{H}_0^{\text{fin}}$ .

We claim that  $\|Y_{\alpha,-n-d} \hat{J}_{-m_1} \cdots \hat{J}_{-m_k} \Omega_0\|^2 \leq C_1(n + C_2)^{4d-2}$  where  $C_1, C_2$  depend on the vector but not on  $n$ . This is clear for  $k = 0$ . To prove the claim by induction on  $k$ , let us

observe that  $Y_{\alpha,-n-d}\hat{J}_{-m_1}\cdots\hat{J}_{-m_k}\Omega_0$  can be reduced, using the commutation relations (4),  $[Y_{\alpha,-n-d}, \hat{J}_m] = -\alpha Y_{\alpha,-n+m-d}$ , as follows:

$$\begin{aligned} Y_{\alpha,-n-d}\hat{J}_{-m_1}\cdots\hat{J}_{-m_k}\Omega_0 &= ([Y_{\alpha,-n-d}, \hat{J}_{-m_1}] + \hat{J}_{-m_1}Y_{\alpha,-n-d})\hat{J}_{-m_2}\cdots\hat{J}_{-m_k}\Omega_0 \\ &= (-\alpha Y_{\alpha,-n+m_1-d} + \hat{J}_{-m_1}Y_{\alpha,-n-d})\hat{J}_{-m_2}\cdots\hat{J}_{-m_k}\Omega_0 \end{aligned}$$

Using  $\|\Psi_1 + \Psi_2\|^2 \leq 2(\|\Psi_1\|^2 + \|\Psi_2\|^2)$ , it is enough to show that the norm of each term decays as desired. The first term decays by the induction hypothesis. Let us calculate the norm of the second term:

$$\begin{aligned} &\|\hat{J}_{-m_1}Y_{\alpha,-n-d}\hat{J}_{-m_2}\cdots\hat{J}_{-m_k}\Omega_0\|^2 \\ &= \langle Y_{\alpha,-n-d}\hat{J}_{-m_2}\cdots\hat{J}_{-m_k}\Omega_0, ([\hat{J}_{m_1}, \hat{J}_{-m_1}] + \hat{J}_{-m_1}\hat{J}_{m_1})Y_{\alpha,-n-d}\hat{J}_{-m_2}\cdots\hat{J}_{-m_k}\Omega_0 \rangle \\ &= \langle Y_{\alpha,-n-d}\hat{J}_{-m_2}\cdots\hat{J}_{-m_k}\Omega_0, (m_1 + \hat{J}_{-m_1}\hat{J}_{m_1})Y_{\alpha,-n-d}\hat{J}_{-m_2}\cdots\hat{J}_{-m_k}\Omega_0 \rangle \\ &= m_1\|Y_{\alpha,-n-d}\hat{J}_{-m_2}\cdots\hat{J}_{-m_k}\Omega_0\|^2 + \|\hat{J}_{m_1}Y_{\alpha,-n-d}\hat{J}_{-m_2}\cdots\hat{J}_{-m_k}\Omega_0\|^2 \\ &= m_1\|Y_{\alpha,-n-d}\hat{J}_{-m_2}\cdots\hat{J}_{-m_k}\Omega_0\|^2 + \|(\alpha Y_{\alpha,-n+m_1-d} + Y_{\alpha,-n-d}\hat{J}_{m_1})\hat{J}_{-m_2}\cdots\hat{J}_{-m_k}\Omega_0\|^2 \\ &\leq m_1\|Y_{\alpha,-n-d}\hat{J}_{-m_2}\cdots\hat{J}_{-m_k}\Omega_0\|^2 \\ &\quad + 2\|\alpha Y_{\alpha,-n+m_1-d}\hat{J}_{-m_2}\cdots\hat{J}_{-m_k}\Omega_0\|^2 + 2\|Y_{\alpha,-n-d}\hat{J}_{m_1}\hat{J}_{-m_2}\cdots\hat{J}_{-m_k}\Omega_0\|^2. \end{aligned}$$

The last term can be further reduced, using  $[\hat{J}_m, \hat{J}_n] = m\delta_{m,-n}$  and  $\hat{J}_m\Omega_0$  (as  $m > 0$ ), to a sum of norms of vectors of the above form. This completes the induction. That is, the norm of  $Y_{\alpha,-n-d}\hat{J}_{-m_1}\cdots\hat{J}_{-m_k}\Omega_0$  is a linear combination of terms that decay like  $(n + C_2)^{2d-1}$ .

When the operator  $Y_{\alpha,-n-d} \otimes Y_{\alpha,s-n-d}$  for a fixed  $s$  is applied to a vector that is the tensor product of two such vectors, the norm decays as  $C_1(n + C_2)^{4d-2}$ , which is summable in  $n$ . Therefore, this operator is defined on  $\bigoplus_{j,\text{alg}} \mathcal{H}_{j\alpha}^{\text{fin}} \otimes_{\text{alg}} \mathcal{H}_{j\alpha}^{\text{fin}}$ .  $\square$

## 5 Commutativity of the time-zero charged field

Now we know that, for  $|\alpha| < \frac{1}{\sqrt{2}}$ ,  $Y_\alpha(w) \otimes Y_\alpha(w^{-1})$  makes sense as a formal series whose coefficients are (unbounded) operators on the dense domain  $\bigoplus_{j,\text{alg}} \mathcal{H}_{j\alpha}^{\text{fin}} \otimes \mathcal{H}_{j\alpha}^{\text{fin}}$ . Next we show that it is not only local but also commutative, and moreover,  $Y_\alpha(w) \otimes Y_\alpha(w^{-1})$  and  $Y_\beta(z) \otimes Y_\beta(z^{-1})$  commute for possibly different  $\alpha, \beta$ . More precisely, we have the following result.

**Theorem 5.1.** *As formal series,  $Y_\alpha(w) \otimes Y_\alpha(w^{-1})$  and  $Y_\beta(z) \otimes Y_\beta(z^{-1})$  commute on the domain  $\bigoplus_{j,\text{alg}} \mathcal{H}_{j\alpha}^{\text{fin}} \otimes_{\text{alg}} \mathcal{H}_{j\alpha}^{\text{fin}}$  weakly, that is, as sesquilinear forms.*

*Proof.* We have  $[\hat{J}_m, Y_{\alpha,s}] = \alpha Y_{\alpha,m+s}$ . In terms of formal series, this amounts to  $[Y_\alpha(w), \hat{J}_m] = -\alpha Y_\alpha(w)w^m$ ,

Let us denote by  $\hat{J}_m \otimes \mathbb{1}$  the operator on  $\bigoplus_{j,\text{alg}} \mathcal{H}_{j\alpha}^{\text{fin}} \otimes_{\text{alg}} \mathcal{H}_{j\alpha}^{\text{fin}}$  that acts as  $\hat{J}_{j\alpha,m} \otimes \mathbb{1}$  on each component  $\mathcal{H}_{j\alpha}^{\text{fin}} \otimes_{\text{alg}} \mathcal{H}_{j\alpha}^{\text{fin}}$ . We calculate the commutator as a sesquilinear form, that is, we apply the operators to a vector and take a scalar product with another vector, but we

omit them. It holds that, by the Jacobi identity,

$$\begin{aligned}
& -[[Y_\alpha(w) \otimes Y_\alpha(w^{-1}), Y_\beta(z) \otimes Y_\beta(z^{-1})], \hat{J}_m \otimes \mathbb{1}] \\
& = [[\hat{J}_m \otimes \mathbb{1}, Y_\alpha(w) \otimes Y_\alpha(w^{-1})], Y_\beta(z) \otimes Y_\beta(z^{-1})] \\
& \quad + [[Y_\beta(z) \otimes Y_\beta(z^{-1}), \hat{J}_m \otimes \mathbb{1}], Y_\alpha(w) \otimes Y_\alpha(w^{-1})] \\
& = (-\alpha z^m - \beta \zeta^m)[Y_\alpha(w) \otimes Y_\alpha(w^{-1}), Y_\beta(z) \otimes Y_\beta(z^{-1})],
\end{aligned}$$

and similarly, with  $\mathbb{1} \otimes \hat{J}_m$  that acts as  $\mathbb{1} \otimes \hat{J}_m$  on each component  $\mathcal{H}_{j\alpha}^{\text{fin}} \otimes_{\text{alg}} \mathcal{H}_{j\alpha}^{\text{fin}}$ ,

$$\begin{aligned}
& -[[Y_\alpha(w) \otimes Y_\alpha(w^{-1}), Y_\beta(z) \otimes Y_\beta(z^{-1})], \mathbb{1} \otimes \hat{J}_m] \\
& = (-\alpha z^{-m} - \beta \zeta^{-m})[Y_\alpha(w) \otimes Y_\alpha(w^{-1}), Y_\alpha(z) \otimes Y_\alpha(z^{-1})].
\end{aligned}$$

We observe that, upon commuting with  $\hat{J}_m \otimes \mathbb{1}$  or  $\mathbb{1} \otimes \hat{J}_m$ , we obtain the same operator  $[Y_\alpha(w) \otimes Y_\alpha(w^{-1}), Y_\alpha(z) \otimes Y_\alpha(z^{-1})]$  multiplied by a scalar.

Now, to show that the commutator  $[Y_\alpha(w) \otimes Y_\alpha(w^{-1}), Y_\alpha(z) \otimes Y_\alpha(z^{-1})]$  vanishes, we only have to check that the matrix components vanishes. It is easy to check that  $Y_\alpha(w) \otimes Y_\alpha(w^{-1})$  commutes with  $c_\alpha \otimes c_\alpha$ , therefore, we only have to consider the pair of vectors in  $\mathcal{H}_0 \otimes \mathcal{H}_0$  and  $\mathcal{H}_\alpha \otimes \mathcal{H}_\alpha$ . Furthermore, due to the above commutation relations, the scalar product

$$\langle \hat{J}_{-m_1} \cdots \hat{J}_{-m_k} \Omega_{2\alpha} \otimes \hat{J}_{-n_1} \cdots \hat{J}_{-n_\ell} \Omega_{2\alpha}, \quad \hat{J}_{-m'_1} \cdots \hat{J}_{-m'_{k'}} \Omega_0 \otimes \hat{J}_{-n'_1} \cdots \hat{J}_{-n'_{\ell'}} \Omega_0 \rangle$$

can be reduced to the case  $\langle \Omega_{2\alpha} \otimes \Omega_{2\alpha}, \cdot \Omega_0 \otimes \Omega_0 \rangle$ .

Let us put  $\underline{Y}_\alpha(z) = E^-(\alpha, z)E^+(\alpha, z)$  and  $\tilde{\underline{Y}}_\alpha(z) = \underline{Y}_\alpha(z) \otimes \underline{Y}_\alpha(z^{-1})$ . As we have  $E^+(\alpha, z)^* = E^-(-\alpha, z)$  (with the convention that  $z^* = z^{-1}$ ), it follows that  $\underline{Y}_\alpha(z)^* = \underline{Y}_{-\alpha}(z)$  and  $\tilde{\underline{Y}}_\alpha(z)^* = \tilde{\underline{Y}}_{-\alpha}(z)$ , or equivalently  $\tilde{\underline{Y}}_{m,\alpha}^\dagger = \tilde{\underline{Y}}_{-m,-\alpha}$ .

Put  $\tilde{\underline{Y}}_\alpha(z) = \sum_m \tilde{\underline{Y}}_{\alpha,m} z^{-m} = \sum_m \sum_k \underline{Y}_{\alpha,k} \otimes \underline{Y}_{\alpha,k+m} z^{-m}$ . Then we have

$$\begin{aligned}
Y_\alpha(z) \otimes Y_\alpha(z^{-1}) & = (c_\alpha \otimes c_\alpha) z^{\alpha J_0} z^{-\alpha J_0} \underline{Y}_\alpha(z) \otimes \underline{Y}_\alpha(z^{-1}) \\
& = (c_\alpha \otimes c_\alpha) \tilde{\underline{Y}}_\alpha(z).
\end{aligned}$$

Therefore, the question is further reduced to  $\langle \Omega_0 \otimes \Omega_0, [\tilde{\underline{Y}}_{\alpha,m}, \tilde{\underline{Y}}_{\beta,n}] \Omega_0 \otimes \Omega_0 \rangle = 0$  for all  $m, n$ .

Note that, with  $F$  the flip operator between the left and right tensor components in  $\mathcal{H}_0 \otimes \mathcal{H}_0$  (which is a unitary operator),

$$\begin{aligned}
\tilde{\underline{Y}}_{\alpha,m} \Omega_0 \otimes \Omega_0 & = \sum_k \underline{Y}_{\alpha,k} \otimes \underline{Y}_{\alpha,k+m} \Omega_0 \otimes \Omega_0 \\
& = F \cdot \sum_k \underline{Y}_{\alpha,k} \otimes \underline{Y}_{\alpha,k-m} \Omega_0 \otimes \Omega_0 = F \cdot \tilde{\underline{Y}}_{\alpha,-m} \Omega_0 \otimes \Omega_0.
\end{aligned}$$

This implies that

$$\begin{aligned}
\langle \tilde{\underline{Y}}_{\alpha,m} \Omega_0 \otimes \Omega_0, \tilde{\underline{Y}}_{\beta,n} \Omega_0 \otimes \Omega_0 \rangle & = \langle F \cdot \tilde{\underline{Y}}_{\alpha,-m} \Omega_0 \otimes \Omega_0, F \cdot \tilde{\underline{Y}}_{\beta,-n} \Omega_0 \otimes \Omega_0 \rangle \\
& = \langle \tilde{\underline{Y}}_{\alpha,-m} \Omega_0 \otimes \Omega_0, \tilde{\underline{Y}}_{\beta,-n} \Omega_0 \otimes \Omega_0 \rangle.
\end{aligned}$$

Moreover, note that the map  $J_m \mapsto -J_m$  is a vacuum-preserving automorphism implemented by a unitary (the multiplication by  $(-1)^k$  on the  $k$ -particle space), and it's tensor product maps

$$\begin{aligned}\tilde{Y}_\alpha(z) &= E^-(\alpha, z)E^+(\alpha, z) \otimes E^-(\alpha, z^{-1})E^+(\alpha, z^{-1}) \\ &\mapsto E^-(-\alpha, z)E^+(-\alpha, z) \otimes E^-(-\alpha, z^{-1})E^+(-\alpha, z^{-1}) = \tilde{Y}_{-\alpha}(z).\end{aligned}$$

That is,  $\tilde{Y}_{\alpha, m}$  is mapped to  $\tilde{Y}_{-\alpha, m}$ , therefore, by the invariance of  $\Omega_0 \otimes \Omega_0$  by this unitary,

$$\langle \tilde{Y}_{\alpha, m} \Omega_0 \otimes \Omega_0, \tilde{Y}_{\beta, n} \Omega_0 \otimes \Omega_0 \rangle = \langle \tilde{Y}_{-\alpha, m} \Omega_0 \otimes \Omega_0, \tilde{Y}_{-\beta, n} \Omega_0 \otimes \Omega_0 \rangle.$$

From this, we can compute the commutator (as a sesquilinear form)

$$\begin{aligned}\langle \Omega_0 \otimes \Omega_0, [\tilde{Y}_{\alpha, m}, \tilde{Y}_{\beta, n}] \Omega_0 \otimes \Omega_0 \rangle &= \langle \Omega_0 \otimes \Omega_0, (\tilde{Y}_{\alpha, m} \tilde{Y}_{\beta, n} - \tilde{Y}_{\beta, n} \tilde{Y}_{\alpha, m}) \Omega_0 \otimes \Omega_0 \rangle \\ &= \langle \tilde{Y}_{-\alpha, -m} \Omega_0 \otimes \Omega_0, \tilde{Y}_{\beta, n} \Omega_0 \otimes \Omega_0 \rangle - \langle \tilde{Y}_{-\beta, -n} \Omega_0 \otimes \Omega_0, \tilde{Y}_{\alpha, m} \Omega_0 \otimes \Omega_0 \rangle \\ &= \langle \tilde{Y}_{-\alpha, m} \Omega_0 \otimes \Omega_0, \tilde{Y}_{\beta, -n} \Omega_0 \otimes \Omega_0 \rangle - \langle \tilde{Y}_{-\beta, n} \Omega_0 \otimes \Omega_0, \tilde{Y}_{\alpha, m} \Omega_0 \otimes \Omega_0 \rangle \\ &= \langle \tilde{Y}_{\alpha, m} \Omega_0 \otimes \Omega_0, \tilde{Y}_{-\beta, -n} \Omega_0 \otimes \Omega_0 \rangle - \langle \tilde{Y}_{-\beta, -n} \Omega_0 \otimes \Omega_0, \tilde{Y}_{\alpha, m} \Omega_0 \otimes \Omega_0 \rangle.\end{aligned}$$

Furthermore, by construction (3) of  $\tilde{Y}_m = \sum_{k \in \mathbb{R}} Y_{\alpha, k} \otimes Y_{\alpha, k+m}$ , these expectation values give only real numbers. Therefore, by hermitianity of the scalar product, this commutator vanishes on the vacuum state.  $\square$

## 6 Perturbation by charged fields

According to the general idea of Section 2, we wish to perturb the net  $\tilde{\mathcal{A}}$  by the field  $\tilde{\phi}(z)$  a la Barata-Jäkel-Mund [BJM23]. That is, while keeping the  $T^{11}$  component of the stress-energy tensor, we add a smeared local field to the  $T^{00}$  component on the time-zero circle  $S^1$ .

The necessary condition for it to work is that the new operators satisfy the Lorentz relations:

$$[\ell_m, \ell_n] = (m - n)\ell_{m, -n}, \quad m, n = -1, 0, 1.$$

As we do not know whether the smeared field can be multiplied on the domain of the old generators, we consider the weak commutation relation: for two vectors  $\Psi_1, \Psi_2$ , we compute  $\langle A^* \Psi_1, B \Psi_2 \rangle - \langle B^* \Psi_1, A \Psi_2 \rangle$ . Obviously, if the commutator  $[A, B]$  can be defined on the domain and calculated, then it implies the weak commutation relation.

We pick the symmetric field

$$\tilde{\psi}(w, z) = Y_\alpha(w) \otimes Y_\alpha(z) + (Y_\alpha(w) \otimes Y_\alpha(z))^* = Y_\alpha(w) \otimes Y_\alpha(z) + Y_{-\alpha}(w) \otimes Y_{-\alpha}(z),$$

restrict it to the time-zero circle  $z = w^{-1}$  and smear it with  $\epsilon_j(\theta)$ ,  $n = -1, 0, 1$ . Correspondingly, we consider the coefficients of  $z^n$ :

$$\sum_{k \in \mathbb{R}} Y_{\alpha, k} \otimes Y_{\alpha, -n+k} + \sum_{s \in \mathbb{R}} Y_{-\alpha, s} \otimes Y_{-\alpha, -n+s} = \sum_{k \in \mathbb{R}, \epsilon = \pm 1} Y_{\epsilon \alpha, k} \otimes Y_{\epsilon \alpha, -n+k}.$$

**Theorem 6.1.** *The Lorentz relations are weakly satisfied for*

$$\begin{aligned} & \hat{L}_1 \otimes \mathbb{1} + \mathbb{1} \otimes \hat{L}_{-1} + \lambda \sum_{k \in \mathbb{R}, \epsilon = \pm 1} Y_{\epsilon\alpha, k} \otimes Y_{\epsilon\alpha, -1+k} \\ & \hat{L}_0 \otimes \mathbb{1} - \mathbb{1} \otimes \hat{L}_0, \\ & \hat{L}_{-1} \otimes \mathbb{1} + \mathbb{1} \otimes \hat{L}_1 + \lambda \sum_{k \in \mathbb{R}, \epsilon = \pm 1} Y_{\epsilon\alpha, k} \otimes Y_{\epsilon\alpha, 1+k} \end{aligned}$$

where  $k$  runs in  $\mathbb{R}$ , but there are only countable nonzero terms on each  $\mathcal{H}_\beta \otimes \mathcal{H}_\beta$ , on the domain  $\bigoplus_{j, \text{alg}} \mathcal{H}_{j\alpha}^{\text{fin}} \otimes_{\text{alg}} \mathcal{H}_{j\alpha}^{\text{fin}}$ .

*Proof.* It is clear that, if  $\lambda = 0$ , these are the old generators and they satisfy the Lorentz relation on the domain. The new terms commute with each other in the weak sense as we have seen in Section 5, hence we only have to check the commutation relations between the old terms and the new terms.

One new term can be applied to a vector in the domain and gives a convergent series, hence we can compute the weak commutator term by term. For a primary field  $Y_\alpha(z)$ , it holds that  $[\hat{L}_m, Y_{\alpha, n}] = ((d-1)m - n)Y_{\alpha, m+n}$  in the operator sense, and hence also in the weak sense. Therefore, we have the following commutation relation in the weak sense:

$$\begin{aligned} & \left[ \hat{L}_m \otimes \mathbb{1} + \mathbb{1} \otimes \hat{L}_{-m}, \sum_{k \in \mathbb{R}} Y_{\alpha, k} \otimes Y_{\alpha, -n+k} \right] \\ &= \sum_{k \in \mathbb{R}} (((d-1)m - k)Y_{\alpha, k+m} \otimes Y_{\alpha, -n+k} + ((d-1)(-m) - (-n+k))Y_{\alpha, k} \otimes Y_{\alpha, -m-n+k}) \\ &= \sum_{k \in \mathbb{R}} (((d-1)m - (k-m))Y_{\alpha, k} \otimes Y_{\alpha, -m-n+k} \\ & \quad + ((d-1)(-m) - (-n+k))Y_{\alpha, k} \otimes Y_{\alpha, -m-n+k}) \\ &= \sum_{k \in \mathbb{R}} (m+n-2k)Y_{\alpha, k} \otimes Y_{\alpha, -m-n+k}. \end{aligned}$$

As  $\alpha$  is arbitrary, this holds even if  $\alpha$  is replaced by  $-\alpha$ . Furthermore,  $\sum_{k \in \mathbb{R}} Y_{\alpha, k} \otimes Y_{\alpha, -n+k}$  and  $\sum_{k \in \mathbb{R}} Y_{-\alpha, k} \otimes Y_{-\alpha, -n+k}$  commute by Theorem 5.1. Altogether,

$$\begin{aligned} & \left[ \hat{L}_m \otimes \mathbb{1} + \mathbb{1} \otimes \hat{L}_{-m} + \lambda \sum_{k \in \mathbb{R}, \epsilon = \pm 1} Y_{\epsilon\alpha, k} \otimes Y_{\epsilon\alpha, -m+k}, \right. \\ & \quad \left. \hat{L}_n \otimes \mathbb{1} + \mathbb{1} \otimes \hat{L}_{-n} + \lambda \sum_{k \in \mathbb{R}, \epsilon = \pm 1} Y_{\epsilon\alpha, k} \otimes Y_{\epsilon\alpha, -n+k} \right] \\ &= (m-n)\hat{L}_{m+n} \otimes \mathbb{1} - (m-n)\mathbb{1} \otimes \hat{L}_{-m-n} \\ & \quad + \lambda \sum_{k \in \mathbb{R}, \epsilon = \pm 1} (m+n-2k)Y_{\epsilon\alpha, k} \otimes Y_{\epsilon\alpha, -m-n+k} - \lambda \sum_{k \in \mathbb{R}, \epsilon = \pm 1} (n+m-2k)Y_{\epsilon\alpha, k} \otimes Y_{\epsilon\alpha, -n-m+k} \\ &= (m-n)\hat{L}_{m+n} \otimes \mathbb{1} - (m-n)\mathbb{1} \otimes \hat{L}_{-m-n}, \end{aligned}$$

and for  $m = 1, n = -1$ , this is  $2(\hat{L}_0 \otimes \mathbb{1} - \mathbb{1} \otimes \hat{L}_0)$ .

On the other hand,

$$\begin{aligned}
& \left[ \hat{L}_0 \otimes \mathbb{1} - \mathbb{1} \otimes \hat{L}_0, \hat{L}_m \otimes \mathbb{1} + \mathbb{1} \otimes \hat{L}_{-m} + \lambda \sum_{k \in \mathbb{R}, \epsilon = \pm 1} Y_{\epsilon\alpha, k} \otimes Y_{\epsilon\alpha, -m+k} \right] \\
&= (-m) \hat{L}_m \otimes \mathbb{1} - (m) \mathbb{1} \otimes \hat{L}_{-m} \\
&\quad + \lambda \sum_{k \in \mathbb{R}, \epsilon = \pm 1} ((-k) Y_{\epsilon\alpha, k} \otimes Y_{\epsilon\alpha, -m+k} - (-(-m+k)) Y_{\epsilon\alpha, k} \otimes Y_{\epsilon\alpha, -m+k}) \\
&= (-m) \left( \hat{L}_m \otimes \mathbb{1} + \mathbb{1} \otimes \hat{L}_{-m} + \lambda \sum_{k \in \mathbb{R}, \epsilon = \pm 1} Y_{\epsilon\alpha, k} \otimes Y_{\epsilon\alpha, -n+k} \right).
\end{aligned}$$

By combining its adjoint as before, we obtain the right commutation relations between  $\ell_0$  and  $\ell_m$  for  $m = 1, -1$ .  $\square$

Note that the Lorentz relations do not extend beyond  $m = 1, 0, -1$ , that is, they do not satisfy the Virasoro relations.

In order to implement the perturbation by this commutative field, we need to solve the following.

**Problems.** Show that the above generators are self-adjoint on a certain domain and generate a dynamics that satisfies finite speed of propagation.

On the other hand, on the time-zero circle, there is a new representation of the Virasoro algebra (non-positive energy) with  $c = 0$ , or the Witt algebra.

**Proposition 6.2.** *The Virasoro relations are weakly satisfied with  $c = 0$  for*

$$\hat{L}_m \otimes \mathbb{1} - \mathbb{1} \otimes \hat{L}_{-m} + i\lambda m \sum_{k \in \mathbb{R}, \epsilon = \pm 1} Y_{\epsilon\alpha, k} \otimes Y_{\epsilon\alpha, -m+k},$$

where  $k$  runs in  $\mathbb{R}$ , but there are only countable nonzero terms on each  $\mathcal{H}_\beta \otimes \mathcal{H}_\beta$ .

*Proof.* As before, we compute the commutation relations weakly. First, with  $d = \frac{\alpha^2}{2} = \frac{(-\alpha)^2}{2}$ ,

$$\begin{aligned}
& \left[ \hat{L}_m \otimes \mathbb{1} - \mathbb{1} \otimes \hat{L}_{-m}, \sum_{k \in \mathbb{R}, \epsilon = \pm 1} Y_{\epsilon\alpha, k} \otimes Y_{\epsilon\alpha, -n+k} \right] \\
&= \sum_{k \in \mathbb{R}, \epsilon = \pm 1} (((d-1)m - k) Y_{\epsilon\alpha, k+m} \otimes Y_{\epsilon\alpha, -n+k} \\
&\quad - ((d-1)(-m) - (-n+k)) Y_{\epsilon\alpha, k} \otimes Y_{\epsilon\alpha, -m-n+k}) \\
&= \sum_{k \in \mathbb{R}, \epsilon = \pm 1} (((d-1)m - (k-m)) Y_{\epsilon\alpha, k} \otimes Y_{\epsilon\alpha, -m-n+k} \\
&\quad - ((d-1)(-m) - (-n+k)) Y_{\epsilon\alpha, k} \otimes Y_{\epsilon\alpha, -m-n+k}) \\
&= \sum_{k \in \mathbb{R}, \epsilon = \pm 1} ((2d-1)m - n) Y_{\epsilon\alpha, k} \otimes Y_{\epsilon\alpha, -m-n+k}
\end{aligned}$$

and by combining this and the one with  $\alpha$  replaced by  $-\alpha$ ,

$$\begin{aligned}
& \left[ \hat{L}_m \otimes \mathbb{1} - \mathbb{1} \otimes \hat{L}_{-m} + i\lambda m \sum_{k \in \mathbb{R}, \epsilon = \pm 1} Y_{\epsilon\alpha, k} \otimes Y_{\epsilon\alpha, -m+k}, \right. \\
& \quad \left. \hat{L}_n \otimes \mathbb{1} - \mathbb{1} \otimes \hat{L}_{-n} + i\lambda n \sum_{k \in \mathbb{R}, \epsilon = \pm 1} Y_{\epsilon\alpha, k} \otimes Y_{\epsilon\alpha, -n+k} \right] \\
&= (m-n) \hat{L}_{m+n} \otimes \mathbb{1} + (-m-n) \mathbb{1} \otimes \hat{L}_{-m-n} \\
& \quad + i\lambda \sum_{k \in \mathbb{R}, \epsilon = \pm 1} (n((2d-1)m-n) - m((2d-1)n-m)) Y_{\epsilon\alpha, k} \otimes Y_{\epsilon\alpha, -m-n+k} \\
&= (m-n) \left( \hat{L}_{m+n} \otimes \mathbb{1} - \mathbb{1} \otimes \hat{L}_{-m-n} + i\lambda(m+n) \sum_{k \in \mathbb{R}, \epsilon = \pm 1} Y_{\epsilon\alpha, k} \otimes Y_{\epsilon\alpha, -m-n+k} \right).
\end{aligned}$$

□

The combination  $i\lambda m$  means that we are taking the derivative  $-i\partial_\theta \tilde{\psi}(e^{i\theta}, e^{-i\theta})$ . The operators  $\hat{L}_m \otimes \mathbb{1} - \mathbb{1} \otimes \hat{L}_{-m}$  is the time-zero Virasoro (Witt) algebra, and this Proposition tells that there are different actions of the Virasoro algebra with  $c = 0$ .

**Proposition 6.3.** *The Virasoro relations are weakly satisfied with  $c = 0$  for*

$$\hat{L}_m \otimes \mathbb{1} - \mathbb{1} \otimes \hat{L}_{-m} + \lambda \sum_{k \in \mathbb{R}, \epsilon = \pm 1} Y_{\epsilon\alpha, k} \otimes Y_{\epsilon\alpha, -m+k}$$

if and only if  $d = \frac{1}{2}$ .

*Proof.* As before, we compute the commutation relations weakly. In Proposition 6.2 we have seen

$$\left[ \hat{L}_m \otimes \mathbb{1} - \mathbb{1} \otimes \hat{L}_{-m}, \sum_{k \in \mathbb{R}, \epsilon = \pm 1} Y_{\epsilon\alpha, k} \otimes Y_{\epsilon\alpha, -n+k} \right] = \sum_{k \in \mathbb{R}, \epsilon = \pm 1} ((2d-1)m-n) Y_{\epsilon\alpha, k} \otimes Y_{\epsilon\alpha, -m-n+k}$$

and by combining this,

$$\begin{aligned}
& \left[ \hat{L}_m \otimes \mathbb{1} - \mathbb{1} \otimes \hat{L}_{-m} + \lambda \sum_{k \in \mathbb{R}, \epsilon = \pm 1} Y_{\epsilon\alpha, k} \otimes Y_{\epsilon\alpha, -m+k}, \right. \\
& \quad \left. \hat{L}_n \otimes \mathbb{1} - \mathbb{1} \otimes \hat{L}_{-n} + \lambda \sum_{k \in \mathbb{R}, \epsilon = \pm 1} Y_{\epsilon\alpha, k} \otimes Y_{\epsilon\alpha, -n+k} \right] \\
&= (m-n) \hat{L}_{m+n} \otimes \mathbb{1} + (-m-n) \mathbb{1} \otimes \hat{L}_{-m-n} \\
& \quad + \lambda \sum_{k \in \mathbb{R}, \epsilon = \pm 1} (((2d-1)m-n) - ((2d-1)n-m)) Y_{\epsilon\alpha, k} \otimes Y_{\epsilon\alpha, -m-n+k} \\
&= (m-n) \left( \hat{L}_{m+n} \otimes \mathbb{1} - \mathbb{1} \otimes \hat{L}_{-m-n} + 2d\lambda \sum_{k \in \mathbb{R}, \epsilon = \pm 1} Y_{\epsilon\alpha, k} \otimes Y_{\epsilon\alpha, -m-n+k} \right),
\end{aligned}$$

and the last expression in the bracket coincides with  $\hat{L}_{m+n} \otimes \mathbb{1} - \mathbb{1} \otimes \hat{L}_{-m-n} + \lambda \sum_{k \in \mathbb{R}, \epsilon = \pm 1} Y_{\epsilon\alpha, k} \otimes Y_{\epsilon\alpha, -m-n+k}$  if and only if  $d = \frac{1}{2}$ . □

The case  $d = \frac{1}{2}$  is related with the free fermion. This might indicate a hidden symmetry for the free fermion.

## 7 Outlook

We need that the above generators are self-adjoint on a certain domain. This is open. To show that they are essentially self-adjoint on our domain, one way would be to use the analytic vector theorem, but it is unclear whether even  $\Omega_0 \otimes \Omega_0$  is an analytic vector. Therefore, we need better estimates of the time-zero restriction  $Y_\alpha(w) \otimes Y_\alpha(w^{-1})$ . Such estimates will be needed also to show that the perturbed Lorentz generators do generate a new representation of the Lorentz group, and that we can construct a new Haag-Kastler net on  $dS^2$  and to find the interacting vacuum. For this purpose, studying the Euclidean models of these two-dimensional CFT might help.

There are many two-dimensional CFTs and some of the charged primary fields have been relatively well-understood. It might be a good idea to take other models where charged fields allow better control.

## Acknowledgements

Y.T. thanks Bin Gui for useful discussions on primary fields and estimates. Y.T. is partially supported by the *MIUR Excellence Department Project MatMod@TOV* awarded to the Department of Mathematics, University of Rome Tor Vergata CUP E83C23000330006 the University of Rome Tor Vergata funding *OAQM* CUP E83C22001800005 and by GNAMPA-INdAM.

## References

- [AGT] Maria Stella Adamo, Luca Giorgetti, and Yoh Tanimoto. Wightman fields for two-dimensional conformal field theories with pointed representation category. in preparation.
- [Bar54] V. Bargmann. On unitary ray representations of continuous groups. *Ann. of Math. (2)*, 59:1–46, 1954. <http://www.jstor.org/stable/1969831>.
- [BB99] H. J. Borchers and D. Buchholz. Global properties of vacuum states in de Sitter space. *Ann. Inst. H. Poincaré Phys. Théor.*, 70(1):23–40, 1999. <https://arxiv.org/abs/gr-qc/9803036>.
- [BGL93] R. Brunetti, D. Guido, and R. Longo. Modular structure and duality in conformal quantum field theory. *Comm. Math. Phys.*, 156(1):201–219, 1993. <http://projecteuclid.org/euclid.cmp/1104253522>.
- [BJM23] João C. A. Barata, Christian Jäkel, and Jens Mund. The  $\mathcal{P}(\varphi)_2$  Model on de Sitter Space. *Mem. Amer. Math. Soc.*, 281, 2023. <https://arxiv.org/abs/1607.02265>.



- [BMT88] Detlev Buchholz, Gerhard Mack, and Ivan Todorov. The current algebra on the circle as a germ of local field theories. *Nuclear Phys. B Proc. Suppl.*, 5B:20–56, 1988. <https://www.researchgate.net/publication/222585851>.
- [Car04] Sebastiano Carpi. On the representation theory of Virasoro nets. *Comm. Math. Phys.*, 244(2):261–284, 2004. <https://arxiv.org/abs/math/0306425>.
- [CKLW18] Sebastiano Carpi, Yasuyuki Kawahigashi, Roberto Longo, and Mihály Weiner. From vertex operator algebras to conformal nets and back. *Mem. Amer. Math. Soc.*, 254(1213):vi+85, 2018. <https://arxiv.org/abs/1503.01260>.
- [CTW19] Sebastiano Carpi, Yoh Tanimoto, and Mihály Weiner. Unitary representations of the  $\mathcal{W}_3$ -algebra with  $c \geq 2$ . 2019. <https://arxiv.org/abs/1910.08334>.
- [FHK75] Rodolfo Figari, Raphael Hoegh-Krohn, and Chiara R. Nappi. Interacting relativistic boson fields in the de Sitter universe with two space-time dimensions. *Comm. Math. Phys.*, 44(3):265–278, 1975. <http://projecteuclid.org/euclid.cmp/1103899347>.
- [GJ72] James Glimm and Arthur Jaffe. The  $\lambda\phi_2^4$  quantum field theory without cutoffs. IV. Perturbations of the Hamiltonian. *J. Mathematical Phys.*, 13:1568–1584, 1972. <https://doi.org/10.1063/1.1665879>.
- [GL03] Daniele Guido and Roberto Longo. A converse Hawking-Unruh effect and  $dS^2$ /CFT correspondence. *Ann. Henri Poincaré*, 4(6):1169–1218, 2003. <https://arxiv.org/abs/gr-qc/0212025>.
- [JM18] Christian D. Jäkel and Jens Mund. The Haag–Kastler Axioms for the  $\mathcal{P}(\varphi)_2$  Model on the De Sitter Space. *Annales Henri Poincaré*, 19(3):959–977, 2018. <https://arxiv.org/abs/1701.08231>.
- [Kac98] Victor Kac. *Vertex algebras for beginners*, volume 10 of *University Lecture Series*. American Mathematical Society, Providence, RI, second edition, 1998. <https://books.google.com/books?id=e-jxBwAAQBAJ>.
- [KL04] Yasuyuki Kawahigashi and Roberto Longo. Classification of two-dimensional local conformal nets with  $c < 1$  and 2-cohomology vanishing for tensor categories. *Comm. Math. Phys.*, 244(1):63–97, 2004. <https://arxiv.org/abs/math-ph/0304022>.
- [LR95] R. Longo and K.-H. Rehren. Nets of subfactors. *Rev. Math. Phys.*, 7(4):567–597, 1995. Workshop on Algebraic Quantum Field Theory and Jones Theory (Berlin, 1994). <https://arxiv.org/abs/hep-th/9411077>.
- [MT19] Vincenzo Morinelli and Yoh Tanimoto. Scale and Möbius covariance in two-dimensional Haag-Kastler net. *Comm. Math. Phys.*, 371(2):619–650, 2019. <https://arxiv.org/abs/1807.04707>.
- [MTW18] Vincenzo Morinelli, Yoh Tanimoto, and Mihály Weiner. Conformal covariance and the split property. *Comm. Math. Phys.*, 357(1):379–406, 2018. <https://arxiv.org/abs/1609.02196>.
- [Reh] Karl-Henning Rehren. Konforme quantenfeldtheorie. Lecture notes. <http://www.theorie.physik.uni-goettingen.de/~rehren/ps/cqft.pdf>.

- [Reh00] K.-H. Rehren. Chiral observables and modular invariants. *Comm. Math. Phys.*, 208(3):689–712, 2000. <https://arxiv.org/abs/hep-th/9903262>.
- [TL97] Valerio Toledano Laredo. Fusion of positive energy representations of  $\mathrm{LSpin}(2n)$ . 1997. Ph.D. Thesis, University of Cambridge. <https://arxiv.org/abs/math/0409044>.
- [TZ12] Michael P. Tuite and Alexander Zuevsky. A generalized vertex operator algebra for Heisenberg intertwiners. *J. Pure Appl. Algebra*, 216(6):1442–1453, 2012. <https://arxiv.org/abs/1106.6149>.
- [Wei05] Mihály Weiner. Conformal covariance and related properties of chiral QFT. 2005. Ph.D. thesis, Università di Roma “Tor Vergata”. <http://arxiv.org/abs/math/0703336>.
- [Wei11] Mihály Weiner. An algebraic version of Haag’s theorem. *Comm. Math. Phys.*, 305(2):469–485, 2011. <https://arxiv.org/abs/1006.4726>.
- [WW21] E. T. Whittaker and G. N. Watson. *A course of modern analysis—an introduction to the general theory of infinite processes and of analytic functions with an account of the principal transcendental functions*. Cambridge University Press, Cambridge, fifth edition, 2021. <https://doi.org/10.1017/9781009004091>.
- [Zam89] A. B. Zamolodchikov. Integrable field theory from conformal field theory. In *Integrable systems in quantum field theory and statistical mechanics*, volume 19 of *Adv. Stud. Pure Math.*, pages 641–674. Academic Press, Boston, MA, 1989. <https://inspirehep.net/literature/279986>.