

Modular operator for null plane algebras in free fields

Vincenzo Morinelli^{1,*}, Yoh Tanimoto^{2,△}, Benedikt Wegener^{2,•}.

¹Department Mathematik, FAU Erlangen-Nürnberg,
Cauerstraße 11 91058 Erlangen, Germany

² Dipartimento di Matematica, Università di Roma Tor Vergata
Via della Ricerca Scientifica 1, I-00133 Roma, Italy

July 2, 2021

Abstract

We consider the algebras generated by observables in quantum field theory localized in regions in the null plane. For a scalar free field theory, we show that the one-particle structure can be decomposed into a continuous direct integral of lightlike fibres, and the modular operator decomposes accordingly. This implies that a certain form of QNEC is valid in free fields involving the causal completions of half-spaces on the null plane (null cuts). We also compute the relative entropy of null cut algebras with respect to the vacuum and some coherent states.

Contents

1	Introduction	2
2	Preliminaries	4
2.1	Abstract one-particle structure	4
2.2	The one-particle structure of the free scalar field	6
2.3	Second quantization and nets of von Neumann algebras	8
2.4	The $U(1)$ -current net	9
3	Free scalar field on the null plane	10
3.1	Direct integrals and decompositions	10
3.2	Decomposition of the one-particle space	11
3.3	The local subspace of the null plane	13
3.4	Duality for null cuts	18
4	Modular operator for null cuts	19

*morinelli@math.fau.de, △ hoyt@mat.uniroma2.it, • wegener@mat.uniroma2.it

5	Relative entropy and Energy bounds	21
5.1	Decomposition of the relative entropy for coherent states	21
5.2	Relative entropy for coherent states on the null plane	23
5.3	The ANEC and the QNEC	24
6	Concluding remarks	28
A	Decomposable Functional calculus	29

1 Introduction

The modular Hamiltonian, or the (logarithm of the half of the) modular operator of local regions in quantum field theory (QFT), has been a focus of attention in recent years (see e.g. [CF20, Lon20, CTT17b]). On one hand, quantum-information aspects, such as the Bekenstein bound, generalized second law of thermodynamics and various null energy conditions, are a rare guidepost in the search of quantum gravity [Cas08, Wal12, FLPW16]. On the other hand, the modular theory of von Neumann algebras allows one to define relative entropy in QFT in a mathematically precise way [OP04], and various modular objects in QFT have been computed in concrete examples [LX18, Hol20, CLR20]. In particular, the modular operator of certain regions in the null plane has played an important role in relation with the quantum null energy condition (QNEC) and the averaged null energy condition (ANEC) [CTT17b, KLLSM18, CF20]. In these works, physicists consider a null cut, a region on the null plane defined by a spacelike curve C and have written a formula for the modular operator for the algebra of a null cut (see e.g. [CTT17b, (1.5)]):

$$\hat{H}_C = 2\pi \int d^{D-2} \mathbf{x}^\perp \int_{-\infty}^{\infty} d\lambda (\lambda - C(\mathbf{x}^\perp)) T_{++}(\lambda, \mathbf{x}^\perp), \quad (1)$$

where T_{++} is the lightlike component of the stress-energy tensor. This suggests that the inclusion of null cut algebras is a half-sided modular inclusion (HSMI) [Wie93]. Based on the latter assumption, a limited version of QNEC has been proved in [CF20]. Therefore, it is crucial to study modular object on the null plane.

Actually, the above formula must be interpreted with care: while it seems reasonable to assume that the stress-energy tensor is an operator-valued distribution (or even a Wightman field), it is unclear whether it can be restricted to a null plane (cf. [Ver00, FR03]). Furthermore, it is integrated against the unbounded function $\lambda - C(\mathbf{x}^\perp)$, that could be even more problematic. For these reasons, (1) cannot be considered directly as an expression for an operator on a Hilbert space. One of the goals of this paper is to partially justify (1) using the modular theory of von Neumann algebras in the case of the free fields.

We observe that the (scalar) free field can be restricted to the null plane, with a slight restriction to the test function. This has been known for a long time, and general properties of observables on the null plane have been studied [SS72, Dri77b, Dri77a, GLRV01, Ull04]. By the Bisognano-Wichmann property [BW76] and the Takesaki theorem [Tak03, Theorem IX.4.2], it is immediate that, if there are enough observables on the null plane, they split into a (continuous) tensor product along the transverse direction. This allows us to consider the

observables on each fibre on the null plane. These observables form a simplified quantum field theory on each lightlike fibre, and we can consider the modular objects there. We will show that the one-particle subspace \mathfrak{H}_m of the free field with mass m disintegrates as follows

$$\mathfrak{H}_m = \int_{\mathbb{R}^{D-1}}^{\oplus_{\mathbb{R}}} \mathcal{H}_{U(1)} d\mathbf{x}_{\perp}. \quad (2)$$

where $\mathcal{H}_{U(1)}$ is the one-particle space of the $U(1)$ -current sitting on the light ray $\{(t, t, \mathbf{x}_{\perp}) \in \mathbb{R}^{1+D} : t \in \mathbb{R}\}$. One can deduce that the modular operator of the region on the null plane is decomposed into the fibres, and its logarithm is written as a direct integral over fibres (see (19)):

$$\log(\Delta_{H(N_C)}) \simeq \int_{\mathbb{R}^{D-1}}^{\oplus} \left(\log(\Delta_{H_{U(1)}}(\mathbb{R}_+)) + 2\pi C(\mathbf{x}_{\perp}) P_{\mathbf{x}_{\perp}} \right) d\mathbf{x}_{\perp},$$

This is a clear analogue of (1), where the logarithm of the modular operator is expressed as the integral in the transverse direction of the stress-energy tensor smeared by the function $\lambda - C(\mathbf{x}_{\perp})$, where the latter is a formal expression for the shifted dilation operator in two-dimensional conformal field theory, which should coincide with the modular operator on each fibre. Our formula, while we avoid taking about the stress-energy tensor, makes explicit the idea that the modular operator decomposes into fibres. While the validity of (1) is believed more generally, we point out that in general there are not many observables that can be restricted to the null plane. We clarify the situation from interacting models in $(1+1)$ -dimensions. Our formula allows a covariant action of such distorted dilations the null plane and, for null cuts with continuous boundary C , on distorted wedge region $W_C = N_C''$.

In the course of the proof (Proposition 4.1), we show that inclusions of the null cut regions are HSMI. By [CF20], this completes the limited version of QNEC as in (24) in the case of the free scalar field (note that the result of [CF20] is based on the assumption that these inclusions are HSMI).

Entropy inequalities can be used to investigate features of quantum systems. For instance on the physical ground the strong subadditive property of the entropy together with the Lorentz covariance leads to a c -theorem for the entanglement entropy in $1+1$ dimensions and connections with the a -theorem are claimed in [CTT17a]. Due to the direct integral disintegration of the one-particle space (2), it is possible to generalize the Buchholz-Mach-Todorov endomorphism β_k [BMT88] to the direct integral of the $U(1)$ -current with $k \in C_0^{\infty}(X_-^0)$. Using the formula contained in [Lon20], we are able to compute the relative entropy with respect the $\omega \circ \beta_k$ and ω and deduce the QNEC – in this case to be intended $S''(t) > 0$ where S is the relative entropy related of the algebra $\mathcal{A}(N_{C+tA})$ with respect to $\omega \circ \beta_k$ and ω and the derivative is with respect to t . We also have a saturation of the strong superadditivity condition of the relative entropy considered.

This paper is organized as follows. In Section 2 we collect the basic notions such as one-particle space in terms of standard subspaces and its second quantization. In Section 3 we discuss observables on the null plane and their transversal decomposition. In Section 4 we obtain the decomposition of the modular operator of null plane regions and prove that inclusions of null plane regions are HSMI. In Section 5, after recalling the notions concerning relative entropy, ANEC, QNEC and some background also from physics, we study

the relative entropy and its relation with the energy inequalities and the saturation of the strong superadditivity condition of the null cut algebras between the BMT type states. In Section 6 we present concluding remarks, including the 1+1 dimensional case.

2 Preliminaries

In this Section we will recall the operator-algebraic formulations of the free field. A free field is constructed from its one-particle structure. A quantum and relativistic particle on Minkowski spacetime is a unitary positive energy representation of the Poincaré group. The localization property of the one-particle states is formulated in terms of the standard subspaces, and it translates to the localization property of the associated free fields through the second quantization. We will further comment on the $U(1)$ -current model, which will be a convenient tool to describe the restriction of the free theory on the null plane.

2.1 Abstract one-particle structure

A real linear, closed subspace H of a complex Hilbert space \mathcal{H} is called **cyclic** if $H + iH$ is dense in \mathcal{H} and **separating** if $H \cap iH = \{0\}$. A **standard subspace** is a real linear, closed subspace that is both cyclic and separating. We recall below some useful properties of standard subspaces, see [Lon08] for details.

It is possible to consider an analogue of the Tomita-Takesaki modular theory for standard subspaces. For a standard subspace H , the Tomita operator S_H is defined to be the closed anti-linear involution with dense domain $H + iH$ acting in the following way:

$$\begin{aligned} S_H : H + iH &\rightarrow H + iH \\ \xi + i\eta &\mapsto \xi - i\eta. \end{aligned}$$

The polar decomposition

$$S_H = \Delta_H^{\frac{1}{2}} J_H$$

defines the modular operator Δ_H and the modular conjugation J_H , and they satisfy the following relations:

$$J_H \Delta_H J_H = \Delta_H^{-1}, \quad \Delta_H^{it} H = H \quad \text{for } t \in \mathbb{R}, \quad J_H H = H',$$

where H' is the symplectic complement of H :

$$H' := \{\xi \in \mathcal{H} : \operatorname{Im} \langle \xi, \eta \rangle = 0 \text{ for } \eta \in H\}.$$

The symplectic complement H' is a standard subspace if and only if so is H , and a standard subspace H satisfies $H = H''$. The Tomita operator of the symplectic complement H' is given by

$$S_{H'} = S_H^* = \Delta_H^{-\frac{1}{2}} J_H = J_H \Delta_H^{\frac{1}{2}}.$$

The assignment $H \mapsto S_H$ of a closed, anti-linear, densely defined involution is one-to-one: For such an operator S , the real closed subspace $\ker(1 - S)$ is a standard subspace.

One can easily deduce the covariance of standard subspaces (see [Mor18, Lemma 2.2]):

Lemma 2.1. *Let $H \subset \mathcal{H}$ be a standard subspace and U be a unitary or anti-unitary operator on \mathcal{H} . Then UH is standard and $U\Delta_H U^* = \Delta_{UH}^{\epsilon(U)}$ and $UJ_H U^* = J_{UH}$ where $\epsilon(U) = 1$ if U is unitary and $\epsilon(U) = -1$ if U is anti-unitary.*

The following is an analogue of Borchers theorem [Bor92, Flo98] for standard subspaces, see [Lon08, Theorem 3.15].

Theorem 2.2. *Let H be a standard subspace of a Hilbert space \mathcal{H} and T a one-parameter group with positive generator such that $T(s)H \subset H$, $s \geq 0$, then the following hold:*

$$\begin{aligned}\Delta_H^{it} T(s) \Delta_H^{-it} &= T(e^{-2\pi t} s) \\ J_H T(s) J_H &= T(-s).\end{aligned}$$

We say that a pair of standard subspaces $K \subset H$ is a **half-sided modular inclusion (HSMI)** if

$$\Delta_H^{-it} K \subset K \quad \text{for } t \geq 0.$$

Let \mathbf{P} be the translation-dilation group, that is, the group of affine transformations of \mathbb{R} , where dilations act by $\mathfrak{d}(2\pi t)x = e^{2\pi t}x$, $x \in \mathbb{R}$ and translations act by $\mathfrak{t}(s)x = x + s$. The group \mathbf{P} contains also dilations centered at 1: $\mathfrak{d}_1(2\pi t)x = e^{2\pi t}(x - 1) + 1$. A HSMI of von Neumann algebras implies the existence of a one-parameter group of unitaries with certain properties [Wie93, AZ05]. The following is its standard subspace version and the first part can be found in [Lon08, Theorem 3.21].

Theorem 2.3. *Let $K \subset H$ be a half-sided modular inclusion of standard subspaces of the Hilbert space \mathcal{H} , then there exists a positive energy representation of the translation-dilation group \mathbf{P} given by*

$$U(\mathfrak{d}(2\pi t)) = \Delta_H^{-it}, \quad U(\mathfrak{d}_1(2\pi t)) = \Delta_K^{-it}$$

In particular, the translations are given by $U(\mathfrak{t}(e^{2\pi t} - 1)) = \Delta_H^{-it} \Delta_K^{it}$, satisfy $U(\mathfrak{t}(s))H \subset H$ for $s \geq 0$, $U(\mathfrak{t}(1))H = K$ and have a positive generator.

Furthermore, the generator P of the translation group is $\frac{1}{2\pi}(\log(\Delta_K) - \log(\Delta_H))$. In general, we have the relation $\log(\Delta_{U(\mathfrak{t}(s))H}) = \text{Ad } U(\mathfrak{t}(s))(\log(\Delta_H)) = \log(\Delta_H) + 2\pi sP$.

Proof. We prove the last statement. The operator $\log(\Delta_K) - \log(\Delta_H)$ is essentially self-adjoint on its natural domain (one can prove this by first taking the Gårding domain). Thus, we can apply Trotter's product formula:

$$\begin{aligned}e^{it(\log(\Delta_H) - \log(\Delta_K))} &= \text{s-lim}_{n \rightarrow \infty} \left(\Delta_H^{i\frac{t}{n}} \Delta_K^{-i\frac{t}{n}} \right)^n = \text{s-lim}_{n \rightarrow \infty} \left(U(\mathfrak{t}(e^{-2\pi\frac{t}{n}} - 1)) \right)^n \\ &= \text{s-lim}_{n \rightarrow \infty} U \left(\mathfrak{t}(n(e^{-2\pi\frac{t}{n}} - 1)) \right) = U(\mathfrak{t}(2\pi t)).\end{aligned}$$

The last relation follows from Lemma 2.1. □

2.2 The one-particle structure of the free scalar field

The relativistic invariance is encoded in the Poincaré group \mathcal{P} on the $(D + 1)$ -dimensional Minkowski space time \mathbb{R}^{D+1} , where $D > 1$. It is the semi-direct product of the full Lorentz group \mathcal{L} and the translation group \mathbb{R}^{D+1} :

$$\mathcal{P} = \mathcal{L} \ltimes \mathbb{R}^{D+1}.$$

The subgroup $\mathcal{P}_+^\uparrow = \mathcal{L}_+^\uparrow \ltimes \mathbb{R}^{D+1}$ of time- and space orientation-preserving transformations gives the relativistic transformations from one inertial frame to another. The causal structure is determined by the Minkowski metric and the causal complement of a region $O \subset \mathbb{R}^{1+D}$ is determined as follows:

$$O' = \{x \in \mathbb{R}^{1+D} : (y - x)^2 < 0, y \in O\}^\circ,$$

where \circ denotes the open kernel.

We restrict the consideration to the scalar representations of \mathcal{P}_+^\uparrow (see e.g. [Var85]). A **scalar representation with mass $m \geq 0$ (for $D > 1$)** is defined on the Hilbert space $\mathfrak{H}_m = L^2(\Omega_m, d\Omega_m)$, where $d\Omega_m$ is the unique (up to a constant) Lorentz-invariant measure on the mass shell $\Omega_m = \{p = (p_0, \dots, p_D) \in \mathbb{R}^{D+1} : p^2 = m^2, p_0 \geq 0\}$:

$$d\Omega_m = \theta(p_0) \delta(p^2 - m^2) d^{D+1}p.$$

Let $\omega_m(\mathbf{p}) = \sqrt{m^2 + |\mathbf{p}|^2}$, then $\Omega_m = \{(\omega_m(\mathbf{p}), \mathbf{p}) : \mathbf{p} \in \mathbb{R}^D\}$ and the measure can be expressed in the \mathbf{p} -coordinates as:

$$d\Omega_m = \frac{d^D \mathbf{p}}{\omega_m(\mathbf{p})}. \quad (3)$$

The action U_m of $(\Lambda, a) \in \mathcal{P}_+^\uparrow = \mathcal{L}_+^\uparrow \ltimes \mathbb{R}^{D+1}$ is given as follows:

$$(U_m((\Lambda, a))\Psi)(p) = e^{ia \cdot p} \Psi(\Lambda^{-1}p), \text{ for } \Psi \in \mathfrak{H}_m. \quad (4)$$

Consider the restriction E of the Fourier transformation on Schwartz functions on \mathbb{R}^{D+1} to Ω_m :

$$E : \mathcal{S}(\mathbb{R}^{D+1}) \rightarrow \mathcal{S}(\Omega_m)$$

$$f \mapsto (Ef)(p) = \int_{\mathbb{R}^{D+1}} e^{ix \cdot p} f(x) d^{D+1}x, \quad p \in \Omega_m$$

(as $D > 1$, this holds even if $m = 0$). Then $E(\mathcal{S}(\mathbb{R}^{D+1}))$ is dense in \mathfrak{H}_m . We refer to p as the momentum variable and x as the position variable. We shall denote by \mathbf{x}_\perp the $D - 1$ coordinate vector (x_2, \dots, x_D) . The action of the Poincaré group on the one-particle space (4) is covariant with respect to the action of \mathcal{P}_+^\uparrow on test functions in $\mathcal{S}(\mathbb{R}^{D+1})$:

$$(U_m((\Lambda, a))Ef)(x) = E(f(\Lambda^{-1}(x - a))).$$

The local structure of the free scalar field is encoded in the local space corresponding to bounded open regions $O \in \mathbb{R}^{D+1}$:

$$H(O) := \overline{\{Ef \in L^2(\Omega_m, d\Omega_m) : f \in \mathcal{S}(\mathbb{R}^{D+1}, \mathbb{R}), \text{supp}(f) \subset O\}}. \quad (5)$$

For an arbitrary regions S with non-empty interior in Minkowski space, its local subspace is generated by the subspaces of bounded open regions contained in it:

$$H(S) = \overline{\bigcup_{O \in S} H(O)}.$$

The map $\mathbb{R}^{D+1} \supset O \mapsto H(O) \subset \mathcal{H}$ and the Poincaré representation U_m define a net of standard subspaces, also called one-particle net or first quantized net, satisfying the following properties (see e.g. [BGL02]):

- (SS1) **Isotony:** $H(O_1) \subset H(O_2)$ for $O_1 \subset O_2$;
- (SS2) **Locality:** if $O_1 \subset O'_2$, then $H(O_1) \subset H(O'_2)$, where O' denotes the spacelike complement of O ;
- (SS3) **Poincaré covariance:** $H(gO) = U(g)H(O)$ for $g \in \mathcal{P}_+^\uparrow$;
- (SS4) **Spectral condition:** the joint spectrum of the translation subgroup in U is contained in the closed forward light cone $\overline{V_+} = \{p \in \mathbb{R}^{D+1} : p^2 \geq 0, p_0 \geq 0\}$.
- (SS5) **Cyclicity:** $H(O)$ are cyclic subspaces.

It is a consequence of locality and cyclicity that $H(O)$ are standard subspaces.

Consider the standard wedge $W_1 = \{x \in \mathbb{R}^{D+1} : |x_0| < x_1\}$ in the x_1 -direction and let

$$\Lambda_{W_1}(t)(x_0, x_1, \mathbf{x}_\perp) = (\cosh(t)x_0 + \sinh(t)x_1, \sinh(t)x_0 + \cosh(t)x_1, \mathbf{x}_\perp)$$

be the one-parameter group of Lorentz boosts fixing W_1 . Any other region of the form $W = gW_1, g \in \mathcal{P}_+^\uparrow$ is also called a wedge, and we put $\Lambda_W(t) = g\Lambda_{W_1}(t)g^{-1}$ with such a g (this is well-defined, because any other g differs only by a Lorentz boost preserving the wedge). We shall denote by \mathcal{W} the set of wedges. Let $H(W) = \overline{\bigcup_{O \in W} H(O)}$ be the subspace associated to the wedge W . The net defined by (5) further satisfies the following properties.

- (SS6) **Bisognano–Wichmann (BW) property:** Let $W \in \mathcal{W}$, it holds that

$$U(\Lambda_W(t)) = \Delta_{H(W)}^{-\frac{it}{2\pi}} \quad \text{for } t \in \mathbb{R},$$

- (SS7) **Haag duality (for wedges):** $H(W') = H(W)'$, for all $W \in \mathcal{W}$.

2.3 Second quantization and nets of von Neumann algebras

Let \mathcal{H} be a Hilbert space and $H \subset \mathcal{H}$ a real linear subspace. The von Neumann algebra $R(H)$, called second quantization algebras, on the symmetric Fock space $\mathcal{F}_+(\mathcal{H})$ generated by the Weyl operators:

$$R(H) \equiv \{w(\xi) : \xi \in H\}'' ,$$

where $w(\xi), \xi \in \mathcal{H}$ are unitary operators on $\mathcal{F}_+(\mathcal{H})$ characterized by

$$w(\xi)e^\eta = e^{-\frac{1}{2}\langle \xi, \xi \rangle - \langle \xi, \eta \rangle} \cdot e^{\xi + \eta},$$

where $e^\eta = 1 \oplus \eta \oplus \frac{1}{2!}\eta^{\otimes 2} \dots$ is a coherent vector in $\mathcal{F}_+(\mathcal{H})$. By strong continuity of the map $\mathcal{H} \ni \xi \mapsto w(\xi) \in \mathcal{F}_+(\mathcal{H})$ we have that

$$R(H) = R(\overline{H}) .$$

Moreover, the Fock vacuum vector $\Omega = e^0$ is cyclic (respectively separating) for $R(H)$ if and only if \overline{H} is cyclic (respectively separating). Therefore one verifies the relation:

$$\langle w(\xi)\Omega, w(\eta)\Omega \rangle = e^{-\frac{1}{2}(\|\xi\|^2 + \|\eta\|^2)} e^{\langle \xi, \eta \rangle} . \quad (6)$$

The second quantization respects the lattice structure [Ara63] and the modular structure [LRT78, LMR16]. We shall denote $J_{R(H), \Omega}$, $\Delta_{R(H), \Omega}$ the Tomita operators associated with $(R_+(H), \Omega)$, and by $\Gamma_+(T)$ the multiplicative second quantization of a one-particle operator T on \mathcal{H} . It holds that $\Gamma_+(T)e^\xi = e^{T\xi}$ for $\xi \in \mathcal{H}$.

Proposition 2.4. [LRT78, LMR16] *Let H and H_a be closed, real linear subspaces of \mathcal{H} . We have*

1. $J_{R(H), \Omega} = \Gamma_+(J_H)$, $\Delta_{R(H), \Omega} = \Gamma_+(\Delta_H)$ if H is standard;
2. $R(H)' = R(H')$;
3. $R(\text{Span}_a H_a) = \bigvee_a R(H_a)$;
4. $R(\bigcap_a H_a) = \bigcap_a R(H_a)$,

where $\bigvee_a R(H_a)$ denotes the von Neumann algebra generated by $R(H_a)$.

In particular, the second quantization promotes the one-particle net defined in (5) to a Haag-Kastler net of local algebras: consider the map $\mathbb{R}^{D+1} \supset O \mapsto \mathcal{A}(O) = R(H(O)) \subset \mathcal{F}_+(\mathcal{H})$ together with the second quantization $U = \Gamma_+(U_m)$ of the one-particle Poincaré representation U_m , then the following hold:

- (HK1) **Isotony:** $\mathcal{A}(O_1) \subset \mathcal{A}(O_2)$ for $O_1 \subset O_2$;
- (HK2) **Locality:** if $O_1 \subset O_2'$, then $\mathcal{A}(O_1) \subset \mathcal{A}(O_2)'$;
- (HK3) **Poincaré covariance:** $U(g)\mathcal{A}(O)U(g)^* = \mathcal{A}(gO)$ for $g \in \mathcal{P}_+^\uparrow$.

(HK4) **Positivity of the energy:** the joint spectrum of the translation subgroup in U is contained in the closed forward light cone $\overline{V_+}$.

(HK5) **Vacuum and the Reeh-Schlieder property:** Ω is the (up to a phase) unique vector such that $U(g)\Omega = \Omega$ for $g \in \mathcal{P}_+^\uparrow$ and is cyclic, $\overline{\mathcal{A}(O)\Omega} = \mathcal{H}$ for any O .

(HK6) **The Bisognano-Wichmann property:** For a wedge $W \in \mathcal{W}$, we put $\mathcal{A}(W) = (\bigvee_{O \subset W} \mathcal{A}(O))''$. Then it holds that

$$U(\Lambda_W(t)) = \Delta_{\mathcal{A}(W), \Omega}^{-\frac{it}{2\pi}},$$

where $\Delta_{\mathcal{A}(W), \Omega}^{it}$ is the modular group of $\mathcal{A}(W)$ with respect to Ω .

2.4 The $U(1)$ -current net

We introduce a family of standard subspaces parametrized by intervals on \mathbb{R} , the $U(1)$ -current. Let us consider $\mathcal{H}_{U(1)} = L^2(\mathbb{R}, d\theta')$ and, for each $I \subset \mathbb{R}$ connected open interval, the subspace

$$H_{U(1)}(I) = \overline{\{\hat{g}(e^{-\theta'}); g \in C_0^\infty(\mathbb{R}, \mathbb{R}), \hat{g}(0) = 0, \text{supp}(g) \subset I\}} \subset L^2(\mathbb{R}, d\theta'),$$

where \hat{g} is the Fourier transform of g . Furthermore, we introduce

$$(U_{U(1)}(\alpha, t)\xi)(\theta') = e^{ite^{-\theta'}} \xi(\theta' - \alpha).$$

This is a unitary representation of the translation-dilation group $\mathbf{P} = \mathbb{R} \ltimes \mathbb{R}$. Let $x \in \mathbb{R}$, then $(\alpha, 0) \in \mathbb{R} \ltimes \mathbb{R}$ corresponds to the dilation $\mathfrak{d}(\alpha)x = e^\alpha x$ and $(0, t) \in \mathbb{R} \ltimes \mathbb{R}$ corresponds to the translations $\mathfrak{t}(t)x = x + t$. It is straightforward to check that $U_{U(1)}(\alpha, t)H(I) = H(e^\alpha I + t)$.

Each $H(I)$ is a standard subspace and for disjoint I_1, I_2 it holds that $H(I_1) \subset H(I_2)'$. Furthermore, the Bisognano-Wichmann property holds: $\Delta_{H(\mathbb{R}_+)}^{it} = U_{U(1)}(-2\pi t, 0)$, $t \in \mathbb{R}$.

Let $f, g \in C_0^\infty(\mathbb{R}, \mathbb{R})$ with $\hat{f}(0) = \hat{g}(0) = 0$ and call the primitives F, G , respectively. The condition $\hat{f}(0) = \hat{g}(0) = 0$ implies $F, G \in C_0^\infty(\mathbb{R}, \mathbb{R})$. We have under the substitution $p = e^{-\theta'}$:

$$\langle \hat{g}(e^{-\theta'}), \hat{f}(e^{-\theta'}) \rangle = \int_{\mathbb{R}_+} \hat{g}(-p) \hat{f}(p) \frac{dp}{p} = \int_{\mathbb{R}_+} \hat{G}(-p) \hat{F}(p) p dp.$$

The imaginary part of the scalar product (symplectic form) plays a crucial role in the second quantization

$$\text{Im} \langle \hat{g}(e^{-\theta'}), \hat{f}(e^{-\theta'}) \rangle = \frac{1}{2i} \int_{\mathbb{R}_+} \left(\hat{G}(-p) \hat{F}(p) - \hat{F}(-p) \hat{G}(p) \right) p dp = \frac{1}{2} \int_{\mathbb{R}} G(x) f(x) dx \quad (7)$$

The family $\{H(I)\}$ and the representation $U_{U(1)}$ are unitarily equivalent to the family of closed real subspaces coming from the $U(1)$ -current conformal net and the one-particle symmetry restricted to the translation-dilation group, see [BT15, Section 5.2]. The intertwining map for $h \in C_0^\infty(\mathbb{R}, \mathbb{R})$ is $\hat{h}(p) \mapsto \hat{h}'(e^{-\theta'})$. Thus to switch to the standard definition of the $U(1)$ -current presented in the literature (see e.g. [Lon08]) one replaces $g \in C_0^\infty(\mathbb{R}, \mathbb{R})$ with $\hat{g}(0) = 0$ with its primitive $G \in C_0^\infty(\mathbb{R}, \mathbb{R})$.

3 Free scalar field on the null plane

3.1 Direct integrals and decompositions

Let us summarize some of the basic notions and results on the direct integral of Hilbert spaces and decomposition of group representations. We follow the conventions of [Dix81], see also [KR97].

Let X be a σ -compact locally compact Borel measure space, ν the completion of a Borel measure on X and $\{\mathcal{K}_\lambda\}$ a family of separable Hilbert spaces indexed by $\lambda \in X$. We say that a separable Hilbert space \mathcal{K} is the **direct integral** of $\{\mathcal{K}_\lambda\}$ over (X, ν) if, to each $\xi \in \mathcal{K}$ there corresponds a function $\lambda \mapsto \xi(\lambda) \in \mathcal{K}_\lambda$ and

1. $\lambda \mapsto \langle \xi(\lambda), \eta(\lambda) \rangle$ is ν -integrable and $\langle \xi, \eta \rangle = \int_X \langle \xi(\lambda), \eta(\lambda) \rangle d\nu(\lambda)$
2. if $\phi_\lambda \in \mathcal{K}_\lambda$ for all λ and $\lambda \mapsto \langle \phi_\lambda, \eta(\lambda) \rangle$ is integrable for all $\eta \in \mathcal{K}$, there is $\phi \in \mathcal{K}$ such that $\phi(\lambda) = \phi_\lambda$ for almost every λ .

In this case, we write:

$$\mathcal{K} = \int_X^\oplus \mathcal{K}_\lambda d\nu(\lambda).$$

An operator $T \in \mathcal{B}(\mathcal{K})$ is said to be **decomposable** when there is a function $\lambda \mapsto T_\lambda$ on X such that $T_\lambda \in \mathcal{B}(\mathcal{K}_\lambda)$ and, for each $\xi \in \mathcal{K}$, $T_\lambda \xi(\lambda) = (T\xi)(\lambda)$ for ν -almost every λ , and in this case we write $T = \int_X^\oplus T_\lambda d\nu(\lambda)$. If $T_\lambda = f(\lambda)1$ for some $f \in L^\infty(X, \nu)$, we say that T is **diagonalizable**. An operator $T \in \mathcal{B}(\mathcal{K})$ is decomposable if and only if T commutes with every diagonalizable operator. Conversely, let $X \ni \lambda \mapsto T_\lambda \in \mathcal{B}(\mathcal{K}_\lambda)$ be a field of bounded operators with $\sup_\lambda \|T_\lambda\| < \infty$. If for any $\xi \in \mathcal{K}$ with $\xi = \int_X^\oplus \xi(\lambda) d\nu(\lambda)$ there exists $\eta \in \mathcal{K}$ such that $\eta(\lambda) = T_\lambda \xi(\lambda)$ for almost every $\lambda \in X$, then $T = \int_X^\oplus T_\lambda d\nu(\lambda) \in \mathcal{B}(\mathcal{K})$ defines a bounded decomposable operator on \mathcal{K} and $\lambda \mapsto T_\lambda$ is called a *measurable field of bounded operators*.

The following Lemma can be found in [MT18, Appendix B].

Lemma 3.1. *Let $\mathcal{K} = \int_X^\oplus \mathcal{K}_\lambda d\nu(\lambda)$ then*

- (a) *Let $H = \int_X^{\oplus \mathbb{R}} H_\lambda d\nu(\lambda) \subset \mathcal{K}$ be a real subspace and the direct integral is taken on \mathbb{R} such that $H_\lambda \subset \mathcal{K}_\lambda$, then $H' = \int_X^{\oplus \mathbb{R}} H'_\lambda d\nu(\lambda)$.*
- (b) *Let $\{H_k\}_{k \in \mathbb{N}}$ be a countable family of real subspaces of \mathcal{K} such that $H_k = \int_X^{\oplus \mathbb{R}} (H_k)_\lambda d\nu(\lambda)$ on \mathbb{R} and $(H_k)_\lambda \subset \mathcal{K}_\lambda$, then $\bigcap_{k \in \mathbb{N}} H_k = \int_X^{\oplus \mathbb{R}} \bigcap_{k \in \mathbb{N}} (H_k)_\lambda d\nu(\lambda)$.*
- (c) *Let $\{H_k\}_{k \in \mathbb{N}}$ be a countable family of real subspaces of \mathcal{K} such that $H_k = \int_X^{\oplus \mathbb{R}} (H_k)_\lambda d\nu(\lambda)$ on \mathbb{R} and $(H_k)_\lambda \subset \mathcal{K}_\lambda$, then $\overline{\text{Span}_{k \in \mathbb{N}} H_k} = \int_X^{\oplus \mathbb{R}} \overline{\text{Span}_{k \in \mathbb{N}} (H_k)_\lambda} d\nu(\lambda)$.*

Let G be a locally compact group and π a continuous unitary representation of G on $\mathcal{K} = \int_X^\oplus \mathcal{K}_\lambda d\nu(\lambda)$. Suppose that, for each $g \in G$, we have $\pi(g) = \int_X^\oplus \pi_\lambda(g) d\mu(\lambda)$, then we say that π is the **direct integral** of the π_λ and write:

$$\pi = \int_X^\oplus \pi_\lambda d\nu(\lambda).$$

Equivalently, π is a direct integral if each $\pi(g)$, $g \in G$, is decomposable. It follows immediately that the direct integral of standard subspaces is again a standard subspaces on the direct integral of Hilbert space, and so is a direct integral of Poincaré covariant nets of standard subspaces.

As a particular case, let $\mathcal{K} = \int_X^\oplus \mathcal{K}_0 d\nu$ be a direct integral Hilbert space over the constant field \mathcal{K}_0 , which is a separable Hilbert space, on X with measure ν . Then it holds that [Dix81, Proposition II.1.8.11, Corollary]

$$\int_X^\oplus \mathcal{K}_0 d\nu(\lambda) \simeq L^2(X, \nu) \otimes \mathcal{K}_0. \quad (8)$$

The isomorphism is given by identifying $L^2(X, \nu) \otimes \mathcal{K}_0$ as the space of \mathcal{K}_0 -valued L^2 -functions. By this isomorphism, we identify $\int_X^\oplus T d\nu(\lambda)$ and $\mathbb{1} \otimes T$, where $T \in \mathcal{B}(\mathcal{K}_0)$ (a constant field of bounded operators).

3.2 Decomposition of the one-particle space

Let us fix $D > 1$. We first consider the $m > 0$ case. In Remark 3.5 we explain the minor modification to deal with the massless case.

The hypersurface $X_-^0 := \{x \in \mathbb{R}^{D+1} : x_0 - x_1 = 0\}$ is called the **null plane** in the x_1 -direction. It is appropriate to consider the coordinate frame $(x_+, x_-, \mathbf{x}_\perp) = (\frac{x_0+x_1}{\sqrt{2}}, \frac{x_0-x_1}{\sqrt{2}}, \mathbf{x}_\perp)$. In these coordinates, the Minkowski product becomes

$$x \cdot p = x_+ p_- + x_- p_+ - \mathbf{x}_\perp \mathbf{p}_\perp \quad (9)$$

and the Minkowski (pseudo)norm $x^2 = 2x_+ x_- - \mathbf{x}_\perp^2$. In the momentum space, the massive hyperboloid is determined by $2p_+ p_- - \mathbf{p}_\perp^2 = m^2$, and for each $(p_-, \mathbf{p}_\perp) \in \mathbb{R}_+ \times \mathbb{R}^{D-1}$ there is one and only one $p_+ \in \mathbb{R}_+$ satisfying this equation. For a test function $f = f(x_+, x_-, \mathbf{x}_\perp)$ on \mathbb{R}^{D+1} , let Ef be its Fourier transform restricted to $\Omega_m = \left\{ p = \left(\frac{m^2 + \mathbf{p}_\perp^2}{2p_-}, p_-, \mathbf{p}_\perp \right) : p_- > 0, \mathbf{p}_\perp \in \mathbb{R}^{D-1} \right\}$ (in the $(p_+, p_-, \mathbf{p}_\perp)$ -coordinates). In the (p_-, \mathbf{p}_\perp) -coordinates, we have, up to a unitary (given by the change of variables),

$$\mathfrak{H}_m \simeq L^2 \left(\mathbb{R}^D, \frac{d^D \mathbf{p}}{\omega_m(\mathbf{p})} \right) \simeq L^2 \left(\mathbb{R}_+ \times \mathbb{R}^{D-1}, \frac{2p_- dp_- d\mathbf{p}_\perp}{m^2 + 2p_-^2 + \mathbf{p}_\perp^2} \right)$$

(since $dp_1 d\mathbf{p}_\perp = \frac{1}{\sqrt{2}} dp_- d\mathbf{p}_\perp$ and $\omega_m(p_-, \mathbf{p}_\perp) = \frac{1}{\sqrt{2}} \left(\frac{m^2 + \mathbf{p}_\perp^2}{2p_-} + p_- \right)$ is the dispersion relation in the (p_-, \mathbf{p}_\perp) -coordinates). For sake of notational simplicity, we will write $\xi(p_1, \mathbf{p}_\perp) \in \mathfrak{H}_m$ or $\xi(p_-, \mathbf{p}_\perp) \in \mathfrak{H}_m$ for $\xi \in \mathfrak{H}_m$ in terms of (p_1, \mathbf{p}_\perp) or (p_-, \mathbf{p}_\perp) , respectively. In the same way, the representation U_m acts on \mathfrak{H}_m in various realizations.

We introduce the map $V_M : \mathfrak{H}_m = L^2(\Omega_m, d\Omega_m) \rightarrow L^2(\mathbb{R}^D, d\theta' d^{D-1} \mathbf{p}_\perp)$ as follows:

$$(V_M^{-1} \xi)(\theta', \mathbf{p}_\perp) = \xi(\operatorname{arcsinh}(\frac{p_1}{\omega_m(\mathbf{p}_\perp)}) - \ln(\omega_m(\mathbf{p}_\perp)) + \ln \sqrt{2}, \mathbf{p}_\perp), \quad \xi \in L^2(\mathbb{R}^D, d\theta' d^{D-1} \mathbf{p}_\perp),$$

where $\omega_m(\mathbf{p}_\perp) := \sqrt{m^2 + \mathbf{p}_\perp^2}$. V_M is a unitary operator. Indeed, first, the change of p_1 to rapidity $\theta = \operatorname{arcsinh}(\frac{p_1}{\omega_m(\mathbf{p}_\perp)})$ for a fixed \mathbf{p}_\perp , or $p_1 = \omega_m(\mathbf{p}_\perp) \sinh \theta$ is a smooth one-to-one map

$\mathbb{R}^D \rightarrow \mathbb{R}^D$. Moreover, with $\frac{\partial \theta}{\partial p_1} = \frac{1}{\omega_m(\mathbf{p}_\perp)} / \sqrt{1 + \frac{p_1^2}{\omega_m(\mathbf{p}_\perp)^2}} = \frac{1}{\omega_m(\mathbf{p})}$, the measure $d\Omega_m$ transforms in the following way, cf. (3):

$$d\Omega_m = \frac{d^D \mathbf{p}}{\omega_m(\mathbf{p})} = d\theta d^{D-1} \mathbf{p}_\perp.$$

Therefore, the pullback of the rapidity substitution imposes the equivalence:

$$\mathfrak{H}_m = L^2(\Omega_m, d\Omega_m) \simeq L^2(\mathbb{R} \times \mathbb{R}^{D-1}, d\theta \times d^{D-1} \mathbf{p}_\perp) (\simeq L^2(\mathbb{R}, d\theta) \otimes L^2(\mathbb{R}^{D-1}, d^{D-1} \mathbf{p}_\perp)).$$

Moreover, the substitution $\theta' = \theta - \ln(\omega_m(\mathbf{p}_\perp)) + \ln \sqrt{2}$ is a smooth one-to-one map: $\mathbb{R} \rightarrow \mathbb{R}$ if $m > 0$ with $d\theta = d\theta'$.

By substituting $\theta = \operatorname{arcsinh}(\frac{p_1}{\omega_m(\mathbf{p}_\perp)})$ or $p_1 = \omega_m(\mathbf{p}_\perp) \sinh \theta$ in $p_- = \frac{1}{\sqrt{2}}(\omega_m(p_1, \mathbf{p}_\perp) - p_1)$ we have $p_- = \frac{1}{\sqrt{2}}\omega_m(\mathbf{p}_\perp)e^{-\theta}$ therefore,

$$p_- = \frac{1}{\sqrt{2}}\omega_m(\mathbf{p}_\perp)e^{-\theta' - \ln(\omega_m(\mathbf{p}_\perp)) + \ln \sqrt{2}} = e^{-\theta'},$$

and

$$(V_M \xi)(\theta', \mathbf{p}_\perp) = \xi(e^{-\theta'}, \mathbf{p}_\perp), \quad \xi \in L^2\left(\mathbb{R}_+ \times \mathbb{R}^{D-1}, \frac{2p_- dp_- d\mathbf{p}_\perp}{m^2 + 2p_-^2 + \mathbf{p}_\perp^2}\right)$$

Proposition 3.2. *We have the following equivalence, where the map is given by V_M composed by the inverse Fourier transform on the perpendicular momenta:*

$$\mathfrak{H}_m \simeq L^2(\mathbb{R}^D, d\theta' d^{D-1} \mathbf{x}_\perp) \simeq \int_{\mathbb{R}^{D-1}}^\oplus L^2(\mathbb{R}, d\theta') d^{D-1} \mathbf{x}_\perp, \quad (10)$$

where the last equivalence is given by (8).

Subsequently, we will refer to these direct integral representations of \mathfrak{H}_m as **spatial decomposition** and denote the intertwining unitary by V_S , while the map V_M is referred to as **momentum decomposition**.

Next we discuss the decomposition of the spacetime symmetry. Let us take the wedge W_1 in the x_1 -direction and consider the subgroup $\Lambda_1 \ltimes \mathfrak{t}_{x_+} \subset \mathcal{P}_+^\uparrow$ consisting of boosts $\Lambda_1 := \Lambda_{W_1}$ along the x_1 -direction and lightlike translations \mathfrak{t}_{x_+} of x_+ . It is straightforward that this group is isomorphic to the translation-dilation group \mathbf{P} of \mathbb{R} . We will show that the restriction of U_m to \mathbf{P} decomposes with respect to the decompositions (10) of the one-particle vectors.

Lemma 3.3. *The unitary V_M between \mathfrak{H}_m and $\int_{\mathbb{R}^{D-1}}^\oplus L^2(\mathbb{R}, d\theta') d^{D-1} \mathbf{p}_\perp$ intertwines the actions of the subgroup \mathbf{P} to:*

$$(V_M U_m(\Lambda_1(\alpha), \mathfrak{t}_{x_+}(a_+))\xi)(\theta', \mathbf{p}_\perp) = e^{ia_+ e^{-\theta'}} (V_M \xi)(\theta' - \alpha, \mathbf{p}_\perp), \quad (11)$$

Proof. We start with the Poincaré group action given by U_m on \mathfrak{H}_m expressed in (4). We recall that $p_- = e^{-\theta'}$. Then the result is immediate by considering (9) and the fact that $a_+ \cdot p_- = a_+ e^{-\theta'}$,

For boosts, the claim follows directly from the following

$$\begin{aligned}\Lambda_1^{-1}(\alpha)p &= \begin{pmatrix} \cosh(\alpha)p_0 - \sinh(\alpha)p_1 \\ -\sinh(\alpha)p_0 + \cosh(\alpha)p_1 \\ \mathbf{p}_\perp \end{pmatrix} \\ &= \begin{pmatrix} \omega_m(\mathbf{p}_\perp) \cosh(\theta' + \ln(\omega_m(\mathbf{p}_\perp)) - \ln \sqrt{2} - \alpha) \\ \omega_m(\mathbf{p}_\perp) \sinh(\theta' + \ln(\omega_m(\mathbf{p}_\perp)) - \ln \sqrt{2} - \alpha) \\ \mathbf{p}_\perp \end{pmatrix}.\end{aligned}$$

□

Proposition 3.4. *Under the spatial decomposition, the representation of U_m to \mathbf{P} is decomposable (in the sense of Section 3.1) and acts as follows:*

$$(U_m((\Lambda_1(\alpha), \mathbf{t}_{x_+}(a_+)))\xi)(\theta', \mathbf{x}_\perp) = e^{ia_+e^{-\theta'}} \xi(\theta' - \alpha, \mathbf{x}_\perp). \quad (12)$$

Proof. The representation U_m decomposes in the momentum decomposition as (11), and the action of U_m does not involve the variable \mathbf{p}_\perp , it commutes with the inverse Fourier transform on \mathbf{p}_\perp . □

Remark 3.5. The disintegration (10) and the identifications (11) and (12) apply also to the $m = 0$ case up to a measure zero set. To see this, note that in the massless case $\Omega_0 = \partial V_+ = \{p \in \mathbb{R}^{D+1} : p^2 = 0, p_0 > 0\}$. The change of variables determining V_M^{-1} (both in (p_1, \mathbf{p}_\perp) or (p_-, \mathbf{p}_\perp) set of coordinates) are 1-1 diffeomorphisms on the set $\Omega_0 \cap \{p \in \mathbb{R}^{D+1} : \mathbf{p}_\perp \neq 0\}$. This set is dense in Ω_0 with respect to the topology induced by the euclidean topology on \mathbb{R}^{D+1} . The complement $N_0 = \{p \in \Omega_0 : \mathbf{p}_\perp = 0\}$ has measure zero with respect to the Lorentz invariant measure on Ω_0 , see (3). We conclude that V_M is a unitary operator. As Λ_1 -boosts fix N_0 , (10), (11) and (12) continue to hold (up to a measure zero set). This is the only modification to have in mind along the paper to adapt the massless case.

3.3 The local subspace of the null plane

Here we show that the free field can be restricted to regions on the null plane if we exclude the lightlike zero-modes (see Lemma 3.6 for the precise restriction, cf. [Ull04]). Recall that the algebra $\mathcal{A}(O)$ is generated by the Weyl operators associated with test functions supported in O . In this spirit, we define the local subspaces of open (with the relative topology) regions R in the null plane $X_-^0 := \{x \in \mathbb{R}^{D+1}; x_- = 0\}$. To make it precise, we consider distributions of the form

$$g_0(x_+, x_-, \mathbf{x}_\perp) = \delta(x_-)g(x_+, \mathbf{x}_\perp), \quad (13)$$

where g is a real-valued function such that $\mathbb{R} \ni x_+ \mapsto g(x_+, \mathbf{x}_\perp)$ is a compactly supported smooth function on X_-^0 . We denote the set of such distributions by $\mathcal{D}(X_-^0, \mathbb{R})$, and call them **thin test functions** supported on X_-^0 .

The Fourier transform of such g_0 is a smooth function and can be restricted to the mass shell, which we denote by Eg_0 . Then it is natural to consider one-particle vectors associated with thin test functions.

Lemma 3.6. *Let $g_0(x_+, x_-, \mathbf{x}_\perp) = \delta(x_-)g(x_+, \mathbf{x}_\perp)$ as in (13). Then $Eg_0 \in L^2(\Omega_m, d\Omega_m)$ (the norm is finite) if and only if $\int dx_+ g(x_+, \mathbf{x}_\perp) = 0$ for each \mathbf{x}_\perp .*

Proof. We take $g_0(x_+, x_-, \mathbf{x}_\perp) = \delta(x_-)g(x_+, \mathbf{x}_\perp)$ as above. The Fourier transform of g_0 is

$$(Eg_0)(p) = \int_{\mathbb{R}^{D+1}} e^{i(x_+p_- + x_-p_+ - \mathbf{x}_\perp \cdot \mathbf{p}_\perp)} \delta(x_-)g(x_+, \mathbf{x}_\perp) dx_+ dx_- d^{D-1}\mathbf{x}_\perp = \hat{g}(p_-, \mathbf{p}_\perp).$$

By rapidity substitution $p_- = \frac{1}{\sqrt{2}}\omega_m(\mathbf{p}_\perp)e^{-\theta}$ and $p_+ = \frac{1}{\sqrt{2}}\omega_m(\mathbf{p}_\perp)e^\theta$ we have:

$$\|Eg_0\|_{\mathfrak{H}_m^-}^2 = \int_{\mathbb{R} \times \mathbb{R}^{D-1}} \left| \hat{g}\left(\frac{1}{\sqrt{2}}\omega_m(\mathbf{p}_\perp)e^{-\theta}, \mathbf{p}_\perp\right) \right|^2 d\theta d^{D-1}\mathbf{p}_\perp = \int_{\mathbb{R} \times \mathbb{R}^{D-1}} |\hat{g}(e^{-\theta'}, \mathbf{p}_\perp)|^2 d\theta' d^{D-1}\mathbf{p}_\perp$$

as we did in (10). Therefore, it is necessary that $\hat{g}(0, \mathbf{p}_\perp) = 0$ for each \mathbf{p}_\perp in order to have $Eg_0 \in \mathfrak{H}_m$ (that is the integral is finite), and equivalently, $\int dx_+ g(x_+, \mathbf{x}_\perp) = 0$ for each \mathbf{x}_\perp .

Conversely, if $\int dx_+ g(x_+, \mathbf{x}_\perp) = 0$ for each \mathbf{x}_\perp , then $\hat{g}(0, \mathbf{p}_\perp) = 0$. In particular, since g is smooth by the Taylor formula $\theta' \mapsto g(e^{-\theta'}, \mathbf{p}_\perp)$ has finite L^2 -norm at $+\infty$. Then since g is compactly supported, then it decays fast when $p_- \rightarrow -\infty$ and $|\mathbf{p}_\perp| \rightarrow +\infty$. As a consequence $Eg_0 \in \mathfrak{H}_m$. \square

The condition $\int dx_+ g(x_+, \mathbf{x}_\perp) = 0$ for each \mathbf{x}_\perp is satisfied if g is the partial derivative in x_+ of another compactly supported smooth function. We define, for an open bounded region R on the null plane X_-^0 , as follows:

$$H(R) := \overline{\text{Span}}^{\mathfrak{H}_m} \{Eg_0 \in L^2(\Omega_m, d\Omega_m) : g_0 = \delta(x_-)\partial_{x_+}g(x_+, \mathbf{x}_\perp) \in \mathcal{D}(X_-^0, \mathbb{R}), \text{supp } g \subset R\}.$$

For a general open region $\tilde{R} \subset X_-^0$, the subspace is generated by the subspaces of open, bounded regions inside \tilde{R} :

$$H(\tilde{R}) = \overline{\bigcup_{R \in \tilde{R}} H(R)}.$$

In particular, a **null cut** is a region on the null plane associated with a continuous function $C : \mathbb{R}^{D-1} \rightarrow \mathbb{R}$, where \mathbb{R}^{D-1} denotes the subspace $\{\mathbf{x} \in \mathbb{R}^{D+1}; x_- = x_+ = 0\}$

$$N_C := \{\mathbf{x} = (x_+, x_-, \mathbf{x}_\perp) \in \mathbb{R}^{D+1}; x_- = 0, x_+ > C(\mathbf{x}_\perp)\}.$$

These subspaces are natural in the view of the following isotony in an extended sense.

Lemma 3.7. *Let $R \subset X_-^0$ be open (relatively in X_-^0) and $O \subset \mathbb{R}^{D+1}$ open such that there is a direction $a \in \mathbb{R}^{D+1}$ where $R \subset O + ta$ with $0 < t < \epsilon$ for sufficiently small ϵ . Then it holds that $H(R) \subset H(O)$.*

Proof. Let g be any smooth function compactly supported in R . Then for g_- a test function on R such that $g_-(t) \geq 0$ and $\int g_-(t)dt = 1$, $g(x_-, x_+, \mathbf{x}_\perp) = g_-(x_-)\partial_{x_+}g(x_+, \mathbf{x}_\perp)$ has a Fourier transform in $\mathfrak{H}_m = L^2(\Omega_m, d\Omega_m)$. Indeed, we know that the Fourier transform of $\partial_{x_+}g(x_+, \mathbf{x}_\perp)$ alone is in \mathfrak{H}_m , and it gets multiplied by $\hat{g}_-(p_+)$, which is rapidly decreasing. If




Figure 1: The geometric setup and illustration of the decomposition of standard subspaces of null cuts.

we consider the scaled function $g_{-,n}(x_-) = ng_-(nx_-)$, its Fourier transform is $\hat{g}_-(\frac{p_+}{n})$, which is bounded and converges to 1, therefore, $E(g_{-,n}g)$ converges to Eg_0 .

On the other hand, for fixed a and t as in the statement, for sufficiently large n , $g_{-,n}g$ is supported in $O + ta$, hence we have $Eg_0 \subset H(O + ta)$ by the closedness of $H(O + ta)$. This implies $H(R) \subset H(O + ta) = U_m(ta, 0)H(O)$, and by the continuity of U_m , we obtain $H(R) \subset H(O)$. \square

The spaces $H(R)$ are real subspaces of \mathfrak{H}_m . Recalling the spatial decomposition (10) of \mathfrak{H}_m , we will show that the same decomposition holds as a real Hilbert space through V_S .

Let $I \subset \mathbb{R}$ a bounded open interval and $I_\perp \subset \mathbb{R}^{D-1}$ a bounded open region. Under the spatial decomposition (10), we consider the local subspace $H(R)$ of rectangular regions as follows

$$R = \{(x_+, x_-, \mathbf{x}_\perp) : x_- = 0, x_+ \in I, \mathbf{x}_\perp \in I_\perp\}.$$

Proposition 3.8. *Let R be a rectangle on the null plane X_-^0 . Under the spatial decomposition (10), the local subspace $H(R)$ decomposes in the following way (see e.g. [MT18, Proposition B.2]):*

$$\begin{aligned} H(R) &\rightarrow \int_{I_\perp}^{\oplus \mathbb{R}} H_{U(1)}(I) d^{D-1} \mathbf{x}_\perp \\ Eg_0 &\mapsto V_S Eg_0, \end{aligned} \tag{14}$$

where $g_0(x_-, x_+, \mathbf{x}_\perp) = \delta(x_-)g(x_+, \mathbf{x}_\perp)$ as in (13) satisfying the condition of Lemma 3.6.

$H(R)$ is separating for any $R = I \times I_\perp$ where I is proper, and standard if I contains an open interval and $I_\perp = \mathbb{R}^{D-1}$.

Proof. By the spatial decomposition (10), we identify the whole Hilbert space \mathfrak{H}_m with $\int_{\mathbb{R}^{D-1}} L^2(\mathbb{R}, d\theta') d\mathbf{x}_\perp$, and hence by Lemma 3.6 and (12) with the constant field $\{H_{U(1)}(I)\}$ of real Hilbert spaces, the right-hand side is identified with $H_{U(1)}(I) \otimes_{\mathbb{R}} L^2(I_\perp, \mathbb{R}, d\mathbf{x}_\perp)$. On the other hand, $H(R)$ is generated by the functions of the form $\hat{g}_+(e^{-\theta'})g_\perp(\mathbf{x}_\perp)$, where $\text{supp } g_+ \subset I, \hat{g}_+(0) = 0$ and $\text{supp } g_\perp \subset I_\perp$, hence they coincide.

The statement about the separating property and cyclicity follows from this decomposition and Lemma 3.1. \square

Next let us consider translations and dilations on each fibre on the null plane. Let $C : \mathbb{R}^{D-1} \rightarrow \mathbb{R}$ be a continuous function. We define the distorted lightlike translations as the following unitary operator on $L^2(\mathbb{R}^D, d\theta' d\mathbf{x}_\perp)$ (and with the identification of \mathfrak{H}_m with it through the spacial decomposition V_S):

$$(T_C \xi)(\theta', \mathbf{x}_\perp) = e^{iC(\mathbf{x}_\perp)e^{-\theta'}} \xi(\theta', \mathbf{x}_\perp). \quad (15)$$

Similarly, we consider the distorted lightlike dilations

$$(D_C \xi)(\theta', \mathbf{x}_\perp) = \xi(\theta' - C(\mathbf{x}_\perp), \mathbf{x}_\perp). \quad (16)$$

If $C = \alpha$ is constant, they coincide with the usual translation and the dilation, respectively, see (12).

Remark 3.9. Note that since C is a continuous function $C, x_\perp \mapsto e^{iC(\mathbf{x}_\perp)te^{-\theta'}}$ is a measurable vector field of bounded operators, that is, $\int_{\mathbb{R}^{D-1}}^\oplus e^{iC(\mathbf{x}_\perp)te^{-\theta'}} \xi(\mathbf{x}_\perp) d\mathbf{x}_\perp \in \mathfrak{H}_m$ for all $\xi \in \mathfrak{H}_m$, with the identification (10). As an intermediate step, if C is a characteristic function, the claim is obvious. For an arbitrary continuous C , there is a family of simple functions $\{C_N\}_{N \in \mathbb{N}}$ converging uniformly to pointwise to C . Then $e^{iC_N(\mathbf{x}_\perp)te^{-\theta'}}$ converges to $e^{iC(\mathbf{x}_\perp)te^{-\theta'}}$ in the strong operator topology. D_C is also a decomposable operator since $(D_C \xi)(\theta', \mathbf{x}_\perp) = \int_{\mathbb{R}^{D-1}}^\oplus D_C(\mathbf{x}_\perp) \xi(\theta', \mathbf{x}_\perp) d\mathbf{x}_\perp$, where $D_C(\mathbf{x}_\perp) \xi(\theta', \mathbf{x}_\perp) = \xi(\theta' - C(\mathbf{x}_\perp), \mathbf{x}_\perp)$, and one can analogously prove that $x_\perp \mapsto D_C(\mathbf{x}_\perp)$ is a measurable vector field of bounded operators passing to the θ' -Fourier transform. In particular, since $H = \int_{\mathbb{R}^{D-1}}^\oplus H(\mathbf{x}_\perp) d\mathbf{x}_\perp \subset \mathfrak{H}_m$ is a standard subspace, the subspaces $T_C H = \int_{\mathbb{R}^{D-1}}^\oplus T_C(\mathbf{x}_\perp) H(\mathbf{x}_\perp) d\mathbf{x}_\perp$ and $D_C H = \int_{\mathbb{R}^{D-1}}^\oplus D_C(\mathbf{x}_\perp) H(\mathbf{x}_\perp) d\mathbf{x}_\perp$ are well-defined and standard as well.

Proposition 3.10. *The family of real subspaces $H(R)$, indexed by open connected regions $R \subset X_-^0$, is covariant with respect to T_C, D_C for a continuous function C , i.e. $T_C H(R) = H(R + C)$ and $D_C H(R) = H(e^C \cdot R)$, where $R + C = \{(x_+ + C(\mathbf{x}_\perp), \mathbf{x}_\perp) : (x_+, \mathbf{x}_\perp) \in R\}$ and $C \cdot R = \{(e^{C(\mathbf{x}_\perp)} x_+, \mathbf{x}_\perp) : (x_+, \mathbf{x}_\perp) \in R\}$.*

Proof. Let g_0 be a thin test function as in (13). The fibre-wise deformed test function $g_{t_C}(x_+, \mathbf{x}_\perp) = g(x_+ - C(\mathbf{x}_\perp), \mathbf{x}_\perp)$ corresponds to the following one-particle vector:

$$\hat{g}_{t_C}(p_-, \mathbf{p}_\perp) = e^{iC(\mathbf{x}_\perp)p_-} (\hat{g})(p_-, \mathbf{p}_\perp),$$

Therefore, we obtain the equality $V_S(Eg_{0,t_C}) = T_C V_S(Eg_0)$ once we prove that $H(R + C)$ contains the vectors which are the Fourier transform of $g_{0,t_C}(x_+, x_-, \mathbf{x}_\perp) = \delta(x_-) g_{t_C}(x_+, \mathbf{x}_\perp)$ where g_{t_C} is of the form $g(x_+ - C(\mathbf{x}_\perp), \mathbf{x}_\perp)$ with a smooth g .

Indeed, we can mollify g_{t_C} by an approximate delta function δ_n on X_-^0 : δ_n is a positive smooth function with $\int \delta_1(x_+, \mathbf{x}_\perp) dx_+ d\mathbf{x}_\perp = 1$ over the Lebesgue measure and $\delta_n(x_+, \mathbf{x}_\perp) = n^D \delta_1(nx_+, n\mathbf{x}_\perp)$. The mollified function $g_{t_C} * \delta_n$ is smooth and supported in $R + C$ for sufficiently large n , its x_- integral vanishes. Therefore, its Fourier transform belongs to $H(R + C)$ when restricted to the mass shell, and is equal to the Fourier transform of g_{t_C} multiplied by that of δ_n . The Fourier transform of δ_n is bounded and converge to 1, therefore, it converges to Eg_{0,t_C} where $g_{0,t_C}(x_+, x_-, \mathbf{x}_\perp) = \delta(x_-)g_{0,t_C}(x_+, \mathbf{x}_\perp)$. As $H(R + C)$ is closed, it contains Eg_{0,t_C} .

Similarly, for $g_{\mathfrak{d}_C}(x_+, \mathbf{x}_\perp) = g(e^{-C(\mathbf{x}_\perp)}x_+, \mathbf{x}_\perp)$, its Fourier transform on x_+ for fixed \mathbf{x}_\perp as a function of p_- is dilated by $e^{C(\mathbf{x}_\perp)}$ and hence

$$(V_S \hat{g}_{\mathfrak{d}_C})(\theta', \mathbf{x}_\perp) = (V_S \hat{g})(\theta' - C(\mathbf{x}_\perp), \mathbf{x}_\perp),$$

that is, it holds that $V_S(Eg_{0,\mathfrak{d}_C}) = D_C V_S(Eg_0)$. Similarly as in the case of g_{t_C} , $Eg_{0,\mathfrak{d}_C} \in H(e^C \cdot R)$, where $g_{0,\mathfrak{d}_C}(x_+, x_-, \mathbf{x}_\perp) = \delta(x_-)g_{\mathfrak{d}_C}(x_+, \mathbf{x}_\perp)$, we obtain the desired covariance. \square

Proposition 3.10 allows to generalize the disintegration (14) to more general open regions.

Proposition 3.11. *Let R be a region on X_-^0 of the form $R = \{(x_+, \mathbf{x}_\perp) : \mathbf{x}_\perp \in I_\perp, C_1(\mathbf{x}_\perp) < x_+ < C_2(\mathbf{x}_\perp)\}$ for an open region $I_\perp \subset \mathbb{R}^{D-1}$ and two continuous functions C_1, C_2 . Then through the spatial decomposition we have*

$$H(R) \simeq \int_{I_\perp}^{\oplus \mathbb{R}} H_{U(1)}((C_1(\mathbf{x}_\perp), C_2(\mathbf{x}_\perp))) d^{D-1} \mathbf{x}_\perp.$$

Similarly, the subspace of a null-cut decomposes:

$$\begin{aligned} H(N_C) &\simeq \int_{\mathbb{R}^{D-1}}^{\oplus \mathbb{R}} H_{U(1)}((C(\mathbf{x}_\perp), \infty)) d^{D-1} \mathbf{x}_\perp, \\ H(N_C^\dagger) &\simeq \int_{\mathbb{R}^{D-1}}^{\oplus \mathbb{R}} H_{U(1)}((-\infty, C(\mathbf{x}_\perp))) d^{D-1} \mathbf{x}_\perp. \end{aligned}$$

Proof. Recall that by Proposition 3.8 the decomposition holds for the case $R = (0, 1) \times I_\perp$. We can dilate this by $D_{C_2-C_1}$ and then translate it by C_1 to arrive at R in the desired form. By the covariance of the whole family $\{H(R)\}$ with respect to fibre-wise translations and dilations by Proposition 3.10, we obtain the desired decomposition.

If we start from the rectangle $\mathbb{R}_\pm \times \mathbb{R}^{D-1}$, then we deform the rectangle via the distorted lightlike translation T_C to the null-cut N_C or the interior of its complement N_C^\dagger , respectively. The interior of the complement N_C^\dagger of the null-cut N_C on the null-plane is:

$$N_C^\dagger = X_-^0 \setminus \overline{N_C} = \{x_+ < C(\mathbf{x}_\perp), x_- = 0, \mathbf{x}_\perp \in \mathbb{R}^{D-1}\}.$$

From the covariance of T_C and Proposition 3.8, we obtain the decompositions:

$$\begin{aligned} H(N_C) &\simeq \int_{\mathbb{R}^{D-1}}^{\oplus \mathbb{R}} H_{U(1)}((C(\mathbf{x}_\perp), \infty)) d^{D-1} \mathbf{x}_\perp \\ H(N_C^\dagger) &\simeq \int_{\mathbb{R}^{D-1}}^{\oplus \mathbb{R}} H_{U(1)}((-\infty, C(\mathbf{x}_\perp))) d^{D-1} \mathbf{x}_\perp. \end{aligned} \tag{17}$$

\square

3.4 Duality for null cuts

Theorem 3.12. *The standard subspaces $H(N_C)$ of continuous C satisfy duality (see Figure 1):*

$$H(N_C) = H(N_C)'' = H(N_C'').$$

Proof. The first equality follows from the fact that $H(N_C)$ is a standard subspace.

Next we show $H(N_C) \subset H(N_C'')$. By any shift ta in the x_1 -direction (or x_- -direction), we have $N_C \subset N_C'' - ta$ for $t > 0$. Indeed, if $x \in N_C, y \in N_C'$, it has $x_- = 0$, and $(x - y)^2 = -2(x_+ - y_+)y_- - (\mathbf{x}_\perp - \mathbf{y}_\perp)^2 < 0$. Furthermore, as x_+ can be arbitrarily large, it must hold that $x_+ - y_+ > 0, y_- > 0$ and hence adding ta to y just decreases the first term. Therefore, for any $y \in N_C', y + ta$ is spacelike from $x \in N_C$, that is, $N_C \subset N_C'' + ta$. Therefore, by Lemma 3.7, we have $H(N_C) \subset H(N_C'')$.

Lastly we prove $H(N_C'') \subset H(N_C)$. It is easy to see that $N_C^\dagger - ta \subset N_C'$ for any $t > 0$, where a is a positive vector in the x_1 -direction, hence we have $H(N_C^\dagger) \subset H(N_C')$ by Lemma 3.7. As we have $H(N_C'') \subset H(N_C')'$ by locality and we have $H(N_C'') \subset H(N_C')' \subset H(N_C^\dagger)' = H(N_C)''$, where the last equality follows from Lemma 3.1 and (17). \square

As an example, it is immediate to see that the causal completion of the zero null-cut $N_0 = N_{C_0}$ with $C_0(\mathbf{x}_\perp) = 0$ is the right wedge: $N_0'' = W_1$, hence $H(N_0) = H(N_0'') = H(W_1)$.

While it is possible to define $H(N_C)$ for non continuous curves (see Section 6), the continuity condition Theorem 3.12 cannot be removed. Indeed, if $C = \chi_{\mathbb{Q}^{D-1}}$ is the characteristic function of \mathbb{Q}^{D-1} then $H(N_C) = H(W_R)$ since the direct integrals

$$H(N_C) = \int_{\mathbb{R}^{D-1}}^{\oplus \mathbb{R}} H_{U(1)}((\chi_{\mathbb{Q}^{D-1}}(\mathbf{x}_\perp), +\infty)) d^{D-1} \mathbf{x}_\perp = \int_{\mathbb{R}^{D-1}}^{\oplus \mathbb{R}} H_{U(1)}((0, +\infty)) d^{D-1} \mathbf{x}_\perp = H(W_1)$$

are equal up to a null measure set, but $H(N_C'')$ is strictly contained in $H(W_R)$ when, again, is considered the definition (5).

Let us define the deformed wedge for a continuous function $C : \mathbb{R}^{D-1} \rightarrow \mathbb{R}$ by

$$W_C := \{\mathbf{x} \in \mathbb{R}^{D+1}, x_- < 0, x_+ > C(\mathbf{x}_\perp)\}.$$

Then we have:

Proposition 3.13. $H(N_C) = H(W_C)$.

Proof. It is clear from the definition that $N_C \subset W_C - tx_-$ for every $t > 0$. From Lemma 3.7, it follows $H(N_C) \subset H(W_C)$.

For the opposite direction, we will deduce $H(W_C) \subset H(N_C'')$. The claim follows from isotony and the previous theorem. Let $y \in W_C$, then we can write $y = x + a$ with $x \in N_C$ and $a = (0, a_-, 0)$. It is enough to prove $N_C - ta \subset N_C''$ for $t > 0$, or equivalently, $N_C \subset N_C'' + ta$ for every $\varepsilon > 0$. This is contained in the proof of Theorem 3.12. \square

4 Modular operator for null cuts

The following is of an independent interest, because it is an assumption of the limited version of QNEC [CF20].

Proposition 4.1. *Let $C_1, C_2 : \mathbb{R}^{D-1} \rightarrow \mathbb{R}$ be continuous functions such that $C_1(\mathbf{x}_\perp) < C_2(\mathbf{x}_\perp)$. Then the inclusion $H(N_{C_2}) \subset H(N_{C_1})$ is a HSMI.*

Proof. Note that, if $K \subset H$ is a HSMI and U is a unitary, then also $UK \subset UH$ is a HSMI by Lemma 2.1.

Recall that¹ $H(W_+) \subset H(W_1)$ is a HSMI, where $C_+(\mathbf{x}_\perp) = 1$ and $W_+ = W_{C_+}$ is the shifted wedge. Put also $C_0(\mathbf{x}_\perp) = 0$, so that $W_1 = W_{C_0}$. By fibre-wise covariance (Proposition 3.10), the inclusion $H(N_{C_2}) \subset H(N_{C_1})$ is unitarily equivalent to $H(W_+) = H(C_+) \subset H(C_0) = H(W_1)$. Indeed, we can take $C_{\text{dl}}(\mathbf{x}_\perp) = \ln(C_2(\mathbf{x}_\perp) - C_1(\mathbf{x}_\perp))$ and $C_{\text{tr}}(\mathbf{x}_\perp) = C_1(\mathbf{x}_\perp)$. Then we have $T_{C_{\text{tr}}} D_{C_{\text{dl}}} H(W_1) = H(C_{\text{tr}} + e^{C_{\text{dl}}} C_0) = H(N_{C_1})$ and $T_{C_{\text{tr}}} D_{C_{\text{dl}}} H(W_+) = H(C_{\text{tr}} + e^{C_{\text{dl}}} C_+) = H(N_{C_2})$. Then the desired HSMI follows from this unitary equivalence. \square

In the rest of this Section, we will study the modular operator of the standard spaces $H(N_C)$. We call Q_S the unitary that results from the concatenation of V_S (cf. Section 3.2) and the fibre-wise unitary between $L^2(\mathbb{R}, d\theta')$ and the $U(1)$ -current mentioned in Section 2.4.

Proposition 4.2. *The unitary operator Q_S between \mathfrak{H}_m and the direct integral of $U(1)$ -currents decomposes the modular operator of $C(N_0)$:*

$$\begin{aligned} \log(\Delta_{H(N_0)}) &= \log(\Delta_{H(W_R)}) \\ &= \text{Ad}_{Q_S} \int_{\mathbb{R}^{D-1}}^{\oplus} \log(\Delta_{H_{U(1)}(\mathbb{R}_+)}) d^{D-1} \mathbf{x}_\perp. \end{aligned}$$

Proof. The causal complement of N_0 is the left wedge W_L :

$$N'_0 = \{x \in \mathbb{R}^{D+1}; x_+ < 0, x_- > 0, \mathbf{x}_\perp \in \mathbb{R}^{D-1}\} = W_L.$$

By Theorem 3.12, we know $H(N_0) = H(N''_0) = H(W_R)$. By the Bisognano-Wichmann property (HK6) in Section 2.2, we know that the modular group of the right wedge coincides with the representation of the boost group along x_1 :

$$\Delta_{H(W_R)}^{it} = U_m(\Lambda_1(-2\pi t)).$$

By (16) the representation of the boost group is equivalent to the constant dilation group of the $U(1)$ -current on each fibre of the spatial decomposition of \mathfrak{H}_m and hence:

$$\text{Ad}_{Q_S} \Delta_{H(W_R)}^{-it} = \int_{\mathbb{R}^{D-1}}^{\oplus} U_{U(1)}(2\pi t, 0) d^{D-1} \mathbf{x}_\perp.$$

The statement follows from Theorem A.1 and the Bisognano-Wichmann property for the $U(1)$ -current (cf. Section 2.4). \square

¹With this slightly unfortunate notations, W_1 is the standard wedge in the x_1 -direction (and not the shifted wedge).

The standard subspace of the constant null cut $C_s(\mathbf{x}_\perp) = s$ is

$$H(N_s) = T_{C_s} H(N_0) = U_m(1, \mathbf{t}_{x_+}(s)) H(W_1) = H(\mathbf{t}_{x_+}(s) W_1).$$

That means that $H(N_s)$ equals the standard subspace associated with the wedge $\mathbf{t}_{x_+}(s) W_1$. According to the Bisognano-Wichmann property, the associated modular group is the boost group leaving the wedge $\mathbf{t}_{x_+}(s) W_1$ invariant:

$$\begin{aligned} \Delta_{H(N_s)}^{it} &= U_m(1, \mathbf{t}_{x_+}(s)) \Delta_{H(N_0)}^{it} U_m(1, \mathbf{t}_{x_+}(s))^* \\ &= U_m(1, \mathbf{t}_{x_+}((1 - e^{2\pi t})s)) \Delta_{H(N_0)}^{it}. \end{aligned}$$

From this, we have the following well-known (cf. Lemma 2.1) relation between generators.

$$\log \Delta_{H(N_s)} = \log \Delta_{H(N_0)} + 2\pi s P, \quad (18)$$

where P is the generator of translations $U_m(\mathbf{t}_{x_+}(s))$ along the light ray x_+ .

The formula (18) has a natural generalization to arbitrary null-cuts. The key structures that we will utilize are half-sided modular inclusion of certain null-cut subspaces. In the following, we will use the notation for half-sided modular inclusions introduced in Section 2.1.

Now we state one of our main results.

Theorem 4.3. *Let $C : \mathbb{R}^{D-1} \rightarrow \mathbb{R}$ be a continuous function, then the generator of the modular group of the associated null-cut is decomposed as follows, where the equivalence is given by the operator Q_S of Proposition 4.2.*

$$\log(\Delta_{H(N_C)}) \simeq \int_{\mathbb{R}^{D-1}}^{\oplus} \left(\log(\Delta_{H_{U(1)}}(\mathbb{R}_+)) + 2\pi C(\mathbf{x}_\perp) P_{\mathbf{x}_\perp} \right) d\mathbf{x}_\perp, \quad (19)$$

where $P_{\mathbf{x}_\perp}$ is the generator of translations on each fibre.

Proof. As we have $N_C = T_C N_0$ (with the notations of Lemma 3.7), we have $\Delta_{H(N_C)}^{it} = \text{Ad } T_C(\Delta_{H(N_0)}^{it})$. Note that T_C does the fibre-wise translation, therefore by the decomposition through Q_S ,

$$\begin{aligned} \Delta_{H(N_C)}^{it} &\simeq \int_{\mathbb{R}^{D-1}}^{\oplus} \text{Ad } U_{U(1)}(0, \mathbf{t}(C(\mathbf{x}_\perp)))(\Delta_{H_{U(1)}}(\mathbb{R}_+)^{it}) d\mathbf{x}_\perp \\ &= \int_{\mathbb{R}^{D-1}}^{\oplus} \exp \left(it \left(\log(\Delta_{H_{U(1)}}(\mathbb{R}_+)) + 2\pi C(\mathbf{x}_\perp) P_{\mathbf{x}_\perp} \right) \right) d\mathbf{x}_\perp, \end{aligned}$$

where the second equation is the fibre-wise transformation law of a simple HSMI (Theorem 2.3). By Proposition A.1 and the Stone theorem, this relation passes to the generators. \square

Let us comment on the second quantized net $(\mathcal{A}, \Gamma_+(U_m), \mathcal{F}_+(\mathfrak{H}_m))$. By Proposition 2.4, a HSMI of standard subspaces promotes to a HSMI of von Neumann algebras, hence the following is an immediate consequence of Proposition 4.1 and Theorem 3.12.




Figure 2: Geometric action of distorted lightlike translations.

Corollary 4.4. *Let $C_1, C_2 : \mathbb{R}^{D-1} \rightarrow \mathbb{R}$ be continuous functions such that $C_1(\mathbf{x}_\perp) < C_2(\mathbf{x}_\perp)$. Then the inclusion $\mathcal{A}(N_{C_2}'') \subset \mathcal{A}(N_{C_1}'')$ is a HSMI.*

It is not easy to write the second-quantized version of (19): while the second quantization of the left-hand side is straightforward $d\Gamma_+(\log \Delta_{H(N_C)}) = \log \Delta_{\mathcal{A}(N_C''), \Omega}$, where $d\Gamma_+(A) = 0 \oplus A \oplus (A \otimes \mathbb{1} \oplus \mathbb{1} \otimes A) \cdots$ is the additive second quantization, the direct-integral structure on the right-hand side translates to the *continuous tensor product* structure in the second quantization [AW66, Nap71]. Conceptually, it should be an “integral” of the second-quantized generators on fibres, but we do not know how to formulate such an integral corresponding to (1). Instead, we were able to formulate the HSMI without reference to the (undefined) stress-energy tensor smeared on the null plane.

On the other hand, for general interacting models, we do not have this simple second-quantization structure, and the validity of HSMI for the inclusions of null cut regions remains open, see Section 6.

5 Relative entropy and Energy bounds

5.1 Decomposition of the relative entropy for coherent states

We recall that the formula for Araki relative entropy of a von Neumann algebra $\mathcal{A} \subset \mathcal{B}(\mathcal{K})$ with respect to two vector states ω_1 and ω_2 is given by

$$S(\omega_1 || \omega_2) = -\langle \xi, \log \Delta_{\eta, \xi} \xi \rangle$$

where $\xi, \eta \in \mathcal{K}$ implement ω_1 and ω_2 on \mathcal{A} , respectively and $S_{\eta, \xi} = J_{\eta, \xi} \Delta_{\eta, \xi}^{\frac{1}{2}}$ is the relative Tomita operator:

$$S_{\eta, \xi} : \mathcal{A}\xi \ni a\xi \mapsto a^* \eta \in \mathcal{A}\eta.$$

It is immediate to see that if $\eta = \xi$, then $\Delta_{\xi,\xi}$ is the modular operator of \mathcal{A} with respect to ξ and $S(\omega_1||\omega_1) = 0$.

Let $H \subset \mathcal{H}$ be a standard subspace and $w(\psi)$ be the Weyl operator of $\psi \in \mathcal{H}$ on $\Gamma_+(\mathcal{H})$. A state on a von Neumann algebra $\mathcal{A} \subset \mathcal{B}(\Gamma_+(\mathcal{H}))$ is called **coherent** if it is given by $\omega_\psi(\cdot) := \omega(w(\psi) \cdot w(\psi)^*)$, where ω is the Fock vacuum.

Following [CLR20], given a standard subspace $H \subset \mathcal{H}$, we can consider the following quantity

$$S_H(\psi) = \text{Im} \langle \psi, P_H i \log \Delta_H \psi \rangle,$$

where P_H is an unbounded real projection called *cutting projection*. The cutting projection is determined by the modular operator and modular conjugation of H . Its explicit form is:

$$P_H = a(\Delta_H) + J_H b(\Delta_H)$$

where $a(\lambda) = \lambda^{-1/2}(\lambda^{-1/2} - \lambda^{1/2})^{-1}$ and $b = (\lambda^{-1/2} - \lambda^{1/2})^{-1}$.

If $h \in H$, then

$$S_H(\psi) = -\langle \psi, \log \Delta_H \psi \rangle.$$

We have the following relation with the relative entropy of second quantization algebra:

$$S_{R(H)}(\omega_\psi||\omega) = S_H(\psi).$$

where $S_{R(H)}(\omega_\psi||\omega)$ is the relative entropy of ω_ψ and ω with respect to $R(H)$ [CLR20, Proposition 4.2].

Lemma 5.1. *Consider a standard subspace H that is a direct integral of standard subspaces $H = \int_X^{\oplus} H(x) d\nu(\lambda) \subset \int_X^{\oplus} \mathcal{K}_\lambda d\nu(\lambda)$ with decomposable modular generator $\log(\Delta_H) = \int_X^{\oplus} \log(\Delta_{H(\lambda)}) d\nu(\lambda)$. Moreover, let $\psi = \int_X^{\oplus} \psi(\lambda) d\nu(\lambda) \in \mathcal{K}$. Then the relative entropy of ω_ψ and the vacuum with respect to H decomposes:*

$$S_{R(H)}(\omega_\psi||\omega) = \int_X S_{R(H(\lambda))}(\omega_{\psi(\lambda)}(\lambda)||\omega(\lambda)) d\nu(\lambda),$$

where $\omega(\lambda)$ denotes the vacuum on the Fock space of \mathcal{K}_λ .

Proof. Following the previous arguments, we have $S_{R(H)}(\omega_\psi||\omega) = S_H(\psi)$. By assumption the modular generator decomposes. It is straightforward to check that $J_H = \int_X^{\oplus} J_{H(\lambda)} d\nu(\lambda)$ decomposes as well. Applying Theorem A.1, the cutting projection decomposes accordingly $P_H = \int_X^{\oplus} P_{H(\lambda)} d\nu(\lambda)$. Thus

$$\langle \psi, P_H i \log \Delta_H \psi \rangle = \int_X \langle \psi(\lambda), P_{H(\lambda)} i \log(\Delta_{H(\lambda)}) \psi(\lambda) \rangle d\nu(\lambda),$$

in particular

$$\begin{aligned} S_H(\psi) &= \text{Im} \int_X \langle \psi(\lambda), P_{H(\lambda)} i \log(\Delta_{H(\lambda)}) \psi(\lambda) \rangle d\nu(\lambda) \\ &= \int_X \text{Im} \langle \psi(\lambda), P_{H(\lambda)} i \log(\Delta_{H(\lambda)}) \psi(\lambda) \rangle d\nu(\lambda) = \int_X S_{H(\lambda)}(\psi(\lambda)). \end{aligned}$$

Since $S_{H(\lambda)}(\psi(\lambda)) = S_{R(H(\lambda))}(\omega_{\psi(\lambda)}(\lambda)||\omega(\lambda))$, we conclude the argument. \square

Remark 5.2. The modular operator of the direct integral of standard subspaces always decomposes into the direct integral of modular operators of the fibre subspaces. This can be seen from the KMS condition for standard subspaces (see [Lon08, Proposition 2.1.8]) and the results in Appendix A.

5.2 Relative entropy for coherent states on the null plane

For a single $U(1)$ -current net, an explicit formula for the relative entropy was computed in [Lon20], where an equivalent definition of the $U(1)$ -current mentioned in Section 2.4 is used. We translate the results to the definition of the $U(1)$ -current that we are working with.

Let $I_t = (t, +\infty) \subset \mathbb{R}$, $H_{U(1)}(I_t)$ be its standard subspace and $\mathcal{A}(I_t) = R(H_{U(1)}(I_t))$ be the second quantization von Neumann algebra. Consider the map $\beta_k(w(\xi)) = e^{-i \int_{\mathbb{R}} k(x) L(x) dx} w(\xi)$ with $k, l \in C_0^\infty(\mathbb{R}, \mathbb{R})$, $\hat{l}(0) = 0$, $C_0^\infty(\mathbb{R}, \mathbb{R}) \ni L(x) = \int_{-\infty}^x l(s) ds$ the primitive of l and $\xi \in H_{U(1)}(I_t)$ being the one-particle vector of l . Then β_k extends to an automorphism of the local algebra $\mathcal{A}(I_t)$, which is called a Buchholz-Mack-Todorov (BMT) automorphism [BMT88]. We denote this extension by β_k as well. One has [Lon20, Theorem 4.7]:

$$S_{\mathcal{A}(I_t)}(\omega \circ \beta_k || \omega) = \pi \int_t^{+\infty} (x - t) k^2(x) dx,$$

where ω is the vacuum state on the $U(1)$ -current.

We can generalize BMT automorphisms to the direct integral of $U(1)$ -currents. Let $h, g \in C_0^\infty(X_-^0, \mathbb{R})$ with $\hat{g}(0) = 0$ and $\xi = V_S E \delta(x_-) g$, we define the automorphism

$$\beta_h(w(\xi)) = e^{-i \int_{X_-^0} h(x) G(x) dx} w(\xi).$$

Let us consider the relative entropy between ω and $\omega \circ \beta_h$ with respect to local algebras of null cuts:

Proposition 5.3. *We have*

$$S_{R(H(N_C))}(\omega \circ \beta_h || \omega) = \pi \int_{\mathbb{R}^{D-1}} \int_{C(\mathbf{x}_\perp)}^\infty (x_+ - C(\mathbf{x}_\perp)) h(x_+, \mathbf{x}_\perp)^2 dx_+ d\mathbf{x}_\perp.$$

Proof. As the support of h is compact, C is bounded on the restriction to \mathbf{x}_\perp of the support of h , and by adding an appropriate smooth function supported in N_C^\dagger , we may assume that $\int h(x_+, \mathbf{x}_\perp) dx_+ = 0$ for every \mathbf{x}_\perp without changing the state $\omega \circ \beta_h$ on $R(H(N_C))$. Then $\delta(x_-)h$ is a thin test function and we call $\mathfrak{h} := V_S E \delta(x_-)h$ its one-particle vector in the spatial decomposition.

In this case, the BMT automorphism β_h is generated by the adjoint action of the Weyl operator $w(\mathfrak{h})$. The symplectic form can be computed fibrewise using (7)

$$\text{Ad } w(\mathfrak{h}) (w(\xi)) = e^{2i \text{Im} \langle \mathfrak{h}, \xi \rangle} w(\xi). \quad (20)$$

$$\text{Im} \langle \mathfrak{h}, \xi \rangle = \int_{\mathbb{R}^{D-1}} \text{Im} \langle \mathfrak{h}(\mathbf{x}_\perp), \xi(\mathbf{x}_\perp) \rangle_{U(1)} d\mathbf{x}_\perp = \frac{1}{2} \int_{\mathbb{R}^D} h(x_+, \mathbf{x}_\perp) G(x_+, \mathbf{x}_\perp) dx_+ d\mathbf{x}_\perp. \quad (21)$$

Thus $\omega \circ \beta_h$ is the coherent state $\omega_{\mathfrak{h}}$ and we can apply Lemma 5.1:

$$S_{R(H(N_C))}(\omega_{\mathfrak{h}}||\omega) = \int_{\mathbb{R}^{D-1}} S_{\mathcal{A}(I_C(\mathbf{x}_{\perp}))}(\omega_{\mathfrak{h}(\mathbf{x}_{\perp})}(\mathbf{x}_{\perp})||\omega_{\mathbf{x}_{\perp}})d\mathbf{x}_{\perp}.$$

The coherent state $\omega_{\mathfrak{h}(\mathbf{x}_{\perp})}(\mathbf{x}_{\perp})$ on the $U(1)$ current is generated by the adjoint action of $w_{U(1)}(\mathfrak{h}(\mathbf{x}_{\perp}))$ on the fibrewise vacuum $\omega(\mathbf{x}_{\perp})$. This in turn coincides with the action of the BMT automorphism $\beta_{h(\mathbf{x}_{\perp})}$ on the $U(1)$ vacuum $\omega(\mathbf{x}_{\perp})$. Therefore, we can insert the results for a single $U(1)$ -current and have:

$$\begin{aligned} S_{R(H(N_C))}(\omega \circ \beta_h||\omega) &= \int_{\mathbb{R}^{D-1}} S_{\mathcal{A}(I_C(\mathbf{x}_{\perp}))}(\omega(\mathbf{x}_{\perp}) \circ \beta_{h(\mathbf{x}_{\perp})}||\omega(\mathbf{x}_{\perp}))d\mathbf{x}_{\perp} \\ &= \pi \int_{\mathbb{R}^{D-1}} \int_{C(\mathbf{x}_{\perp})}^{\infty} (x_+ - C(\mathbf{x}_{\perp}))h(x_+, \mathbf{x}_{\perp})^2 dx_+ d\mathbf{x}_{\perp}. \end{aligned}$$

□

5.3 The ANEC and the QNEC

Energy conditions in QFT One of the motivations to study quantum energy conditions comes from energy constraints in General Relativity, see e.g. [Few12]. Indeed, pointlike positivity of the stress energy tensor, assumed in general relativity to avoid singular spacetimes, is not possible in QFT. It is not difficult to see that the stress energy smeared by a positive test function can not be positive but only bounded from below, see e.g. [FH05]. On the other hand, the ANEC claims that the the integral of the expectation value of the energy-momentum tensor in any physical state, along any complete, lightlike geodesic γ is always non-negative (see e.g. [FLPW16, (3)], [Ver00]):

$$\langle \Phi | \int_{\mathbb{R}} ds T_{\mu,\nu}(\gamma(s)) k^{\mu} k^{\nu} \Phi \rangle \geq 0. \quad (22)$$

where κ^{μ} is the tangent of γ . We can write T_{++} when we consider the null x_+ -direction $(x_+, 0, \mathbf{0})$. It is expected that (22) is satisfied for a dense set of vectors in a certain formal sense.

The Quantum Null Energy (QNEC) is a local bound on the expectation value of the null energy density given by the von Neumann entropy of the null cut (the following is only symbolic and we do not attempt to justify the expressions):

$$\langle T_{++}(x) \rangle_{\Phi} \geq \frac{1}{2\pi} S''_{\mathcal{A}(N_C), \Phi},$$

where N_C is a null cut and x is on the boundary of the cut C , and the derivative is taken in the sense of the deformation of C at the point x (see [BFKW19]).

In [BFLW16] for null cuts N_C and in [LLSM18] for deformed wedges W_C it is claimed that the QNEC is equivalent to the positivity of the second derivative of the relative entropy between the vacuum and another state with respect to null-deformations:

$$S'' \geq 0. \quad (23)$$

The equivalence is reached through the expression of the modular operator as in (1). With this motivation the positivity of S'' has been checked rigorously in some families of AQFT models (see e.g. [Lon20, CLR20, Lon19, Pan20]) and is expected to hold in general in the AQFT context.

In [CF20], the authors assume that the inclusions of null cut algebras are HSMI. For continuous functions $C_1 \geq C_2$ and $A \geq 0$, they showed

$$\partial_t^+ S_{R(N_{C_1+tA})}(\Psi||\Omega)\Big|_{t=0} - \partial_t^- S_{R(N_{C_2+tA})}(\Psi||\Omega)\Big|_{t=0} \geq 0, \quad (24)$$

where ∂_t^\pm denote half-sided partial derivatives. This statement was proved for the class of vectors satisfying (22) with respect to T_{++} and having finite relative entropy with respect to the vacuum and the algebra of the zero null cut.

We will verify the QNEC (23) for the free scalar field and for some states explicitly. In order to do this, we take the decomposition of the modular operator (cf. Theorem 4.3) and exploit the result by [Lon20] on each fiber. In this sense, our findings constitute a generalization of the results to a continuum family of $U(1)$ -currents.

The ANEC To make contact with the physical literature, let us consider distorted lightlike translations T_A by $A \geq 0$, which generate half-sided modular inclusions of standard subspaces $H(N_{C+A}) \subset H(N_C)$ of null-cuts. It is claimed, e.g. [CTT17b, (5.6) and (5.18)], that the following operators are positive.

$$P_{\mathbf{x}_\perp} = \int dx_+ T_{++}(x_+, \mathbf{x}_\perp)$$

$$H_A = \int d^{D-1} \mathbf{x}_\perp A(\mathbf{x}_\perp) \int dx_+ T_{++}(x_+, \mathbf{x}_\perp) = \int d^{D-1} \mathbf{x}_\perp A(\mathbf{x}_\perp) P_{\mathbf{x}_\perp}.$$

Furthermore, it is argued that positivity of H_A should imply ANEC (22) in [FLPW16, Section 3.2].

Note that the last expression does not involve T_{++} , and we can make sense of it in the free field if we interpret these relations at the one-particle level. Indeed, it is simply a weighted integral of the generator $P_{\mathbf{x}_\perp}$ of translations in the $U(1)$ -current, which is positive. The operator H_A is also positive as it is the second quantization of this positive operator.

Following [Lon20], we define the vacuum energy associated to null deformations by $A \geq 0$ of a normal and faithful state φ , that has a vector representative η , by

$$E_A(\eta) = \langle \eta, H_A \eta \rangle, \quad (25)$$

where $H_A = \frac{1}{2\pi} (\log(\Delta_{R(H(N_{C+A})), \Omega}) - \log(\Delta_{R(H(N_C)), \Omega}))$. The expression (25) is positive for the dense set of vectors in $\mathcal{D}(H_A)$ (cf. Appendix A for explicit form of $\mathcal{D}(H_A)$) since H_A is a positive operator. In [Lon20] and [CF20], the positivity of (25) is considered as form of the ANEC.

The QNEC A BMT automorphism of the form (20) generates a coherent state $\omega_h(\cdot) = \omega(w(\mathfrak{h}) \cdot w(\mathfrak{h})^*)$ when restricted to a null cut. The GNS representation space of $\omega \circ \beta_h$ is the

Fock space. The representation of \mathbf{P} transforms by the adjoint action of $w(\mathfrak{h})$. We call this representation the β_h -representation.

Let $\omega \circ \beta_{h_1}$ and $\omega \circ \beta_{h_2}$ be two coherent states that are unitarily generated by the adjoint action of $w(\mathfrak{h}_1)$ and $w(\mathfrak{h}_2)$, respectively. The relative entropy between these states is:

$$S(\omega \circ \beta_{h_1} || \omega \circ \beta_{h_2}) = S(\omega \circ \beta_{h_1} || \omega \cdot \text{Ad } w(\mathfrak{h}_2)) = S(\omega \cdot \text{Ad } w(\mathfrak{h}_1 - \mathfrak{h}_2) || \omega) = S(\omega \circ \beta_{h_1 - h_2} || \omega).$$

To study the relative entropy between these states, we can therefore restrict to $h_2 = 0$, i.e. $\omega \circ \beta_{h_2}$ being the vacuum state.

A distorted lightlike translation T_{tA} by A maps $H(N_C)$ to $H(N_{C+tA})$ and the relative entropy of $\omega \circ \beta_h$ and ω changes accordingly (cf. Proposition 5.3). The differentiation with respect to the deformation parameter t gives:

$$\begin{aligned} \frac{d}{dt} S_{R(N_{C+tA})}(\omega \circ \beta_h || \omega) &= -\pi \int_{\mathbb{R}^{D-1}} \int_{C(\mathbf{x}_\perp) + tA(\mathbf{x}_\perp)}^\infty A(\mathbf{x}_\perp) h(x_+, \mathbf{x}_\perp)^2 dx_+ d\mathbf{x}_\perp \\ \frac{d^2}{dt^2} S_{R(N_{C+tA})}(\omega \circ \beta_h || \omega) &= \pi \int_{\mathbb{R}^{D-1}} A(\mathbf{x}_\perp)^2 h(C(\mathbf{x}_\perp) + tA(\mathbf{x}_\perp), \mathbf{x}_\perp)^2 d\mathbf{x}_\perp \geq 0 \end{aligned} \quad (26)$$

We applied the differentiation fibrewise and used

$$\frac{d}{dt} \int_{C+tA(\mathbf{x}_\perp)}^\infty (x_+ - (C + tA(\mathbf{x}_\perp))) h(x_+, \mathbf{x}_\perp)^2 dx_+ = \int_{C+tA(\mathbf{x}_\perp)}^\infty A(\mathbf{x}_\perp) h(x_+, \mathbf{x}_\perp)^2 dx_+ \quad (27)$$

$$\frac{d^2}{dt^2} \int_{C+tA(\mathbf{x}_\perp)}^\infty A(\mathbf{x}_\perp) h(x_+, \mathbf{x}_\perp)^2 dx_+ = A(\mathbf{x}_\perp)^2 h(C + tA(\mathbf{x}_\perp), \mathbf{x}_\perp)^2. \quad (28)$$

To justify the interchange of the \mathbf{x}_\perp -integral and the $\frac{d}{dt}$ -derivative, we verify that (27), (28) $\in L^1(\mathbb{R}^{D-1}, d^{D-1}\mathbf{x}_\perp)$. We estimate (27) $\leq \|A(\mathbf{x}_\perp) h(x_+, \mathbf{x}_\perp)^2\|_1$. Since A is continuous and h is compactly supported and smooth, the estimate is bounded and compactly supported in \mathbb{R}^{D-1} and as such it is integrable. For (28), we note that $C + tA(\mathbf{x}_\perp)$ is continuous on \mathbb{R}^{D-1} and accordingly maps compact sets to compact sets. Hence, the product in (28) is compactly supported in \mathbb{R}^{D-1} and bounded and therefore integrable.

In summary, we have proven the following form of the Quantum Null Energy Condition:

Corollary 5.4. *For the coherent states $\omega \circ \beta_h$ considered here, the QNEC holds:*

$$\frac{1}{2\pi} \frac{d^2}{dt^2} S_{R(N_{C+tA})}(\omega \circ \beta_h || \omega) \geq 0.$$

This inequality is not saturated at every point of positive energy density. Following the arguments in Theorem 3.12 and Proposition 3.13, we can replace the region N_{C+tA} with N''_{C+tA} and W_{C+tA} , respectively.

As expected in [BFK⁺16], we recover the ANEC by integrating the QNEC along a null-

direction for coherent states by (25):

$$\begin{aligned}
& \frac{1}{2\pi} \int_{\mathbb{R}} dt \frac{d^2}{dt^2} S_{R(N_{C+tA})}(\omega \circ \beta_h || \omega) \\
&= \frac{1}{2} \int_{\mathbb{R}} dt \int_{\mathbb{R}^{D-1}} A(\mathbf{x}_{\perp})^2 h(C(\mathbf{x}_{\perp}) + tA(\mathbf{x}_{\perp}), \mathbf{x}_{\perp})^2 d\mathbf{x}_{\perp} \\
&= \frac{1}{2} \int_{\mathbb{R}} dx_+ \int_{\mathbb{R}^{D-1}} A(\mathbf{x}_{\perp}) h(x_+, \mathbf{x}_{\perp})^2 d\mathbf{x}_{\perp} \\
&= \text{Im} \langle \mathfrak{h}, iPA\mathfrak{h} \rangle = i \left. \frac{d}{ds} \right|_{s=0} \langle \mathfrak{h}, U_A(s)\mathfrak{h} \rangle \\
&= i \left. \frac{d}{ds} \right|_{s=0} \left(e^{-\frac{1}{2}(\|\mathfrak{h}\|^2 + \|U_A(s)\mathfrak{h}\|^2)} e^{\langle \mathfrak{h}, U_A(s)\mathfrak{h} \rangle} \right) = i \left. \frac{d}{ds} \right|_{s=0} \langle w(\mathfrak{h})\Omega, \Gamma(U_A(s))w(\mathfrak{h})\Omega \rangle \\
&= \langle w(\mathfrak{f})\Omega, H_A w(\mathfrak{h})\Omega \rangle,
\end{aligned}$$

where we used that the self-adjoint fibrewise "momentum operator" $\int_{\mathbb{R}^{D-1}}^{\oplus} A(\mathbf{x}_{\perp}) P_{\mathbf{x}_{\perp}} d\mathbf{x}_{\perp}$ is the generator of $U_A(s)$, similar arguments as in (7) and (21) and the general identity (6) for the Weyl unitaries. This is in agreement with [Lon20, Corollary 3.10 and (46)].

We interpret $A(\mathbf{x}_{\perp})h(x_+, \mathbf{x}_{\perp})^2$ as vacuum energy density at the point $(x_+, \mathbf{x}_{\perp}) \in X_-^0$ of the state $w(\mathfrak{h})\Omega$ with respect to null-deformations A . This is the vacuum energy density of the state $w(\mathfrak{h}(\mathbf{x}_{\perp}))\Omega_{\mathbf{x}_{\perp}}$ of the $U(1)$ -current at \mathbf{x}_{\perp} -fibre with respect to deformations by $A(\mathbf{x}_{\perp})$ (cf. [Lon20]). With this interpretation, the energy averaged over the a null-cut N_C is:

$$E_{A, \beta_h}(N_C) = \int_{\mathbb{R}^{D-1}} \int_{C(\mathbf{x}_{\perp})}^{\infty} A(\mathbf{x}_{\perp}) h(x_+, \mathbf{x}_{\perp})^2 d\mathbf{x}_{\perp}, \quad (29)$$

and we have the following connection between the first derivative of the relative entropy and the energy localised in the null-cut by comparing (26) and (29):

$$\frac{d}{dt} S_{R(N_{C+tA})}(\omega \circ \beta_h || \omega) = E_{A, \beta_h}(N_{C+tA}).$$

Strong superadditivity of relative entropy Consider two null-cuts N_{C_1} and N_{C_2} and the null-cuts $N_{C_{\cup}}$ and $N_{C_{\cap}}$ generated by $C_{\cup}(\mathbf{x}_{\perp}) = \min\{C_1(\mathbf{x}_{\perp}), C_2(\mathbf{x}_{\perp})\}$ and $C_{\cap}(\mathbf{x}_{\perp}) = \max\{C_1(\mathbf{x}_{\perp}), C_2(\mathbf{x}_{\perp})\}$ respectively. Then the strong superadditivity of relative entropy is:

$$S_{R(N_{C_{\cup}})}(\Psi || \Omega) + S_{R(N_{C_{\cap}})}(\Psi || \Omega) \geq S_{R(N_{C_1})}(\Psi || \Omega) + S_{R(N_{C_2})}(\Psi || \Omega).$$

As proven in [CF20, Section 3.2] the QNEC (24) implies this strong superadditivity for a state Ψ with finite QNEC (24). We can show that the states considered in Section 5.3 saturate the strong superadditivity of relative entropy. Indeed, we apply Proposition 5.3 and have:

$$\begin{aligned}
& S_{R(H(N_{C_{\cup}}))}(\omega \circ \beta_h || \omega) + S_{R(H(N_{C_{\cap}}))}(\omega \circ \beta_h || \omega) - S_{R(H(N_{C_1}))}(\omega \circ \beta_h || \omega) - S_{R(H(N_{C_2}))}(\omega \circ \beta_h || \omega) \\
&= \pi \int_{\mathbb{R}^{D-1}} \left(\int_{C_{\cup}(\mathbf{x}_{\perp})}^{\infty} (x_+ - C_{\cup}(\mathbf{x}_{\perp})) + \int_{C_{\cap}(\mathbf{x}_{\perp})}^{\infty} (x_+ - C_{\cap}(\mathbf{x}_{\perp})) \right. \\
&\quad \left. - \int_{C_1(\mathbf{x}_{\perp})}^{\infty} (x_+ - C_1(\mathbf{x}_{\perp})) - \int_{C_2(\mathbf{x}_{\perp})}^{\infty} (x_+ - C_2(\mathbf{x}_{\perp})) \right) h(x_+, \mathbf{x}_{\perp})^2 dx_+ d\mathbf{x}_{\perp} = 0
\end{aligned}$$

since $C_{\cup}(\mathbf{x}_{\perp}) + C_{\cap}(\mathbf{x}_{\perp}) - C_1(\mathbf{x}_{\perp}) - C_2(\mathbf{x}_{\perp}) = 0$ for all \mathbf{x}_{\perp} .

6 Concluding remarks

Let us close this paper with a few more remarks.

- It is possible to extend the definition of local subspaces on the null plane to measurable functions C . The strategy is to loosen the assumption on "thin test functions" (cf. Section 3.3) in the sense that $\mathbf{x}_\perp \mapsto g(x_+, \mathbf{x}_\perp) \in L^1(\mathbb{R}^{D-1}) \cap L^2(\mathbb{R}^{D-1})$ for almost every x_+ . The real subspaces are covariant with respect to distorted lightlike translations and dilations (cf. (15) and (16)) for measurable functions C . Also in this case T_C and D_C are determined by measurable vector fields of bounded operators since Remark 3.9 obviously extends. It constitutes an extension of the present analysis in the sense that for continuous functions the definitions coincide. Many results such as the decomposition of the subspaces (Proposition 3.11) and modular operators (Theorem 4.3) hold for the resulting subspaces as well. However, isotony (Lemma 3.7) and duality for null-cuts (Theorem 3.12) do not hold for arbitrary measurable functions.
- The modular operator of a direct integral of standard subspaces decomposes into the direct integral of modular operators. Therefore, the modular operator of $H(R)$ from Proposition 3.11 decomposes into the direct integral of $U(1)$ -modular operators of the interval $(C_1(\mathbf{x}_\perp), C_2(\mathbf{x}_\perp))$.
- The fact that there are observables that can be restricted to null plane shows that the minimal localization region in the sense of [Kuc00] can have empty interior.
- In a general Haag-Kastler net, we do not expect that there are sufficiently many observables (e.g. in the sense of the Reeh-Schlieder property) that can be restricted to the null plane. With additional assumptions (including the \mathbb{C} -number commutation relations), it is shown that only free fields can be directly restricted on the null plane [Dri77a], cf. [Wal12, Section 5] [BCFM15, A]. In addition, in the two-dimensional spacetime, there are interacting Haag-Kastler nets [Tan14] where observables on the lightray generate a proper subspace of the Hilbert space from the vacuum [BT15, Section 5.3, Trivial examples]. In these cases, different ideas are required to justify (1).

On the other hand, it has been suggested that there could be bounded operators localized on the null plane in the sense of Haag duality [Sch05]. If so, two-dimensional conformal field theory is hidden on each lightlike fibre [BLM11].

Acknowledgements

We thank Roberto Longo and Aron Wall for inspiring discussions.

VM is supported by Alexander-von-Humboldt Foundation and was supported by the European Research Council Advanced Grant 669240 QUEST until March 2021. BW is supported by the INdAM Doctoral programme and has received funding from the European Union's 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 713485.

We acknowledge the MIUR Excellence Department Project awarded to the Department of Mathematics, University of Rome "Tor Vergata" CUP E83C18000100006 and the University of Rome "Tor Vergata" funding scheme "Beyond Borders" CUP E84I19002200005.

A Decomposable Functional calculus

Assume the same notations as in Section 3.1. We call an (possibly unbounded) operator T on the direct integral of Hilbert spaces $\mathcal{K} = \int_X^\oplus \mathcal{K}_\lambda d\nu(\lambda)$ decomposable when there is a function $\lambda \mapsto T_\lambda$, such that T_λ is an (possibly unbounded) operator on \mathcal{K}_λ , and for each $\xi \in \mathcal{D}(T)$, $\xi(\lambda) \in \mathcal{D}(T_\lambda)$ and $(T\xi)(\lambda) = T_\lambda\xi(\lambda)$ for ν -almost every λ .

Theorem A.1. *Let T be a self-adjoint operator on the direct integral of Hilbert spaces $\mathcal{K} = \int_X^\oplus \mathcal{K}_\lambda d\nu(\lambda)$. The following are equivalent:*

1. *The projection-valued measure E^T associated to T decomposes in the direct integral of projection-valued measures:*

$$E^T = \int_X^\oplus E^{T_\lambda} d\nu(\lambda) =: E \quad (30)$$

for some self-adjoint operators T_λ on \mathcal{K}_λ .

2. *The Borel functional calculus of T decomposes in the direct sum of Borel functional calculi of self-adjoint operators T_λ on \mathcal{K}_λ :*

$$f(T) = \int_X^\oplus f(T_\lambda) d\nu(\lambda).$$

3. *$T = \int_X^\oplus T_\lambda d\nu(\lambda)$ is decomposable. Each T_λ is a self-adjoint operator on \mathcal{K}_λ .*

4. *The one-parameter group of unitaries $U(t) := e^{itT}$ decomposes:*

$$U(t) = \int_X^\oplus U_\lambda(t) d\nu(\lambda),$$

where $U_\lambda(t)$ is a one-parameter group of unitaries on \mathcal{K}_λ .

Proof. 2 \Rightarrow 3,4: clear.

3 \Leftrightarrow 1 \Rightarrow 2: We show that E (cf. (30)) is the spectral measure associated to $\int_X^\oplus T_\lambda d\nu(\lambda)$. It is a projection-valued measure defined by:

$$\Omega \mapsto E(\Omega) := \int_X^\oplus E^{T_\lambda}(\Omega) d\nu(\lambda).$$

That is because the multiplication of decomposable operators is defined fibre-wise. It is clear that each element is a projection and that $E(\emptyset) = 0$, $E(\mathbb{R}) = 1$ and $E(\Omega_1 \cap \Omega_2) = E(\Omega_1)E(\Omega_2)$. For mutually disjoint Ω_n the associated projections are orthogonal. It follows that their sum is again a projection. Hence, we can use dominated convergence, to deduce the convergence $E(\cup^N \Omega_n) \xrightarrow{N \rightarrow \infty} E(\Omega)$ for a countable union $\Omega = \cup \Omega_n$ from the fibre-wise convergence. The latter is true because E^{T_λ} is a spectral measure. The spectral measure E_φ for $\varphi \in \mathcal{K}$ is for a Borel set Ω :

$$E_\varphi(\Omega) = \langle \varphi, E(\Omega)\varphi \rangle = \int_X E_{\varphi_\lambda}^{T_\lambda}(\Omega) d\nu(\lambda).$$

For integration, we denote it by $dE_\varphi(x)$. For a simple function $s = \sum a_i \chi_{\Omega_i}$ on a compact set $\Omega = \cup_i \Omega_i$ it holds:

$$\begin{aligned} \int_{\mathbb{R}} s(x) dE_\varphi(x) &= \sum_i a_i \int_X E_{\varphi_\lambda}^{T_\lambda}(\Omega_i) d\nu(\lambda) \\ &= \int_X \int_{\mathbb{R}} s(x) dE_{\varphi_\lambda}^{T_\lambda}(x) d\nu(\lambda) = \langle \varphi, \left(\int_X^\oplus s(T_\lambda) d\nu(\lambda) \right) \varphi \rangle. \end{aligned}$$

The integral of a general bounded Borel function is the difference of the integrals over the positive and negative part. The integral over a positive bounded Borel function f with respect to dE_φ is defined as the supremum of integral over simple functions s that are locally bounded by f :

$$\int_{\mathbb{R}} f(x) dE_\varphi(x) = \sup_{0 \leq s \leq f} \int_X \int_{\mathbb{R}} s(x) dE_{\varphi_\lambda}^{T_\lambda}(x) d\nu(\lambda) \leq \int_X \sup_{0 \leq s \leq f} \int_{\mathbb{R}} s(x) dE_{\varphi_\lambda}^{T_\lambda}(x) d\nu(\lambda)$$

For a positive, Borel measurable function f , there is a non-decreasing sequence s_n of simple functions converging from below pointwise. We can utilize dominated convergence (spectral measures are finite) to show

$$\lim_{n \rightarrow \infty} \int_X \int_{\mathbb{R}} s_n(x) dE_{\varphi_\lambda}(x) d\nu(\lambda) = \int_X \int_{\mathbb{R}} f(x) dE_{\varphi_\lambda}(x) d\nu(\lambda).$$

Hence, we have for bounded Borel functions:

$$\int_X \int_{\mathbb{R}} f(x) dE_{\varphi_\lambda}^{T_\lambda}(x) d\nu(\lambda) = \int_X f(x) dE_\varphi(x). \quad (31)$$

For each $\varphi \in \mathcal{K}$, the measure E_φ is inner regular and countably additive. The second property was already discussed. The inner regularity follows from the finiteness of E_φ . Hence, the two integrals in (31) define the same bounded, linear functional on the compactly supported, continuous functions. From the uniqueness in the Riesz-Markov theorem (cf. Lemma A.2), we deduce the equality of the measures. This means for each $\varphi \in \mathcal{K}$ and Borel function h :

$$\int_{\mathbb{R}} h(x) dE_\varphi = \int_X \int_{\mathbb{R}} h(x) dE_{\varphi_\lambda}^{T_\lambda} d\nu(\lambda) = \langle \varphi, \left(\int_X^\oplus h(T_\lambda) d\nu(\lambda) \right) \varphi \rangle.$$

In the usual notation, this amounts to:

$$\int_{\mathbb{R}} h(x) dE(x) = \int_X^\oplus h(T_\lambda) d\nu(\lambda).$$

The domain of the operator $h(T)$ is:

$$\begin{aligned} \mathcal{D}(h(T)) &= \{ \varphi \in \mathcal{K}; \int_{\mathbb{R}} |f(k)|^2 dE_\varphi^T(k) < \infty \} \\ &= \{ \varphi \in \mathcal{K}; \int_X \int_{\mathbb{R}} |f(k)|^2 dE_{\varphi_\lambda}^{T_\lambda}(k) d\nu(\lambda) < \infty \} \\ &= \{ \varphi \in \mathcal{K}; \int_X \|h(T_\lambda)\varphi(\lambda)\|^2 d\nu(\lambda) < \infty \} = \int_X^\oplus \mathcal{D}(h(T_\lambda)) d\nu(\lambda). \end{aligned}$$

For a closed operator A on a Hilbert space \mathcal{H} its domain $\mathcal{D}(A)$ is a Hilbert space w.r.t. the Graph inner product:

$$\langle \xi, \eta \rangle_A := \langle \xi, \eta \rangle_{\mathcal{H}} + \langle A\xi, A\eta \rangle_{\mathcal{H}}.$$

Since T_λ is self-adjoint for (ν -almost) every λ , the domain of $h(T_\lambda)$ is a Hilbert space w.r.t. the Graph inner product for almost every λ . The domain $\mathcal{D}(h(T))$ embeds in \mathcal{K} trivially. If T is decomposable, then we have for all vectors $\varphi \in \mathcal{K}$:

$$\int_{\mathbb{R}} \lambda dE_\varphi^T(\lambda) = \int_X \int_{\mathbb{R}} \lambda dE_{\varphi_\lambda}^{T_\lambda} d\nu(\lambda).$$

4 \Rightarrow 1: Let $\xi, \eta \in \mathcal{K}$ and $g \in \mathcal{S}(\mathbb{R})$. Then by Fubini's theorem the operator $g(T)$ decomposes:

$$\begin{aligned} \langle \xi, g(T)\eta \rangle &= \int_{\mathbb{R}} \tilde{g}(t) \langle \xi, U(t)\eta \rangle dt \\ &= \int_X \langle \xi(\lambda), \left(\int_{\mathbb{R}} \tilde{g}(t) U_\lambda(t) dt \right) \eta(\lambda) \rangle d\nu(\lambda) \\ &= \langle \xi, \left(\int_X^\oplus g(T_\lambda) d\nu(\lambda) \right) \eta \rangle. \end{aligned}$$

Every characteristic function χ_S of a set S with finite measure can be expressed as the pointwise limit of a sequence $g_n \subset \mathcal{S}(\mathbb{R})$ of uniformly bounded functions. In the functional calculus this amounts to the strong convergence:

$$g_n(T) \xrightarrow{n \rightarrow \infty} \chi_S(T) = E^T(S),$$

where $E^T(S)$ denotes the spectral projection of T with respect to the set S .

Since g_n is uniformly bounded, we can apply the theorem of dominated convergence:

$$\begin{aligned} \langle \xi, E_S^T \eta \rangle &= \lim_{n \rightarrow \infty} \int_X \langle \xi(\lambda), g_n(T_\lambda) \eta(\lambda) \rangle d\nu(\lambda) \\ &= \int_X \langle \xi(\lambda), \chi_S(T_\lambda) \eta(\lambda) \rangle d\nu(\lambda) \\ &= \langle \xi, \left(\int_X^\oplus E^{T_\lambda}(S) d\nu(\lambda) \right) \eta \rangle. \end{aligned}$$

Hence, the spectral measure E_φ^T associated to any vector $\varphi \in \mathcal{K}$ decomposes in the following sense for any bounded Borel function f :

$$\langle \varphi, f(T)\varphi \rangle = \int_{\mathbb{R}} f(k) dE_\varphi^T(k) \tag{32}$$

$$= \int_X \int_{\mathbb{R}} f(k) dE_{\varphi_\lambda}^{T_\lambda}(k) = \langle \varphi, \left(\int_X^\oplus f(T_\lambda) d\nu(\lambda) \right) \varphi \rangle. \tag{33}$$

The two expressions (32) and (33) define the same linear functional on the continuous, compactly supported functions. By the Riesz-Markov theorem such a functional is determined by a unique countably additive, inner regular measure on \mathbb{R} . Since both measures in (32) and (33) are countably additive and inner regular, they coincide. \square

Lemma A.2. [*Riesz-Markov theorem*] Let X be a locally compact Hausdorff space and Ψ a bounded linear functional on $\mathcal{C}_c(X)$, then there exists a unique inner regular and countably additive measure μ on X such that:

$$\Psi(f) = \int_X f d\mu.$$

References

- [Ara63] Huzihiro Araki. A lattice of von Neumann algebras associated with the quantum theory of a free Bose field. *J. Mathematical Phys.*, 4:1343–1362, 1963. <https://doi.org/10.1063/1.1703912>.
- [AW66] Huzihiro Araki and E. J. Woods. Complete Boolean algebras of type I factors. *Publ. Res. Inst. Math. Sci. Ser. A*, 2:157–242, 1966. <https://doi.org/10.2977/prims/1195195888>.
- [AZ05] Huzihiro Araki and László Zsidó. Extension of the structure theorem of Borchers and its application to half-sided modular inclusions. *Rev. Math. Phys.*, 17(5):491–543, 2005. <https://arxiv.org/abs/math/0412061>.
- [BCFM15] Raphael Bousso, Horacio Casini, Zachary Fisher, and Juan Maldacena. Entropy on a null surface for interacting quantum field theories and the Bousso bound. *Phys. Rev. D*, 91(8):084030, 17, 2015. <https://arxiv.org/abs/1406.4545>.
- [BFK⁺16] Raphael Bousso, Zachary Fisher, Jason Koeller, Stefan Leichenauer, and Aron C. Wall. Proof of the quantum null energy condition. *Physical Review D*, 93(2), Jan 2016. <https://arxiv.org/abs/1509.02542>.
- [BFKW19] Srivatsan Balakrishnan, Thomas Faulkner, Zuhair U. Khandker, and Huajia Wang. A general proof of the quantum null energy condition. *Journal of High Energy Physics*, 2019(9), Sep 2019. <https://arxiv.org/abs/1706.09432>.
- [BFLW16] Raphael Bousso, Zachary Fisher, Stefan Leichenauer, and Aron C. Wall. Quantum focusing conjecture. *Physical Review D*, 93(6), Mar 2016. <https://arxiv.org/abs/1506.02669>.
- [BGL02] R. Brunetti, D. Guido, and R. Longo. Modular localization and Wigner particles. *Rev. Math. Phys.*, 14(7-8):759–785, 2002. <https://arxiv.org/abs/math-ph/0203021>.
- [BLM11] Henning Bostelmann, Gandalf Lechner, and Gerardo Morsella. Scaling limits of integrable quantum field theories. *Rev. Math. Phys.*, 23(10):1115–1156, 2011. <https://arxiv.org/abs/1105.2781>.
- [BMT88] Detlev Buchholz, Gerhard Mack, and Ivan Todorov. The current algebra on the circle as a germ of local field theories. *Nuclear Phys. B Proc. Suppl.*, 5B:20–56, 1988. <https://www.researchgate.net/publication/222585851>.
- [Bor92] H.-J. Borchers. The CPT-theorem in two-dimensional theories of local observables. *Comm. Math. Phys.*, 143(2):315–332, 1992. <http://projecteuclid.org/euclid.cmp/1104248958>.

- [BT15] Marcel Bischoff and Yoh Tanimoto. Integrable QFT and Longo-Witten endomorphisms. *Ann. Henri Poincaré*, 16(2):569–608, 2015. <https://arxiv.org/abs/1305.2171>.
- [BW76] Joseph J. Bisognano and Eyvind H. Wichmann. On the duality condition for quantum fields. *J. Mathematical Phys.*, 17(3):303–321, 1976. <https://doi.org/10.1063/1.522898>.
- [Cas08] H. Casini. Relative entropy and the Bekenstein bound. *Classical Quantum Gravity*, 25(20):205021, 12, 2008. <https://arxiv.org/abs/0804.2182>.
- [CF20] Fikret Ceyhan and Thomas Faulkner. Recovering the QNEC from the ANEC. *Comm. Math. Phys.*, 377(2):999–1045, 2020. <https://arxiv.org/abs/1812.04683>.
- [CLR20] Fabio Ciolli, Roberto Longo, and Giuseppe Ruzzi. The Information in a Wave. *Comm. Math. Phys.*, 379(3):979–1000, 2020. <https://arxiv.org/abs/1703.10656>.
- [CTT17a] Horacio Casini, Eduardo Testé, and Gonzalo Torroba. Markov property of the conformal field theory vacuum and the a theorem. *Phys. Rev. Lett.*, 118(26):261602, 5, 2017. <https://arxiv.org/abs/1704.01870>.
- [CTT17b] Horacio Casini, Eduardo Testé, and Gonzalo Torroba. Modular Hamiltonians on the null plane and the Markov property of the vacuum state. *J. Phys. A*, 50(36):364001, 29, 2017. <https://arxiv.org/abs/1703.10656>.
- [Dix81] Jacques Dixmier. *von Neumann algebras*, volume 27 of *North-Holland Mathematical Library*. North-Holland Publishing Co., Amsterdam, 1981. <https://books.google.com/books?id=8xSoAAAAIAAJ>.
- [Dri77a] W. Driessler. On the structure of fields and algebras on null planes. I. Local algebras. *Acta Phys. Austriaca*, 46(2):63–96, 1976/77.
- [Dri77b] W. Driessler. On the structure of fields and algebras on null planes. II. Field structure. *Acta Phys. Austriaca*, 46(3):163–196, 1976/77.
- [Few12] C. Fewster. Lectures on quantum energy inequalities. 2012. <https://arxiv.org/abs/1208.5399>.
- [FH05] Christopher J. Fewster and Stefan Hollands. Quantum energy inequalities in two-dimensional conformal field theory. *Rev. Math. Phys.*, 17(5):577–612, 2005. <https://arxiv.org/abs/math-ph/0412028>.
- [Flo98] Martin Florig. On Borchers’ theorem. *Lett. Math. Phys.*, 46(4):289–293, 1998. <https://dx.doi.org/10.1023/A:1007546507392>.
- [FLPW16] Thomas Faulkner, Robert G. Leigh, Onkar Parrikar, and Huajia Wang. Modular Hamiltonians for deformed half-spaces and the averaged null energy condition. *Journal of High Energy Physics*, 2016(9), Sep 2016. <https://arxiv.org/abs/1605.08072>.
- [FR03] Christopher J. Fewster and Thomas A. Roman. Null energy conditions in quantum field theory. *Phys. Rev. D (3)*, 67(4):044003, 11, 2003. <https://arxiv.org/abs/gr-qc/0209036>.

- [GLRV01] D. Guido, R. Longo, J. E. Roberts, and R. Verch. Charged sectors, spin and statistics in quantum field theory on curved spacetimes. *Rev. Math. Phys.*, 13(2):125–198, 2001. <https://arxiv.org/abs/math-ph/9906019>.
- [Hol20] Stefan Hollands. Relative entropy for coherent states in chiral CFT. *Lett. Math. Phys.*, 110(4):713–733, 2020. <https://arxiv.org/abs/1903.07508>.
- [KLLSM18] Jason Koeller, Stefan Leichenauer, Adam Levine, and Arvin Shahbazi-Moghaddam. Local modular Hamiltonians from the quantum null energy condition. *Phys. Rev. D*, 97(6):065011, 6, 2018. <https://arxiv.org/abs/1702.00412>.
- [KR97] Richard V. Kadison and John R. Ringrose. *Fundamentals of the theory of operator algebras. Vol. II*, volume 16 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 1997. <https://books.google.com/books?id=6eorDAAAQBAJ>.
- [Kuc00] Bernd Kuckert. Localization regions of local observables. *Comm. Math. Phys.*, 215(1):197–216, 2000. <https://arxiv.org/abs/math-ph/0002040>.
- [LLSM18] Stefan Leichenauer, Adam Levine, and Arvin Shahbazi-Moghaddam. Energy density from second shape variations of the von neumann entropy. *Physical Review D*, 98(8), Oct 2018. <https://arxiv.org/abs/1802.02584>.
- [LMR16] Roberto Longo, Vincenzo Morinelli, and Karl-Henning Rehren. Where infinite spin particles are localizable. *Comm. Math. Phys.*, 345(2):587–614, 2016. <https://arxiv.org/abs/1505.01759>.
- [Lon08] Roberto Longo. Real Hilbert subspaces, modular theory, $SL(2, \mathbf{R})$ and CFT. In *Von Neumann algebras in Sibiu: Conference Proceedings*, pages 33–91. Theta, Bucharest, 2008. https://www.mat.uniroma2.it/longo/Lecture-Notes_files/LN-Part1.pdf.
- [Lon19] Roberto Longo. Entropy of coherent excitations. *Lett. Math. Phys.*, 109(12):2587–2600, 2019. <https://arxiv.org/abs/1901.02366>.
- [Lon20] Roberto Longo. Entropy distribution of localised states. *Comm. Math. Phys.*, 373(2):473–505, 2020. <https://arxiv.org/abs/1809.03358>.
- [LRT78] Pen Leyland, John Roberts, and Daniel Testard. Duality for quantum free fields. *unpublished manuscript, Marseille*, 1978.
- [LX18] Roberto Longo and Feng Xu. Relative entropy in CFT. *Adv. Math.*, 337:139–170, 2018. <https://arxiv.org/abs/1712.07283>.
- [Mor18] Vincenzo Morinelli. The Bisognano-Wichmann property on nets of standard subspaces, some sufficient conditions. *Ann. Henri Poincaré*, 19(3):937–958, 2018. <https://arxiv.org/abs/1703.06831>.
- [MT18] Vincenzo Morinelli and Yoh Tanimoto. Scale and Möbius covariance in two-dimensional Haag-Kastler net. 2018. <https://arxiv.org/abs/1807.04707>.
- [Nap71] Kazimierz Napiórkowski. Continuous tensor products of Hilbert spaces and product operators. *Studia Math.*, 39:307–327. (errata insert), 1971. <http://matwbn.icm.edu.pl/ksiazki/sm/sm39/sm39122.pdf>.

- [OP04] M. Ohya and D. Petz. *Quantum Entropy and Its Use*. Theoretical and Mathematical Physics. Springer Berlin Heidelberg, 2004. <https://books.google.co.jp/books?id=r2u11NVyESQC>.
- [Pan20] Lorenzo Panebianco. A formula for the relative entropy in chiral CFT. *Lett. Math. Phys.*, 110(9):2363–2381, 2020. <https://arxiv.org/abs/1911.10136>.
- [Sch05] Bert Schroer. Constructive proposals for QFT based on the crossing property and on lightfront holography. *Ann. Physics*, 319(1):48–91, 2005. <https://arxiv.org/abs/hep-th/0406016>.
- [SS72] S. Schlieder and E. Seiler. Some remarks on the “null plane development” of a relativistic quantum field theory. *Comm. Math. Phys.*, 25:62–72, 1972. <http://projecteuclid.org/euclid.cmp/1103857838>.
- [Tak03] M. Takesaki. *Theory of operator algebras. III*, volume 127 of *Encyclopaedia of Mathematical Sciences*. Springer-Verlag, Berlin, 2003. <https://books.google.com/books?id=MGgHL15Ggg4C>.
- [Tan14] Yoh Tanimoto. Construction of two-dimensional quantum field models through Longo-Witten endomorphisms. *Forum Math. Sigma*, 2:e7, 31, 2014. <https://arxiv.org/abs/1301.6090>.
- [Ull04] Peter Ullrich. On the restriction of quantum fields to a lightlike surface. *J. Math. Phys.*, 45(8):3109–3145, 2004. <https://doi.org/10.1063/1.1765746>.
- [Var85] V.S. Varadarajan. *Geometry of quantum theory*. Springer-Verlag, New York, second edition edition, 1985. <https://books.google.com/books?id=AbbMLuXC9tsC>.
- [Ver00] Rainer Verch. The averaged null energy condition for general quantum field theories in two dimensions. *J. Math. Phys.*, 41(1):206–217, 2000. <https://arxiv.org/abs/math-ph/9904036>.
- [Wal12] Aron C. Wall. Proof of the generalized second law for rapidly changing fields and arbitrary horizon slices. *Phys. Rev. D*, 85:104049, May 2012. <https://arxiv.org/abs/1105.3445>.
- [Wie93] Hans-Werner Wiesbrock. Half-sided modular inclusions of von-Neumann-algebras. *Comm. Math. Phys.*, 157(1):83–92, 1993. <https://projecteuclid.org/euclid.cmp/1104253848>.